

Numerical Analysis & Scientific Computing II

Lesson 4

Numerical Solution of PDE

4.1 BVP for 2nd Order Elliptic PDE



Akash Anand
MATH, IIT KANPUR

Numerical Analysis & Scientific Computing II

Lesson 4

Numerical Solution of PDE

4.1 BVP for 2nd Order Elliptic PDE **- Finite Difference Method**



Akash Anand
MATH, IIT KANPUR

Numerical Methods for PDE: 2nd Order Elliptic PDE



A natural generalization to the two-point BVP

$$\begin{aligned}u'' &= f(t), & a < t < b, \\u(a) &= 0, & u(b) = 0,\end{aligned}$$

to two dimensions is

$$\begin{aligned}\Delta u &:= u_{x_1x_1} + u_{x_2x_2} = f, & \text{in } \Omega, \\u &= g, & \text{on } \Gamma.\end{aligned}$$

Numerical Methods for PDE: 2nd Order Elliptic PDE


A natural generalization to the two-point BVP

$$\begin{aligned} u'' &= f(t), & a < t < b, \\ u(a) &= 0, & u(b) = 0, \end{aligned}$$

to two dimensions is

$$\begin{aligned} \Delta u &:= u_{x_1 x_1} + u_{x_2 x_2} = f, & \text{in } \Omega, \\ u &= g, & \text{on } \Gamma. \end{aligned}$$

For simplicity, we will first consider a very simple domain $\Omega = (0,1) \times (0,1)$.



Numerical Methods for PDE: 2nd Order Elliptic PDE

A natural generalization to the two-point BVP

$$\begin{aligned}u'' &= f(t), & a < t < b, \\u(a) &= 0, & u(b) = 0,\end{aligned}$$

to two dimensions is


$$\begin{aligned}\Delta u &:= u_{x_1x_1} + u_{x_2x_2} = f, & \text{in } \Omega, \\u &= g, & \text{on } \Gamma.\end{aligned}$$

For simplicity, we will first consider a very simple domain $\Omega = (0,1) \times (0,1)$.

Let N be a positive integer and set $h = 1/N$. Consider the mesh in \mathbb{R}^2

$$\mathbb{R}_h^2 = \{(mh, nh) : m, n \in \mathbb{Z}\}.$$

Let $\Omega_h = \Omega \cap \mathbb{R}_h^2$, the set of interior mesh points.



Numerical Methods for PDE: 2nd Order Elliptic PDE

A natural generalization to the two-point BVP

$$\begin{aligned}u'' &= f(t), & a < t < b, \\u(a) &= 0, & u(b) = 0,\end{aligned}$$

to two dimensions is


$$\begin{aligned}\Delta u &:= u_{x_1x_1} + u_{x_2x_2} = f, & \text{in } \Omega, \\u &= g, & \text{on } \Gamma.\end{aligned}$$

For simplicity, we will first consider a very simple domain $\Omega = (0,1) \times (0,1)$.

Let N be a positive integer and set $h = 1/N$. Consider the mesh in \mathbb{R}^2

$$\mathbb{R}_h^2 = \{(mh, nh) : m, n \in \mathbb{Z}\}.$$

Let $\Omega_h = \Omega \cap \mathbb{R}_h^2$, the set of interior mesh points.



Numerical Methods for PDE: 2nd Order Elliptic PDE

A natural generalization to the two-point BVP

$$\begin{aligned} u'' &= f(t), & a < t < b, \\ u(a) &= 0, & u(b) &= 0, \end{aligned}$$

to two dimensions is

$$\begin{aligned} \Delta u &:= u_{x_1 x_1} + u_{x_2 x_2} = f, & \text{in } \Omega, \\ u &= g, & \text{on } \Gamma. \end{aligned}$$


For simplicity, we will first consider a very simple domain $\Omega = (0,1) \times (0,1)$.

Let N be a positive integer and set $h = 1/N$. Consider the mesh in \mathbb{R}^2

$$\mathbb{R}_h^2 = \{(mh, nh) : m, n \in \mathbb{Z}\}.$$

Let $\Omega_h = \Omega \cap \mathbb{R}_h^2$, the set of interior mesh points.

Note that for each $x \in \mathbb{R}_h^2$ has a set of four nearest neighbors in \mathbb{R}_h^2 , one each to the left, right, above and below. We define Γ_h as the set of mesh points in \mathbb{R}_h^2 which is not in Ω_h but has a nearest neighbor in Ω_h .



Numerical Methods for PDE: 2nd Order Elliptic PDE

A natural generalization to the two-point BVP

$$\begin{aligned} u'' &= f(t), & a < t < b, \\ u(a) &= 0, & u(b) &= 0, \end{aligned}$$

to two dimensions is

$$\begin{aligned} \Delta u &:= u_{x_1 x_1} + u_{x_2 x_2} = f, & \text{in } \Omega, \\ u &= g, & \text{on } \Gamma. \end{aligned}$$


For simplicity, we will first consider a very simple domain $\Omega = (0,1) \times (0,1)$.

Let N be a positive integer and set $h = 1/N$. Consider the mesh in \mathbb{R}^2

$$\mathbb{R}_h^2 = \{(mh, nh) : m, n \in \mathbb{Z}\}.$$

Let $\Omega_h = \Omega \cap \mathbb{R}_h^2$, the set of interior mesh points.

Note that for each $x \in \mathbb{R}_h^2$ has a set of four nearest neighbors in \mathbb{R}_h^2 , one each to the left, right, above and below. We define Γ_h as the set of mesh points in \mathbb{R}_h^2 which is not in Ω_h but has a nearest neighbor in Ω_h .



Numerical Methods for PDE: 2nd Order Elliptic PDE

A natural generalization to the two-point BVP

$$\begin{aligned} u'' &= f(t), & a < t < b, \\ u(a) &= 0, & u(b) &= 0, \end{aligned}$$

to two dimensions is

$$\begin{aligned} \Delta u &:= u_{x_1 x_1} + u_{x_2 x_2} = f, & \text{in } \Omega, \\ u &= g, & \text{on } \Gamma. \end{aligned}$$

For simplicity, we will first consider a very simple domain $\Omega = (0,1) \times (0,1)$.


Let N be a positive integer and set $h = 1/N$. Consider the mesh in \mathbb{R}^2

$$\mathbb{R}_h^2 = \{(mh, nh) : m, n \in \mathbb{Z}\}.$$

Let $\Omega_h = \Omega \cap \mathbb{R}_h^2$, the set of interior mesh points.

Note that for each $x \in \mathbb{R}_h^2$ has a set of four nearest neighbors in \mathbb{R}_h^2 , one each to the left, right, above and below. We define Γ_h as the set of mesh points in \mathbb{R}_h^2 which is not in Ω_h but has a nearest neighbor in Ω_h .

Also let $\bar{\Omega}_h = \Omega_h \cup \Gamma_h$.



Numerical Methods for PDE: 2nd Order Elliptic PDE

To discretize

$$\Delta u = f, \quad \text{in } \Omega,$$

$$u = g, \quad \text{on } \Gamma,$$


we seek a function $u_h: \overline{\Omega_h} \rightarrow \mathbb{R}$ satisfying

$$\Delta_h u_h = f, \quad \text{on } \Omega_h,$$

$$u_h = g, \quad \text{on } \Gamma_h.$$

where, writing $v_{m,n} = v(mh, nh)$, we have the **5-point Laplacian**

$$\Delta_h v(mh, nh) = \frac{v_{m+1,n} - 2v_{m,n} + v_{m-1,n}}{h^2} + \frac{v_{m,n+1} - 2v_{m,n} + v_{m,n-1}}{h^2}$$



Numerical Methods for PDE: 2nd Order Elliptic PDE

To discretize

$$\Delta u = f, \quad \text{in } \Omega,$$

$$u = g, \quad \text{on } \Gamma,$$


we seek a function $u_h: \overline{\Omega_h} \rightarrow \mathbb{R}$ satisfying

$$\Delta_h u_h = f, \quad \text{on } \Omega_h,$$

$$u_h = g, \quad \text{on } \Gamma_h.$$

where, writing $v_{m,n} = v(mh, nh)$, we have the **5-point Laplacian**

$$\begin{aligned} \Delta_h v(mh, nh) &= \frac{v_{m+1,n} - 2v_{m,n} + v_{m-1,n}}{h^2} + \frac{v_{m,n+1} - 2v_{m,n} + v_{m,n-1}}{h^2} \\ &= \frac{v_{m+1,n} + v_{m-1,n} + v_{m,n+1} + v_{m,n-1} - 4v_{m,n}}{h^2}. \end{aligned}$$



Numerical Methods for PDE: 2nd Order Elliptic PDE

To discretize

$$\Delta u = f, \quad \text{in } \Omega,$$

$$u = g, \quad \text{on } \Gamma,$$

we seek a function $u_h: \overline{\Omega}_h \rightarrow \mathbb{R}$ satisfying

$$\Delta_h u_h = f, \quad \text{on } \Omega_h,$$

$$u_h = g, \quad \text{on } \Gamma_h.$$


where, writing $v_{m,n} = v(mh, nh)$, we have the **5-point Laplacian**

$$\begin{aligned} \Delta_h v(mh, nh) &= \frac{v_{m+1,n} - 2v_{m,n} + v_{m-1,n}}{h^2} + \frac{v_{m,n+1} - 2v_{m,n} + v_{m,n-1}}{h^2} \\ &= \frac{v_{m+1,n} + v_{m-1,n} + v_{m,n+1} + v_{m,n-1} - 4v_{m,n}}{h^2}. \end{aligned}$$

From the error estimate in one-dimensional case, we can easily get that for $v \in C^4(\overline{\Omega})$,

$$\Delta_h v(mh, nh) - \Delta v(mh, nh) = \frac{h^2}{12} \left[\frac{\partial^4 v}{\partial x_1^4}(\xi, nh) + \frac{\partial^4 v}{\partial x_2^4}(mh, \eta) \right]$$

for some ξ, η .



Numerical Methods for PDE: 2nd Order Elliptic PDE

Theorem

If $v \in C^2(\overline{\Omega})$, then

$$\lim_{h \rightarrow 0} \|\Delta_h v - \Delta v\|_{\infty, \Omega_h} = 0.$$

If $v \in C^4(\overline{\Omega})$, then

$$\|\Delta_h v - \Delta v\|_{\infty, \Omega_h} \leq \frac{h^2}{6} M_4,$$

where

$$M_4 = \max \left\{ \left\| \frac{\partial^4 v}{\partial x_1^4} \right\|_{\infty, \overline{\Omega}}, \left\| \frac{\partial^4 v}{\partial x_2^4} \right\|_{\infty, \overline{\Omega}} \right\}.$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

Theorem

If $v \in C^2(\overline{\Omega})$, then

$$\lim_{h \rightarrow 0} \|\Delta_h v - \Delta v\|_{\infty, \Omega_h} = 0.$$

If $v \in C^4(\overline{\Omega})$, then

$$\|\Delta_h v - \Delta v\|_{\infty, \Omega_h} \leq \frac{h^2}{6} M_4,$$

where

$$M_4 = \max \left\{ \left\| \frac{\partial^4 v}{\partial x_1^4} \right\|_{\infty, \overline{\Omega}}, \left\| \frac{\partial^4 v}{\partial x_2^4} \right\|_{\infty, \overline{\Omega}} \right\}.$$

Note that the discrete PDE $\Delta_h u_h = f$, on Ω_h is a system of $(N - 1)^2$ equations in $(N - 1)^2$ unknowns.

Numerical Methods for PDE: 2nd Order Elliptic PDE

Theorem

If $v \in C^2(\overline{\Omega})$, then

$$\lim_{h \rightarrow 0} \|\Delta_h v - \Delta v\|_{\infty, \Omega_h} = 0.$$

If $v \in C^4(\overline{\Omega})$, then

$$\|\Delta_h v - \Delta v\|_{\infty, \Omega_h} \leq \frac{h^2}{6} M_4,$$

where

$$M_4 = \max \left\{ \left\| \frac{\partial^4 v}{\partial x_1^4} \right\|_{\infty, \overline{\Omega}}, \left\| \frac{\partial^4 v}{\partial x_2^4} \right\|_{\infty, \overline{\Omega}} \right\}.$$

Note that the discrete PDE $\Delta_h u_h = f$, on Ω_h is a system of $(N-1)^2$ equations in $(N-1)^2$ unknowns.

Theorem (Discrete Maximum Principle)

Let v be a function on $\overline{\Omega}_h$ satisfying $\Delta_h v \geq 0$ on Ω_h . Then $\max_{\Omega_h} v \leq \max_{\Gamma_h} v$. Equality holds if and only if v is constant.

Numerical Methods for PDE: 2nd Order Elliptic PDE

Theorem

If $v \in C^2(\overline{\Omega})$, then

$$\lim_{h \rightarrow 0} \|\Delta_h v - \Delta v\|_{\infty, \Omega_h} = 0.$$

If $v \in C^4(\overline{\Omega})$, then

$$\|\Delta_h v - \Delta v\|_{\infty, \Omega_h} \leq \frac{h^2}{6} M_4,$$

where

$$M_4 = \max \left\{ \left\| \frac{\partial^4 v}{\partial x_1^4} \right\|_{\infty, \overline{\Omega}}, \left\| \frac{\partial^4 v}{\partial x_2^4} \right\|_{\infty, \overline{\Omega}} \right\}.$$

Note that the discrete PDE $\Delta_h u_h = f$, on Ω_h is a system of $(N - 1)^2$ equations in $(N - 1)^2$ unknowns.

Theorem (Discrete Maximum Principle)

Let v be a function on $\overline{\Omega}_h$ satisfying $\Delta_h v \geq 0$ on Ω_h . Then $\max_{\Omega_h} v \leq \max_{\Gamma_h} v$. Equality holds if and only if v is constant.

Proof: Exercise. **HINT:** Use $4v(x_0) = \sum_{i=1}^4 v(x_i) - h^2 \Delta_h v(x_0)$ where x_1, x_2, x_3, x_4 are neighbors of x_0 .

Numerical Methods for PDE: 2nd Order Elliptic PDE



Theorem (Discrete Minimum Principle)

Let v be a function on $\overline{\Omega}_h$ satisfying $\Delta_h v \leq 0$ on Ω_h . Then $\min_{\Omega_h} v \geq \min_{\Gamma_h} v$. Equality holds if and only if v is constant.



Theorem (Discrete Minimum Principle)

Let v be a function on $\overline{\Omega}_h$ satisfying $\Delta_h v \leq 0$ on Ω_h . Then $\min_{\Omega_h} v \geq \min_{\Gamma_h} v$. Equality holds if and only if v is constant.

Proof: Exercise.



Numerical Methods for PDE: 2nd Order Elliptic PDE

Theorem (Discrete Minimum Principle)

Let v be a function on $\overline{\Omega}_h$ satisfying $\Delta_h v \leq 0$ on Ω_h . Then $\min_{\Omega_h} v \geq \min_{\Gamma_h} v$. Equality holds if and only if v is constant.

Proof: Exercise.

Theorem

There is a unique solution to the discrete BVP

$$\begin{aligned} \Delta_h u_h &= f, & \text{on } \Omega_h, \\ u_h &= g, & \text{on } \Gamma_h. \end{aligned}$$



Numerical Methods for PDE: 2nd Order Elliptic PDE

Theorem (Discrete Minimum Principle)

Let v be a function on $\overline{\Omega}_h$ satisfying $\Delta_h v \leq 0$ on Ω_h . Then $\min_{\Omega_h} v \geq \min_{\Gamma_h} v$. Equality holds if and only if v is constant.

Proof: Exercise.

Theorem

There is a unique solution to the discrete BVP

$$\begin{aligned} \Delta_h u_h &= f, & \text{on } \Omega_h, \\ u_h &= g, & \text{on } \Gamma_h. \end{aligned}$$

Proof: Exercise.

Numerical Methods for PDE: 2nd Order Elliptic PDE

Theorem (Discrete Minimum Principle)

Let v be a function on $\bar{\Omega}_h$ satisfying $\Delta_h v \leq 0$ on Ω_h . Then $\min_{\Omega_h} v \geq \min_{\Gamma_h} v$. Equality holds if and only if v is constant.

Proof: Exercise.

Theorem

There is a unique solution to the discrete BVP

$$\begin{aligned} \Delta_h u_h &= f, & \text{on } \Omega_h, \\ u_h &= g, & \text{on } \Gamma_h. \end{aligned}$$

Proof: Exercise.

Theorem

The solution u_h to

$$\begin{aligned} \Delta_h u_h &= f, & \text{on } \Omega_h, \\ u_h &= g, & \text{on } \Gamma_h. \end{aligned}$$

satisfies

$$\|u_h\|_{\infty, \bar{\Omega}_h} \leq \frac{1}{8} \|f\|_{\infty, \Omega_h} + \|g\|_{\infty, \Gamma_h}.$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

Theorem (Discrete Minimum Principle)

Let v be a function on $\bar{\Omega}_h$ satisfying $\Delta_h v \leq 0$ on Ω_h . Then $\min_{\Omega_h} v \geq \min_{\Gamma_h} v$. Equality holds if and only if v is constant.

Proof: Exercise.

Theorem

There is a unique solution to the discrete BVP

$$\begin{aligned} \Delta_h u_h &= f, & \text{on } \Omega_h, \\ u_h &= g, & \text{on } \Gamma_h. \end{aligned}$$

Proof: Exercise.

Theorem

The solution u_h to

$$\begin{aligned} \Delta_h u_h &= f, & \text{on } \Omega_h, \\ u_h &= g, & \text{on } \Gamma_h. \end{aligned}$$

satisfies

$$\|u_h\|_{\infty, \bar{\Omega}_h} \leq \frac{1}{8} \|f\|_{\infty, \Omega_h} + \|g\|_{\infty, \Gamma_h}.$$

Proof: Exercise. **HINT:** Use $w(x_1, x_2) = [(x_1 - 1/2)^2 + (x_2 - 1/2)^2]/4$.

Numerical Methods for PDE: 2nd Order Elliptic PDE

Theorem

Let u be the solution to

$$\begin{aligned}\Delta u &= f, & \text{in } \Omega, \\ u &= g, & \text{on } \Gamma,\end{aligned}$$

and u_h be the solution to the corresponding discrete problem

$$\begin{aligned}\Delta_h u_h &= f, & \text{on } \Omega_h, \\ u_h &= g, & \text{on } \Gamma_h.\end{aligned}$$

Then,

$$\|u_h - u\|_{\infty, \bar{\Omega}_h} \leq \frac{1}{8} \|\Delta u - \Delta_h u\|_{\infty, \bar{\Omega}_h}.$$

Proof: Exercise.

Corollary

If $u \in C^2(\bar{\Omega}_h)$, then

$$\lim_{h \rightarrow 0} \|u_h - u\|_{\infty, \bar{\Omega}_h} = 0.$$

If $u \in C^4(\bar{\Omega}_h)$, then

$$\|u_h - u\|_{\infty, \bar{\Omega}_h} \leq \frac{h^2}{48} M_4, \quad M_4 = \max \left\{ \left\| \frac{\partial^4 v}{\partial x_1^4} \right\|_{\infty, \bar{\Omega}}, \left\| \frac{\partial^4 v}{\partial x_2^4} \right\|_{\infty, \bar{\Omega}} \right\}.$$