

Lesson 4

Numerical Solution of PDE

4.1 BVP for 2nd Order Elliptic PDE

- Finite Difference Method

- More Stability Analysis – Fourier Analysis



Numerical Methods for PDE: 2nd Order Elliptic PDE



Stability Analysis using Fourier Analysis



Numerical Methods for PDE: 2nd Order Elliptic PDE

Stability Analysis using Fourier Analysis

First we consider the one dimensional case. With $h = 1/N$, let $I_h = \{h, 2h, \dots, (N-1)h\}$ and let

$$L(I_h) = \{u: \bar{I}_h \rightarrow \mathbb{R} : u(0) = 0, u(1) = 0\}.$$

Clearly, $L(I_h)$ is isomorphic to \mathbb{R}^{N-1} .

Numerical Methods for PDE: 2nd Order Elliptic PDE

Stability Analysis using Fourier Analysis

First we consider the one dimensional case. With $h = 1/N$, let $I_h = \{h, 2h, \dots, (N-1)h\}$ and let

$$L(I_h) = \{u: \bar{I}_h \rightarrow \mathbb{R} : u(0) = 0, u(1) = 0\}.$$

Clearly, $L(I_h)$ is isomorphic to \mathbb{R}^{N-1} . On $L(I_h)$, we define the inner product

$$\langle u, v \rangle_h = h \sum_{k=1}^{N-1} u(kh)v(kh)$$

with the corresponding norm $\|v\|_h$.

Numerical Methods for PDE: 2nd Order Elliptic PDE

Stability Analysis using Fourier Analysis

First we consider the one dimensional case. With $h = 1/N$, let $I_h = \{h, 2h, \dots, (N-1)h\}$ and let

$$L(I_h) = \{u: \bar{I}_h \rightarrow \mathbb{R} : u(0) = 0, u(1) = 0\}.$$

Clearly, $L(I_h)$ is isomorphic to \mathbb{R}^{N-1} . On $L(I_h)$, we define the inner product

$$\langle u, v \rangle_h = h \sum_{k=1}^{N-1} u(kh)v(kh)$$

with the corresponding norm $\|v\|_h$.

Define $\varphi_m \in L(I_h)$ by $\varphi_m(x) = \sin \pi m x$.

Numerical Methods for PDE: 2nd Order Elliptic PDE

Stability Analysis using Fourier Analysis

First we consider the one dimensional case. With $h = 1/N$, let $I_h = \{h, 2h, \dots, (N-1)h\}$ and let

$$L(I_h) = \{u: \bar{I}_h \rightarrow \mathbb{R} : u(0) = 0, u(1) = 0\}.$$

Clearly, $L(I_h)$ is isomorphic to \mathbb{R}^{N-1} . On $L(I_h)$, we define the inner product

$$\langle u, v \rangle_h = h \sum_{k=1}^{N-1} u(kh)v(kh)$$

with the corresponding norm $\|v\|_h$.

Define $\varphi_m \in L(I_h)$ by $\varphi_m(x) = \sin \pi m x$. Then

$$D_h^2 \varphi_m(x) = \frac{\sin \pi m(x+h) - 2\sin \pi m x + \sin \pi m(x-h)}{h^2}$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

Stability Analysis using Fourier Analysis

First we consider the one dimensional case. With $h = 1/N$, let $I_h = \{h, 2h, \dots, (N-1)h\}$ and let

$$L(I_h) = \{u: \bar{I}_h \rightarrow \mathbb{R} : u(0) = 0, u(1) = 0\}.$$

Clearly, $L(I_h)$ is isomorphic to \mathbb{R}^{N-1} . On $L(I_h)$, we define the inner product

$$\langle u, v \rangle_h = h \sum_{k=1}^{N-1} u(kh)v(kh)$$

with the corresponding norm $\|v\|_h$.

Define $\varphi_m \in L(I_h)$ by $\varphi_m(x) = \sin \pi m x$. Then

$$D_h^2 \varphi_m(x) = \frac{\sin \pi m(x+h) - 2\sin \pi m x + \sin \pi m(x-h)}{h^2} = \frac{2}{h^2} (\cos \pi m h - 1) \sin \pi m x.$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

Stability Analysis using Fourier Analysis

First we consider the one dimensional case. With $h = 1/N$, let $I_h = \{h, 2h, \dots, (N-1)h\}$ and let

$$L(I_h) = \{u: \bar{I}_h \rightarrow \mathbb{R} : u(0) = 0, u(1) = 0\}.$$

Clearly, $L(I_h)$ is isomorphic to \mathbb{R}^{N-1} . On $L(I_h)$, we define the inner product

$$\langle u, v \rangle_h = h \sum_{k=1}^{N-1} u(kh)v(kh)$$

with the corresponding norm $\|v\|_h$.

Define $\varphi_m \in L(I_h)$ by $\varphi_m(x) = \sin \pi m x$. Then

$$D_h^2 \varphi_m(x) = \frac{\sin \pi m(x+h) - 2\sin \pi m x + \sin \pi m(x-h)}{h^2} = \frac{2}{h^2} (\cos \pi m h - 1) \sin \pi m x.$$

Thus,

$$D_h^2 \varphi_m = -\lambda_m \varphi_m, \quad \lambda_m = \frac{2}{h^2} (1 - \cos \pi m h) = \frac{4}{h^2} \sin^2 \frac{\pi m h}{2}.$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

Stability Analysis using Fourier Analysis

First we consider the one dimensional case. With $h = 1/N$, let $I_h = \{h, 2h, \dots, (N-1)h\}$ and let

$$L(I_h) = \{u: \bar{I}_h \rightarrow \mathbb{R} : u(0) = 0, u(1) = 0\}.$$

Clearly, $L(I_h)$ is isomorphic to \mathbb{R}^{N-1} . On $L(I_h)$, we define the inner product

$$\langle u, v \rangle_h = h \sum_{k=1}^{N-1} u(kh)v(kh)$$

with the corresponding norm $\|v\|_h$.

Define $\varphi_m \in L(I_h)$ by $\varphi_m(x) = \sin \pi m x$. Then

$$D_h^2 \varphi_m(x) = \frac{\sin \pi m(x+h) - 2\sin \pi m x + \sin \pi m(x-h)}{h^2} = \frac{2}{h^2} (\cos \pi m h - 1) \sin \pi m x.$$

Thus,

$$D_h^2 \varphi_m = -\lambda_m \varphi_m, \quad \lambda_m = \frac{2}{h^2} (1 - \cos \pi m h) = \frac{4}{h^2} \sin^2 \frac{\pi m h}{2}.$$

Note that $0 < \lambda_1 < \lambda_2 < \dots < \lambda_{N-1} < 4/h^2$.

Numerical Methods for PDE: 2nd Order Elliptic PDE

Stability Analysis using Fourier Analysis

First we consider the one dimensional case. With $h = 1/N$, let $I_h = \{h, 2h, \dots, (N-1)h\}$ and let

$$L(I_h) = \{u: \bar{I}_h \rightarrow \mathbb{R} : u(0) = 0, u(1) = 0\}.$$

Clearly, $L(I_h)$ is isomorphic to \mathbb{R}^{N-1} . On $L(I_h)$, we define the inner product

$$\langle u, v \rangle_h = h \sum_{k=1}^{N-1} u(kh)v(kh)$$

with the corresponding norm $\|v\|_h$.

Define $\varphi_m \in L(I_h)$ by $\varphi_m(x) = \sin \pi m x$. Then

$$D_h^2 \varphi_m(x) = \frac{\sin \pi m(x+h) - 2\sin \pi m x + \sin \pi m(x-h)}{h^2} = \frac{2}{h^2} (\cos \pi m h - 1) \sin \pi m x.$$

Thus,

$$D_h^2 \varphi_m = -\lambda_m \varphi_m, \quad \lambda_m = \frac{2}{h^2} (1 - \cos \pi m h) = \frac{4}{h^2} \sin^2 \frac{\pi m h}{2}.$$

Note that $0 < \lambda_1 < \lambda_2 < \dots < \lambda_{N-1} < 4/h^2$. Also, $\lambda_1 = 8$ for $N = 2$ and λ_1 increases with N .

Numerical Methods for PDE: 2nd Order Elliptic PDE

Stability Analysis using Fourier Analysis

First we consider the one dimensional case. With $h = 1/N$, let $I_h = \{h, 2h, \dots, (N-1)h\}$ and let

$$L(I_h) = \{u: \bar{I}_h \rightarrow \mathbb{R} : u(0) = 0, u(1) = 0\}.$$

Clearly, $L(I_h)$ is isomorphic to \mathbb{R}^{N-1} . On $L(I_h)$, we define the inner product

$$\langle u, v \rangle_h = h \sum_{k=1}^{N-1} u(kh)v(kh)$$

with the corresponding norm $\|v\|_h$.

Define $\varphi_m \in L(I_h)$ by $\varphi_m(x) = \sin \pi m x$. Then

$$D_h^2 \varphi_m(x) = \frac{\sin \pi m(x+h) - 2\sin \pi m x + \sin \pi m(x-h)}{h^2} = \frac{2}{h^2} (\cos \pi m h - 1) \sin \pi m x.$$

Thus,

$$D_h^2 \varphi_m = -\lambda_m \varphi_m, \quad \lambda_m = \frac{2}{h^2} (1 - \cos \pi m h) = \frac{4}{h^2} \sin^2 \frac{\pi m h}{2}.$$

Note that $0 < \lambda_1 < \lambda_2 < \dots < \lambda_{N-1} < 4/h^2$. Also, $\lambda_1 = 8$ for $N = 2$ and λ_1 increases with N . Moreover, for $m \neq n$,

$$\langle \varphi_m, \varphi_n \rangle_h = h \sum_{k=1}^{N-1} \sin \pi m k h \sin \pi n k h$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

Stability Analysis using Fourier Analysis

First we consider the one dimensional case. With $h = 1/N$, let $I_h = \{h, 2h, \dots, (N-1)h\}$ and let

$$L(I_h) = \{u: \bar{I}_h \rightarrow \mathbb{R} : u(0) = 0, u(1) = 0\}.$$

Clearly, $L(I_h)$ is isomorphic to \mathbb{R}^{N-1} . On $L(I_h)$, we define the inner product

$$\langle u, v \rangle_h = h \sum_{k=1}^{N-1} u(kh)v(kh)$$

with the corresponding norm $\|v\|_h$.

Define $\varphi_m \in L(I_h)$ by $\varphi_m(x) = \sin \pi m x$. Then

$$D_h^2 \varphi_m(x) = \frac{\sin \pi m(x+h) - 2\sin \pi m x + \sin \pi m(x-h)}{h^2} = \frac{2}{h^2} (\cos \pi m h - 1) \sin \pi m x.$$

Thus,

$$D_h^2 \varphi_m = -\lambda_m \varphi_m, \quad \lambda_m = \frac{2}{h^2} (1 - \cos \pi m h) = \frac{4}{h^2} \sin^2 \frac{\pi m h}{2}.$$

Note that $0 < \lambda_1 < \lambda_2 < \dots < \lambda_{N-1} < 4/h^2$. Also, $\lambda_1 = 8$ for $N = 2$ and λ_1 increases with N . Moreover, for $m \neq n$,

$$\langle \varphi_m, \varphi_n \rangle_h = h \sum_{k=1}^{N-1} \sin \pi m k h \sin \pi n k h = \frac{h}{2} \sum_{k=1}^{N-1} \cos \pi(m-n)kh - \frac{h}{2} \sum_{k=1}^{N-1} \cos \pi(m+n)kh$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

Define $\varphi_m \in L(I_h)$ by $\varphi_m(x) = \sin \pi m x$. Then

$$D_h^2 \varphi_m(x) = \frac{\sin \pi m(x+h) - 2\sin \pi m x + \sin \pi m(x-h)}{h^2} = \frac{2}{h^2} (\cos \pi m h - 1) \sin \pi m x.$$

Thus,

$$D_h^2 \varphi_m = -\lambda_m \varphi_m, \quad \lambda_m = \frac{2}{h^2} (1 - \cos \pi m h) = \frac{4}{h^2} \sin^2 \frac{\pi m h}{2}.$$

Note that $0 < \lambda_1 < \lambda_2 < \dots < \lambda_{N-1} < 4/h^2$. Also, $\lambda_1 = 8$ for $N = 2$ and λ_1 increases with N . Moreover, for $m \neq n$,

$$\langle \varphi_m, \varphi_n \rangle_h = h \sum_{k=1}^{N-1} \sin \pi m k h \sin \pi n k h = \frac{h}{2} \sum_{k=1}^{N-1} \cos \pi(m-n) k h - \frac{h}{2} \sum_{k=1}^{N-1} \cos \pi(m+n) k h$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

Define $\varphi_m \in L(I_h)$ by $\varphi_m(x) = \sin \pi m x$. Then

$$D_h^2 \varphi_m(x) = \frac{\sin \pi m(x+h) - 2\sin \pi m x + \sin \pi m(x-h)}{h^2} = \frac{2}{h^2} (\cos \pi m h - 1) \sin \pi m x.$$

Thus,

$$D_h^2 \varphi_m = -\lambda_m \varphi_m, \quad \lambda_m = \frac{2}{h^2} (1 - \cos \pi m h) = \frac{4}{h^2} \sin^2 \frac{\pi m h}{2}.$$

Note that $0 < \lambda_1 < \lambda_2 < \dots < \lambda_{N-1} < 4/h^2$. Also, $\lambda_1 = 8$ for $N = 2$ and λ_1 increases with N . Moreover, for $m \neq n$,

$$\begin{aligned} \langle \varphi_m, \varphi_n \rangle_h &= h \sum_{k=1}^{N-1} \sin \pi m k h \sin \pi n k h = \frac{h}{2} \sum_{k=1}^{N-1} \cos \pi(m-n) k h - \frac{h}{2} \sum_{k=1}^{N-1} \cos \pi(m+n) k h \\ &= \frac{h}{2} \sum_{k=0}^{N-1} \cos \pi(m-n) k h - \frac{h}{2} \sum_{k=0}^{N-1} \cos \pi(m+n) k h \end{aligned}$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

Define $\varphi_m \in L(I_h)$ by $\varphi_m(x) = \sin \pi m x$. Then

$$D_h^2 \varphi_m(x) = \frac{\sin \pi m(x+h) - 2\sin \pi m x + \sin \pi m(x-h)}{h^2} = \frac{2}{h^2} (\cos \pi m h - 1) \sin \pi m x.$$

Thus,

$$D_h^2 \varphi_m = -\lambda_m \varphi_m, \quad \lambda_m = \frac{2}{h^2} (1 - \cos \pi m h) = \frac{4}{h^2} \sin^2 \frac{\pi m h}{2}.$$

Note that $0 < \lambda_1 < \lambda_2 < \dots < \lambda_{N-1} < 4/h^2$. Also, $\lambda_1 = 8$ for $N = 2$ and λ_1 increases with N . Moreover, for $m \neq n$,

$$\begin{aligned} \langle \varphi_m, \varphi_n \rangle_h &= h \sum_{k=1}^{N-1} \sin \pi m k h \sin \pi n k h = \frac{h}{2} \sum_{k=1}^{N-1} \cos \pi(m-n) k h - \frac{h}{2} \sum_{k=1}^{N-1} \cos \pi(m+n) k h \\ &= \frac{h}{2} \sum_{k=0}^{N-1} \cos \pi(m-n) k h - \frac{h}{2} \sum_{k=0}^{N-1} \cos \pi(m+n) k h = \frac{h}{2} \operatorname{Re} \left[\sum_{k=0}^{N-1} e^{i\pi(m-n) k h} - \sum_{k=0}^{N-1} e^{i\pi(m+n) k h} \right] \end{aligned}$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

Define $\varphi_m \in L(I_h)$ by $\varphi_m(x) = \sin \pi m x$. Then

$$D_h^2 \varphi_m(x) = \frac{\sin \pi m(x+h) - 2\sin \pi m x + \sin \pi m(x-h)}{h^2} = \frac{2}{h^2} (\cos \pi m h - 1) \sin \pi m x.$$

Thus,

$$D_h^2 \varphi_m = -\lambda_m \varphi_m, \quad \lambda_m = \frac{2}{h^2} (1 - \cos \pi m h) = \frac{4}{h^2} \sin^2 \frac{\pi m h}{2}.$$

Note that $0 < \lambda_1 < \lambda_2 < \dots < \lambda_{N-1} < 4/h^2$. Also, $\lambda_1 = 8$ for $N = 2$ and λ_1 increases with N . Moreover, for $m \neq n$,

$$\begin{aligned} \langle \varphi_m, \varphi_n \rangle_h &= h \sum_{k=1}^{N-1} \sin \pi m k h \sin \pi n k h = \frac{h}{2} \sum_{k=1}^{N-1} \cos \pi(m-n) k h - \frac{h}{2} \sum_{k=1}^{N-1} \cos \pi(m+n) k h \\ &= \frac{h}{2} \sum_{k=0}^{N-1} \cos \pi(m-n) k h - \frac{h}{2} \sum_{k=0}^{N-1} \cos \pi(m+n) k h = \frac{h}{2} \operatorname{Re} \left[\sum_{k=0}^{N-1} e^{i\pi(m-n) k h} - \sum_{k=0}^{N-1} e^{i\pi(m+n) k h} \right] \\ &= \frac{h}{2} \operatorname{Re} \left[\frac{e^{i\pi(m-n) N h} - 1}{e^{i\pi(m-n) h} - 1} - \frac{e^{i\pi(m+n) N h} - 1}{e^{i\pi(m+n) h} - 1} \right] \end{aligned}$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

Define $\varphi_m \in L(I_h)$ by $\varphi_m(x) = \sin \pi m x$. Then

$$D_h^2 \varphi_m(x) = \frac{\sin \pi m(x+h) - 2\sin \pi m x + \sin \pi m(x-h)}{h^2} = \frac{2}{h^2} (\cos \pi m h - 1) \sin \pi m x.$$

Thus,

$$D_h^2 \varphi_m = -\lambda_m \varphi_m, \quad \lambda_m = \frac{2}{h^2} (1 - \cos \pi m h) = \frac{4}{h^2} \sin^2 \frac{\pi m h}{2}.$$

Note that $0 < \lambda_1 < \lambda_2 < \dots < \lambda_{N-1} < 4/h^2$. Also, $\lambda_1 = 8$ for $N = 2$ and λ_1 increases with N . Moreover, for $m \neq n$,

$$\begin{aligned} \langle \varphi_m, \varphi_n \rangle_h &= h \sum_{k=1}^{N-1} \sin \pi m k h \sin \pi n k h = \frac{h}{2} \sum_{k=1}^{N-1} \cos \pi(m-n) k h - \frac{h}{2} \sum_{k=1}^{N-1} \cos \pi(m+n) k h \\ &= \frac{h}{2} \sum_{k=0}^{N-1} \cos \pi(m-n) k h - \frac{h}{2} \sum_{k=0}^{N-1} \cos \pi(m+n) k h = \frac{h}{2} \operatorname{Re} \left[\sum_{k=0}^{N-1} e^{i\pi(m-n) k h} - \sum_{k=0}^{N-1} e^{i\pi(m+n) k h} \right] \\ &= \frac{h}{2} \operatorname{Re} \left[\frac{e^{i\pi(m-n) N h} - 1}{e^{i\pi(m-n) h} - 1} - \frac{e^{i\pi(m+n) N h} - 1}{e^{i\pi(m+n) h} - 1} \right] = \frac{h}{2} \operatorname{Re} \left[\frac{(-1)^{(m-n)} - 1}{e^{i\pi(m-n) h} - 1} - \frac{(-1)^{(m+n)} - 1}{e^{i\pi(m+n) h} - 1} \right] \end{aligned}$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

Define $\varphi_m \in L(I_h)$ by $\varphi_m(x) = \sin \pi m x$. Then

$$D_h^2 \varphi_m(x) = \frac{\sin \pi m(x+h) - 2\sin \pi m x + \sin \pi m(x-h)}{h^2} = \frac{2}{h^2} (\cos \pi m h - 1) \sin \pi m x.$$

Thus,

$$D_h^2 \varphi_m = -\lambda_m \varphi_m, \quad \lambda_m = \frac{2}{h^2} (1 - \cos \pi m h) = \frac{4}{h^2} \sin^2 \frac{\pi m h}{2}.$$

Note that $0 < \lambda_1 < \lambda_2 < \dots < \lambda_{N-1} < 4/h^2$. Also, $\lambda_1 = 8$ for $N = 2$ and λ_1 increases with N . Moreover, for $m \neq n$,

$$\begin{aligned} \langle \varphi_m, \varphi_n \rangle_h &= h \sum_{k=1}^{N-1} \sin \pi m k h \sin \pi n k h = \frac{h}{2} \sum_{k=1}^{N-1} \cos \pi(m-n) k h - \frac{h}{2} \sum_{k=1}^{N-1} \cos \pi(m+n) k h \\ &= \frac{h}{2} \sum_{k=0}^{N-1} \cos \pi(m-n) k h - \frac{h}{2} \sum_{k=0}^{N-1} \cos \pi(m+n) k h = \frac{h}{2} \operatorname{Re} \left[\sum_{k=0}^{N-1} e^{i\pi(m-n) k h} - \sum_{k=0}^{N-1} e^{i\pi(m+n) k h} \right] \\ &= \frac{h}{2} \operatorname{Re} \left[\frac{e^{i\pi(m-n) N h} - 1}{e^{i\pi(m-n) h} - 1} - \frac{e^{i\pi(m+n) N h} - 1}{e^{i\pi(m+n) h} - 1} \right] = \frac{h}{2} \operatorname{Re} \left[\frac{(-1)^{(m-n)} - 1}{e^{i\pi(m-n) h} - 1} - \frac{(-1)^{(m+n)} - 1}{e^{i\pi(m+n) h} - 1} \right] \\ &= \frac{h}{2} \left((-1)^{(m-n)} - 1 \right) \left[\frac{\cos \pi(m-n) h - 1}{2 - 2 \cos \pi(m-n) h} - \frac{\cos \pi(m+n) h - 1}{2 - 2 \cos \pi(m+n) h} \right] \end{aligned}$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

Define $\varphi_m \in L(I_h)$ by $\varphi_m(x) = \sin \pi m x$. Then

$$D_h^2 \varphi_m(x) = \frac{\sin \pi m(x+h) - 2\sin \pi m x + \sin \pi m(x-h)}{h^2} = \frac{2}{h^2} (\cos \pi m h - 1) \sin \pi m x.$$

Thus,

$$D_h^2 \varphi_m = -\lambda_m \varphi_m, \quad \lambda_m = \frac{2}{h^2} (1 - \cos \pi m h) = \frac{4}{h^2} \sin^2 \frac{\pi m h}{2}.$$

Note that $0 < \lambda_1 < \lambda_2 < \dots < \lambda_{N-1} < 4/h^2$. Also, $\lambda_1 = 8$ for $N = 2$ and λ_1 increases with N . Moreover, for $m \neq n$,

$$\begin{aligned} \langle \varphi_m, \varphi_n \rangle_h &= h \sum_{k=1}^{N-1} \sin \pi m k h \sin \pi n k h = \frac{h}{2} \sum_{k=1}^{N-1} \cos \pi(m-n) k h - \frac{h}{2} \sum_{k=1}^{N-1} \cos \pi(m+n) k h \\ &= \frac{h}{2} \sum_{k=0}^{N-1} \cos \pi(m-n) k h - \frac{h}{2} \sum_{k=0}^{N-1} \cos \pi(m+n) k h = \frac{h}{2} \operatorname{Re} \left[\sum_{k=0}^{N-1} e^{i\pi(m-n) k h} - \sum_{k=0}^{N-1} e^{i\pi(m+n) k h} \right] \\ &= \frac{h}{2} \operatorname{Re} \left[\frac{e^{i\pi(m-n) N h} - 1}{e^{i\pi(m-n) h} - 1} - \frac{e^{i\pi(m+n) N h} - 1}{e^{i\pi(m+n) h} - 1} \right] = \frac{h}{2} \operatorname{Re} \left[\frac{(-1)^{(m-n)} - 1}{e^{i\pi(m-n) h} - 1} - \frac{(-1)^{(m+n)} - 1}{e^{i\pi(m+n) h} - 1} \right] \\ &= \frac{h}{2} \left((-1)^{(m-n)} - 1 \right) \left[\frac{\cos \pi(m-n) h - 1}{2 - 2 \cos \pi(m-n) h} - \frac{\cos \pi(m+n) h - 1}{2 - 2 \cos \pi(m+n) h} \right] = 0 \end{aligned}$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

Define $\varphi_m \in L(I_h)$ by $\varphi_m(x) = \sin \pi m x$. Then

$$D_h^2 \varphi_m(x) = \frac{\sin \pi m(x+h) - 2\sin \pi m x + \sin \pi m(x-h)}{h^2} = \frac{2}{h^2} (\cos \pi m h - 1) \sin \pi m x.$$

Thus,

$$D_h^2 \varphi_m = -\lambda_m \varphi_m, \quad \lambda_m = \frac{2}{h^2} (1 - \cos \pi m h) = \frac{4}{h^2} \sin^2 \frac{\pi m h}{2}.$$

Note that $0 < \lambda_1 < \lambda_2 < \dots < \lambda_{N-1} < 4/h^2$. Also, $\lambda_1 = 8$ for $N = 2$ and λ_1 increases with N . Moreover, for $m \neq n$,

$$\langle \varphi_m, \varphi_n \rangle_h = 0.$$

Theorem

The functions $\varphi_m, m = 1, 2, \dots, N-1$ form an orthogonal basis of $L(I_h)$. Consequently, any function $v \in L(I_h)$ can be expanded as $v = \sum_{m=1}^{N-1} a_m \varphi_m$ with

$$a_m = \frac{\langle v, \varphi_m \rangle_h}{\|\varphi_m\|_h^2}, \quad \|v\|_h^2 = \sum_{m=1}^{N-1} a_m^2 \|\varphi_m\|_h^2.$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

Define $\varphi_m \in L(I_h)$ by $\varphi_m(x) = \sin \pi m x$. Then

$$D_h^2 \varphi_m(x) = \frac{\sin \pi m(x+h) - 2\sin \pi m x + \sin \pi m(x-h)}{h^2} = \frac{2}{h^2} (\cos \pi m h - 1) \sin \pi m x.$$

Thus,

$$D_h^2 \varphi_m = -\lambda_m \varphi_m, \quad \lambda_m = \frac{2}{h^2} (1 - \cos \pi m h) = \frac{4}{h^2} \sin^2 \frac{\pi m h}{2}.$$

Note that $0 < \lambda_1 < \lambda_2 < \dots < \lambda_{N-1} < 4/h^2$. Also, $\lambda_1 = 8$ for $N = 2$ and λ_1 increases with N . Moreover, for $m \neq n$,

$$\langle \varphi_m, \varphi_n \rangle_h = 0.$$

Theorem

The functions $\varphi_m, m = 1, 2, \dots, N-1$ form an orthogonal basis of $L(I_h)$. Consequently, any function $v \in L(I_h)$ can be expanded as $v = \sum_{m=1}^{N-1} a_m \varphi_m$ with

$$a_m = \frac{\langle v, \varphi_m \rangle_h}{\|\varphi_m\|_h^2}, \quad \|v\|_h^2 = \sum_{m=1}^{N-1} a_m^2 \|\varphi_m\|_h^2.$$

From this, we obtain the stability result for the one-dimensional Laplacian: if $f = D_h^2 v = -\sum_{m=1}^{N-1} \lambda_m a_m \varphi_m$, then

$$\|f\|_h^2 = \sum_{m=1}^{N-1} \lambda_m^2 a_m^2 \|\varphi_m\|_h^2 \geq 8^2 \|v\|_h^2$$



Thus, we have the stability estimate

$$\|v\|_h \leq \frac{1}{8} \|f\|_h.$$



Thus, we have the stability estimate

$$\|v\|_h \leq \frac{1}{8} \|f\|_h.$$

The extension to two-dimensional case is straightforward.

Let $L(\Omega_h) = \{u: \bar{\Omega}_h \rightarrow \mathbb{R} : u(x) = 0, x \in \Gamma_h\}$ so that $L(\Omega_h)$ is isomorphic to \mathbb{R}^M , $M = (N - 1)^2$.



Thus, we have the stability estimate

$$\|v\|_h \leq \frac{1}{8} \|f\|_h.$$

The extension to two-dimensional case is straightforward.

Let $L(\Omega_h) = \{u: \bar{\Omega}_h \rightarrow \mathbb{R} : u(x) = 0, x \in \Gamma_h\}$ so that $L(\Omega_h)$ is isomorphic to \mathbb{R}^M , $M = (N - 1)^2$.

We use the basis

$$\varphi_{mn}(x_1, x_2) = \varphi_m(x_1)\varphi_n(x_2), \quad m, n = 1, \dots, N - 1.$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

Thus, we have the stability estimate

$$\|v\|_h \leq \frac{1}{8} \|f\|_h.$$

The extension to two-dimensional case is straightforward.

Let $L(\Omega_h) = \{u: \bar{\Omega}_h \rightarrow \mathbb{R} : u(x) = 0, x \in \Gamma_h\}$ so that $L(\Omega_h)$ is isomorphic to \mathbb{R}^M , $M = (N - 1)^2$.

We use the basis

$$\varphi_{mn}(x_1, x_2) = \varphi_m(x_1)\varphi_n(x_2), \quad m, n = 1, \dots, N - 1.$$

It is easy to check (*exercise*) that these $(N - 1)^2$ functions form an orthogonal basis for $L(\Omega_h)$ equipped with the inner product

$$\langle u, v \rangle_h = h^2 \sum_{m=1}^{N-1} \sum_{n=1}^{N-1} u(mh, nh)v(mh, nh)$$

and the corresponding norm $\|\cdot\|_h$.

Numerical Methods for PDE: 2nd Order Elliptic PDE

Thus, we have the stability estimate

$$\|v\|_h \leq \frac{1}{8} \|f\|_h.$$

The extension to two-dimensional case is straightforward.

Let $L(\Omega_h) = \{u: \bar{\Omega}_h \rightarrow \mathbb{R} : u(x) = 0, x \in \Gamma_h\}$ so that $L(\Omega_h)$ is isomorphic to \mathbb{R}^M , $M = (N - 1)^2$.

We use the basis

$$\varphi_{mn}(x_1, x_2) = \varphi_m(x_1)\varphi_n(x_2), \quad m, n = 1, \dots, N - 1.$$

It is easy to check (*exercise*) that these $(N - 1)^2$ functions form an orthogonal basis for $L(\Omega_h)$ equipped with the inner product

$$\langle u, v \rangle_h = h^2 \sum_{m=1}^{N-1} \sum_{n=1}^{N-1} u(mh, nh)v(mh, nh)$$

and the corresponding norm $\|\cdot\|_h$. Moreover, we have

$$\Delta_h \varphi_{mn} = -\lambda_{mn} \varphi_{mn}$$

where $\lambda_{mn} = \lambda_m + \lambda_n \geq 16$.

Numerical Methods for PDE: 2nd Order Elliptic PDE

Thus, we have the stability estimate

$$\|v\|_h \leq \frac{1}{8} \|f\|_h.$$

The extension to two-dimensional case is straightforward.

Let $L(\Omega_h) = \{u: \bar{\Omega}_h \rightarrow \mathbb{R} : u(x) = 0, x \in \Gamma_h\}$ so that $L(\Omega_h)$ is isomorphic to \mathbb{R}^M , $M = (N - 1)^2$.

We use the basis

$$\varphi_{mn}(x_1, x_2) = \varphi_m(x_1)\varphi_n(x_2), \quad m, n = 1, \dots, N - 1.$$

It is easy to check (*exercise*) that these $(N - 1)^2$ functions form an orthogonal basis for $L(\Omega_h)$ equipped with the inner product

$$\langle u, v \rangle_h = h^2 \sum_{m=1}^{N-1} \sum_{n=1}^{N-1} u(mh, nh) v(mh, nh)$$

and the corresponding norm $\|\cdot\|_h$. Moreover, we have

$$\Delta_h \varphi_{mn} = -\lambda_{mn} \varphi_{mn}$$

where $\lambda_{mn} = \lambda_m + \lambda_n \geq 16$.

Theorem

We have $\|v\|_h \leq \frac{1}{16} \|f\|_h$ as the stability estimate where v solves the discrete problem $\Delta_h v = f$, on Ω_h , $v = 0$, on Γ_h .