

# Initial Value Problems: Linear Multistep Methods

## Theorem

A linear multistep is consistent if and only if

$$\sum_{j=-1}^k a_j = 0, \quad \sum_{j=-1}^k j a_j + \sum_{j=-1}^k b_j = 0.$$

The method is of order  $p$  if and only if

$$\sum_{j=-1}^k (-j)^m a_j - m \sum_{j=-1}^k (-j)^{m-1} b_j = 0, \quad m = 0, 1, \dots, p.$$

## Example

Construct the 2-step method of highest order!

# Initial Value Problems: Linear Multistep Methods

## Theorem

A linear multistep is consistent if and only if

$$\sum_{j=-1}^k a_j = 0, \quad \sum_{j=-1}^k j a_j + \sum_{j=-1}^k b_j = 0.$$

The method is of order  $p$  if and only if

$$\sum_{j=-1}^k (-j)^m a_j - m \sum_{j=-1}^k (-j)^{m-1} b_j = 0, \quad m = 0, 1, \dots, p.$$

## Example

Construct the 2-step method of highest order!

For a 2-step method, there are 5 undetermined coefficients:  $a_0, a_1, b_{-1}, b_0, b_1$ .

# Initial Value Problems: Linear Multistep Methods

## Theorem

A linear multistep is consistent if and only if

$$\sum_{j=-1}^k a_j = 0, \quad \sum_{j=-1}^k j a_j + \sum_{j=-1}^k b_j = 0.$$

The method is of order  $p$  if and only if

$$\sum_{j=-1}^k (-j)^m a_j - m \sum_{j=-1}^k (-j)^{m-1} b_j = 0, \quad m = 0, 1, \dots, p.$$

## Example

Construct the 2-step method of highest order!

For a 2-step method, there are 5 undetermined coefficients:  $a_0, a_1, b_{-1}, b_0, b_1$ . The first five order conditions are

$$1 + a_0 + a_1 = 0,$$

# Initial Value Problems: Linear Multistep Methods

## Theorem

A linear multistep is consistent if and only if

$$\sum_{j=-1}^k a_j = 0, \quad \sum_{j=-1}^k j a_j + \sum_{j=-1}^k b_j = 0.$$

The method is of order  $p$  if and only if

$$\sum_{j=-1}^k (-j)^m a_j - m \sum_{j=-1}^k (-j)^{m-1} b_j = 0, \quad m = 0, 1, \dots, p.$$

## Example

Construct the 2-step method of highest order!

For a 2-step method, there are 5 undetermined coefficients:  $a_0, a_1, b_{-1}, b_0, b_1$ . The first five order conditions are

$$1 + a_0 + a_1 = 0, \quad 1 - a_1 - b_{-1} - b_0 - b_1 = 0,$$

# Initial Value Problems: Linear Multistep Methods

## Theorem

A linear multistep is consistent if and only if

$$\sum_{j=-1}^k a_j = 0, \quad \sum_{j=-1}^k j a_j + \sum_{j=-1}^k b_j = 0.$$

The method is of order  $p$  if and only if

$$\sum_{j=-1}^k (-j)^m a_j - m \sum_{j=-1}^k (-j)^{m-1} b_j = 0, \quad m = 0, 1, \dots, p.$$

## Example

Construct the 2-step method of highest order!

For a 2-step method, there are 5 undetermined coefficients:  $a_0, a_1, b_{-1}, b_0, b_1$ . The first five order conditions are

$$1 + a_0 + a_1 = 0, \quad 1 - a_1 - b_{-1} - b_0 - b_1 = 0, \quad 1 + a_1 - 2b_{-1} + 2b_1 = 0,$$

# Initial Value Problems: Linear Multistep Methods

## Theorem

A linear multistep is consistent if and only if

$$\sum_{j=-1}^k a_j = 0, \quad \sum_{j=-1}^k j a_j + \sum_{j=-1}^k b_j = 0.$$

The method is of order  $p$  if and only if

$$\sum_{j=-1}^k (-j)^m a_j - m \sum_{j=-1}^k (-j)^{m-1} b_j = 0, \quad m = 0, 1, \dots, p.$$

## Example

Construct the 2-step method of highest order!

For a 2-step method, there are 5 undetermined coefficients:  $a_0, a_1, b_{-1}, b_0, b_1$ . The first five order conditions are

$$\begin{aligned} 1 + a_0 + a_1 &= 0, & 1 - a_1 - b_{-1} - b_0 - b_1 &= 0, & 1 + a_1 - 2b_{-1} + 2b_1 &= 0, \\ 1 - a_1 - 3b_{-1} - 3b_1 &= 0, \end{aligned}$$

# Initial Value Problems: Linear Multistep Methods

## Theorem

A linear multistep is consistent if and only if

$$\sum_{j=-1}^k a_j = 0, \quad \sum_{j=-1}^k j a_j + \sum_{j=-1}^k b_j = 0.$$

The method is of order  $p$  if and only if

$$\sum_{j=-1}^k (-j)^m a_j - m \sum_{j=-1}^k (-j)^{m-1} b_j = 0, \quad m = 0, 1, \dots, p.$$

## Example

Construct the 2-step method of highest order!

For a 2-step method, there are 5 undetermined coefficients:  $a_0, a_1, b_{-1}, b_0, b_1$ . The first five order conditions are

$$\begin{aligned} 1 + a_0 + a_1 &= 0, & 1 - a_1 - b_{-1} - b_0 - b_1 &= 0, & 1 + a_1 - 2b_{-1} + 2b_1 &= 0, \\ 1 - a_1 - 3b_{-1} - 3b_1 &= 0, & 1 + a_1 - 4b_{-1} + 4b_1 &= 0. \end{aligned}$$

# Initial Value Problems: Linear Multistep Methods

## Theorem

A linear multistep is consistent if and only if

$$\sum_{j=-1}^k a_j = 0, \quad \sum_{j=-1}^k j a_j + \sum_{j=-1}^k b_j = 0.$$

The method is of order  $p$  if and only if

$$\sum_{j=-1}^k (-j)^m a_j - m \sum_{j=-1}^k (-j)^{m-1} b_j = 0, \quad m = 0, 1, \dots, p.$$

## Example

Construct the 2-step method of highest order!

For a 2-step method, there are 5 undetermined coefficients:  $a_0, a_1, b_{-1}, b_0, b_1$ . The first five order conditions are

$$\begin{aligned} 1 + a_0 + a_1 &= 0, & 1 - a_1 - b_{-1} - b_0 - b_1 &= 0, & 1 + a_1 - 2b_{-1} + 2b_1 &= 0, \\ 1 - a_1 - 3b_{-1} - 3b_1 &= 0, & 1 + a_1 - 4b_{-1} + 4b_1 &= 0. \end{aligned}$$

This system of linear equation has a unique solution

$$a_0 = 0, a_1 = -1, b_{-1} = \frac{1}{3}, b_0 = \frac{4}{3}, b_1 = \frac{1}{3}.$$



# Initial Value Problems: Linear Multistep Methods

## Theorem

A linear multistep is consistent if and only if

$$\sum_{j=-1}^k a_j = 0, \quad \sum_{j=-1}^k j a_j + \sum_{j=-1}^k b_j = 0.$$

The method is of order  $p$  if and only if

$$\sum_{j=-1}^k (-j)^m a_j - m \sum_{j=-1}^k (-j)^{m-1} b_j = 0, \quad m = 0, 1, \dots, p.$$

## Example

Construct the 2-step method of highest order!

For a 2-step method, there are 5 undetermined coefficients:  $a_0, a_1, b_{-1}, b_0, b_1$ . The first five order conditions are

$$\begin{aligned} 1 + a_0 + a_1 &= 0, & 1 - a_1 - b_{-1} - b_0 - b_1 &= 0, & 1 + a_1 - 2b_{-1} + 2b_1 &= 0, \\ 1 - a_1 - 3b_{-1} - 3b_1 &= 0, & 1 + a_1 - 4b_{-1} + 4b_1 &= 0. \end{aligned}$$

This system of linear equation has a unique solution

$$a_0 = 0, a_1 = -1, b_{-1} = \frac{1}{3}, b_0 = \frac{4}{3}, b_1 = \frac{1}{3}.$$

This scheme is known as Milne-Simpson method and it is the unique fourth order 2-step method.

# Initial Value Problems: Linear Multistep Methods

## Theorem

A linear multistep is consistent if and only if

$$\sum_{j=-1}^k a_j = 0, \quad \sum_{j=-1}^k j a_j + \sum_{j=-1}^k b_j = 0.$$

The method is of order  $p$  if and only if

$$\sum_{j=-1}^k (-j)^m a_j - m \sum_{j=-1}^k (-j)^{m-1} b_j = 0, \quad m = 0, 1, \dots, p.$$

## Example

Construct the explicit 2-step method of highest order!

# Initial Value Problems: Linear Multistep Methods

## Theorem

A linear multistep is consistent if and only if

$$\sum_{j=-1}^k a_j = 0, \quad \sum_{j=-1}^k j a_j + \sum_{j=-1}^k b_j = 0.$$

The method is of order  $p$  if and only if

$$\sum_{j=-1}^k (-j)^m a_j - m \sum_{j=-1}^k (-j)^{m-1} b_j = 0, \quad m = 0, 1, \dots, p.$$

## Example

Construct the explicit 2-step method of highest order!

There are 4 undetermined coefficients:  $a_0, a_1, b_0, b_1$ .

# Initial Value Problems: Linear Multistep Methods

## Theorem

A linear multistep is consistent if and only if

$$\sum_{j=-1}^k a_j = 0, \quad \sum_{j=-1}^k j a_j + \sum_{j=-1}^k b_j = 0.$$

The method is of order  $p$  if and only if

$$\sum_{j=-1}^k (-j)^m a_j - m \sum_{j=-1}^k (-j)^{m-1} b_j = 0, \quad m = 0, 1, \dots, p.$$

## Example

Construct the explicit 2-step method of highest order!

There are 4 undetermined coefficients:  $a_0, a_1, b_0, b_1$ . The first four order conditions are

$$\begin{aligned} 1 + a_0 + a_1 &= 0, & 1 - a_1 - b_0 - b_1 &= 0, \\ 1 + a_1 + 2b_1 &= 0, & 1 - a_1 - 3b_1 &= 0, \end{aligned}$$

# Initial Value Problems: Linear Multistep Methods

## Theorem

A linear multistep is consistent if and only if

$$\sum_{j=-1}^k a_j = 0, \quad \sum_{j=-1}^k j a_j + \sum_{j=-1}^k b_j = 0.$$

The method is of order  $p$  if and only if

$$\sum_{j=-1}^k (-j)^m a_j - m \sum_{j=-1}^k (-j)^{m-1} b_j = 0, \quad m = 0, 1, \dots, p.$$

## Example

Construct the explicit 2-step method of highest order!

There are 4 undetermined coefficients:  $a_0, a_1, b_0, b_1$ . The first four order conditions are

$$\begin{aligned} 1 + a_0 + a_1 &= 0, & 1 - a_1 - b_0 - b_1 &= 0, \\ 1 + a_1 + 2b_1 &= 0, & 1 - a_1 - 3b_1 &= 0, \end{aligned}$$

which gives

$$a_0 = 4, a_1 = -5, b_0 = 4, b_1 = 2.$$

# Initial Value Problems: Linear Multistep Methods

## Theorem

A linear multistep is consistent if and only if

$$\sum_{j=-1}^k a_j = 0, \quad \sum_{j=-1}^k j a_j + \sum_{j=-1}^k b_j = 0.$$

The method is of order  $p$  if and only if

$$\sum_{j=-1}^k (-j)^m a_j - m \sum_{j=-1}^k (-j)^{m-1} b_j = 0, \quad m = 0, 1, \dots, p.$$

## Example

Construct the explicit 2-step method of highest order!

There are 4 undetermined coefficients:  $a_0, a_1, b_0, b_1$ . The first four order conditions are

$$\begin{aligned} 1 + a_0 + a_1 &= 0, & 1 - a_1 - b_0 - b_1 &= 0, \\ 1 + a_1 + 2b_1 &= 0, & 1 - a_1 - 3b_1 &= 0, \end{aligned}$$

which gives

$$a_0 = 4, a_1 = -5, b_0 = 4, b_1 = 2.$$

Thus, the unique explicit 2-step method of order 3 is  $y_{n+1} + 4y_n - 5y_{n-1} = h(4f_n + 2f_{n-1})$ .

# Initial Value Problems: Linear Multistep Methods

## Example

Construct the explicit 2-step method of highest order!

There are 4 undetermined coefficients:  $a_0, a_1, b_0, b_1$ . The first four order conditions are

$$\begin{aligned} 1 + a_0 + a_1 &= 0, & 1 - a_1 - b_0 - b_1 &= 0, \\ 1 + a_1 + 2b_1 &= 0, & 1 - a_1 - 3b_1 &= 0, \end{aligned}$$

which gives

$$a_0 = 4, a_1 = -5, b_0 = 4, b_1 = 2.$$

Thus, the unique explicit 2-step method of order 3 is  $y_{n+1} + 4y_n - 5y_{n-1} = h(4f_n + 2f_{n-1})$ .

Does this method converge?

# *Numerical Analysis & Scientific Computing II*

## *Lesson 2*

# *Initial Value Problems*

*2.4 Implicit method*

*2.5 Stiffness*

**2.6 Linear Multistep Methods**

**- Convergence**



*Akash Anand*  
MATH, IIT KANPUR



# Initial Value Problems: Linear Multistep Methods

## Example

Construct the explicit 2-step method of highest order!

There are 4 undetermined coefficients:  $a_0, a_1, b_0, b_1$ . The first four order conditions are

$$\begin{aligned} 1 + a_0 + a_1 &= 0, & 1 - a_1 - b_0 - b_1 &= 0, \\ 1 + a_1 + 2b_1 &= 0, & 1 - a_1 - 3b_1 &= 0, \end{aligned}$$

which gives

$$a_0 = 4, a_1 = -5, b_0 = 4, b_1 = 2.$$

Thus, the unique explicit 2-step method of order 3 is  $y_{n+1} + 4y_n - 5y_{n-1} = h(4f_n + 2f_{n-1})$ .

Does this method converge?

## Convergence

A linear multistep method is **convergent** if whenever the initial values  $y_n$  are chosen such that  $\max_{0 \leq n \leq k} |e_n| \rightarrow 0$ , as  $h \rightarrow 0$ , then  $\max_{0 \leq n \leq N} |e_n| \rightarrow 0$ .

# Initial Value Problems: Linear Multistep Methods

## Convergence

A linear multistep method is **convergent** if whenever the initial values  $y_n$  are chosen such that  $\max_{0 \leq n \leq k} |e_n| \rightarrow 0$ , as  $h \rightarrow 0$ , then  $\max_{0 \leq n \leq N} |e_n| \rightarrow 0$ .

## Example

Does

$$y_{n+1} + 4y_n - 5y_{n-1} = h(4f_n + 2f_{n-1}).$$

this method converge?

# Initial Value Problems: Linear Multistep Methods

## Convergence

A linear multistep method is **convergent** if whenever the initial values  $y_n$  are chosen such that  $\max_{0 \leq n \leq k} |e_n| \rightarrow 0$ , as  $h \rightarrow 0$ , then  $\max_{0 \leq n \leq N} |e_n| \rightarrow 0$ .

## Example

Does

$$y_{n+1} + 4y_n - 5y_{n-1} = h(4f_n + 2f_{n-1}).$$

this method converge?

Apply the method to the IVP  $y' = 0$ ,  $y(0) = 0$ .

# Initial Value Problems: Linear Multistep Methods

## Convergence

A linear multistep method is **convergent** if whenever the initial values  $y_n$  are chosen such that  $\max_{0 \leq n \leq k} |e_n| \rightarrow 0$ , as  $h \rightarrow 0$ , then  $\max_{0 \leq n \leq N} |e_n| \rightarrow 0$ .

## Example

Does

$$y_{n+1} + 4y_n - 5y_{n-1} = h(4f_n + 2f_{n-1}).$$

this method converge?

Apply the method to the IVP  $y' = 0$ ,  $y(0) = 0$ . We have

$$y_{n+1} = -4y_n + 5y_{n-1}.$$

# Initial Value Problems: Linear Multistep Methods

## Convergence

A linear multistep method is **convergent** if whenever the initial values  $y_n$  are chosen such that  $\max_{0 \leq n \leq k} |e_n| \rightarrow 0$ , as  $h \rightarrow 0$ , then  $\max_{0 \leq n \leq N} |e_n| \rightarrow 0$ .

## Example

Does

$$y_{n+1} + 4y_n - 5y_{n-1} = h(4f_n + 2f_{n-1}).$$

this method converge?

Apply the method to the IVP  $y' = 0$ ,  $y(0) = 0$ . We have

$$y_{n+1} = -4y_n + 5y_{n-1}.$$

To solve for  $y_n$ , we need to solve the difference equation.

# Initial Value Problems: Linear Multistep Methods

## Convergence

A linear multistep method is **convergent** if whenever the initial values  $y_n$  are chosen such that  $\max_{0 \leq n \leq k} |e_n| \rightarrow 0$ , as  $h \rightarrow 0$ , then  $\max_{0 \leq n \leq N} |e_n| \rightarrow 0$ .

## Example

Does

$$y_{n+1} + 4y_n - 5y_{n-1} = h(4f_n + 2f_{n-1}).$$

this method converge?

Apply the method to the IVP  $y' = 0$ ,  $y(0) = 0$ . We have

$$y_{n+1} = -4y_n + 5y_{n-1}.$$

To solve for  $y_n$ , we need to solve the difference equation. Toward this, introduce

$$v_n = \begin{bmatrix} y_{n+1} \\ y_n \end{bmatrix}$$

to obtain the equation  $v_{n+1} = Av_n$ , where

$$A = \begin{bmatrix} -4 & 5 \\ 1 & 0 \end{bmatrix}$$

# Initial Value Problems: Linear Multistep Methods

## Convergence

A linear multistep method is **convergent** if whenever the initial values  $y_n$  are chosen such that  $\max_{0 \leq n \leq k} |e_n| \rightarrow 0$ , as  $h \rightarrow 0$ , then  $\max_{0 \leq n \leq N} |e_n| \rightarrow 0$ .

## Example

Does

$$y_{n+1} + 4y_n - 5y_{n-1} = h(4f_n + 2f_{n-1}).$$

this method converge?

Apply the method to the IVP  $y' = 0$ ,  $y(0) = 0$ . We have

$$y_{n+1} = -4y_n + 5y_{n-1}.$$

To solve for  $y_n$ , we need to solve the difference equation. Toward this, introduce

$$v_n = \begin{bmatrix} y_{n+1} \\ y_n \end{bmatrix}$$

to obtain the equation  $v_{n+1} = Av_n$ , where

$$A = \begin{bmatrix} -4 & 5 \\ 1 & 0 \end{bmatrix} = \frac{1}{6} \begin{bmatrix} 1 & 5 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & -5 \end{bmatrix} \begin{bmatrix} 1 & 5 \\ 1 & -1 \end{bmatrix}.$$

# Initial Value Problems: Linear Multistep Methods

## Convergence

A linear multistep method is **convergent** if whenever the initial values  $y_n$  are chosen such that  $\max_{0 \leq n \leq k} |e_n| \rightarrow 0$ , as  $h \rightarrow 0$ , then  $\max_{0 \leq n \leq N} |e_n| \rightarrow 0$ .

## Example

Does

$$y_{n+1} + 4y_n - 5y_{n-1} = h(4f_n + 2f_{n-1}).$$

this method converge?

Apply the method to the IVP  $y' = 0$ ,  $y(0) = 0$ . We have

$$y_{n+1} = -4y_n + 5y_{n-1}.$$

To solve for  $y_n$ , we need to solve the difference equation. Toward this, introduce

$$v_n = \begin{bmatrix} y_{n+1} \\ y_n \end{bmatrix}$$

to obtain the equation  $v_{n+1} = Av_n$ , where

$$A = \begin{bmatrix} -4 & 5 \\ 1 & 0 \end{bmatrix} = \frac{1}{6} \begin{bmatrix} 1 & 5 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & -5 \end{bmatrix} \begin{bmatrix} 1 & 5 \\ 1 & -1 \end{bmatrix}.$$

Using,  $v_n = A^n v_0$  and  $y_0 = 0$ , we get

$$y_n = (1 - (-5)^n)y_1/6$$



# Initial Value Problems: Linear Multistep Methods

## Convergence

A linear multistep method is **convergent** if whenever the initial values  $y_n$  are chosen such that  $\max_{0 \leq n \leq k} |e_n| \rightarrow 0$ , as  $h \rightarrow 0$ , then  $\max_{0 \leq n \leq N} |e_n| \rightarrow 0$ .

## Example

Does

$$y_{n+1} + 4y_n - 5y_{n-1} = h(4f_n + 2f_{n-1}).$$

this method converge?

Apply the method to the IVP  $y' = 0$ ,  $y(0) = 0$ . We have

$$y_{n+1} = -4y_n + 5y_{n-1}.$$

Solving the difference equation yields

$$y_n = (1 - (-5)^n)y_1/6.$$

# Initial Value Problems: Linear Multistep Methods

## Convergence

A linear multistep method is **convergent** if whenever the initial values  $y_n$  are chosen such that  $\max_{0 \leq n \leq k} |e_n| \rightarrow 0$ , as  $h \rightarrow 0$ , then  $\max_{0 \leq n \leq N} |e_n| \rightarrow 0$ .

## Example

Does

$$y_{n+1} + 4y_n - 5y_{n-1} = h(4f_n + 2f_{n-1}).$$

this method converge?

Apply the method to the IVP  $y' = 0$ ,  $y(0) = 0$ . We have

$$y_{n+1} = -4y_n + 5y_{n-1}.$$

Solving the difference equation yields

$$y_n = (1 - (-5)^n)y_1/6.$$

If  $h = 1/N$ , then

$$e_N = y_N - y(1)$$

# Initial Value Problems: Linear Multistep Methods

## Convergence

A linear multistep method is **convergent** if whenever the initial values  $y_n$  are chosen such that  $\max_{0 \leq n \leq k} |e_n| \rightarrow 0$ , as  $h \rightarrow 0$ , then  $\max_{0 \leq n \leq N} |e_n| \rightarrow 0$ .

## Example

Does

$$y_{n+1} + 4y_n - 5y_{n-1} = h(4f_n + 2f_{n-1}).$$

this method converge?

Apply the method to the IVP  $y' = 0$ ,  $y(0) = 0$ . We have

$$y_{n+1} = -4y_n + 5y_{n-1}.$$

Solving the difference equation yields

$$y_n = (1 - (-5)^n)y_1/6.$$

If  $h = 1/N$ , then

$$e_N = y_N - y(1) = (1 - (-5)^N)y_1/6$$

# Initial Value Problems: Linear Multistep Methods

## Convergence

A linear multistep method is **convergent** if whenever the initial values  $y_n$  are chosen such that  $\max_{0 \leq n \leq k} |e_n| \rightarrow 0$ , as  $h \rightarrow 0$ , then  $\max_{0 \leq n \leq N} |e_n| \rightarrow 0$ .

## Example

Does

$$y_{n+1} + 4y_n - 5y_{n-1} = h(4f_n + 2f_{n-1}).$$

this method converge?

Apply the method to the IVP  $y' = 0$ ,  $y(0) = 0$ . We have

$$y_{n+1} = -4y_n + 5y_{n-1}.$$

Solving the difference equation yields

$$y_n = (1 - (-5)^n)y_1/6.$$

If  $h = 1/N$ , then

$$e_N = y_N - y(1) = (1 - (-5)^N)y_1/6 = (1 - (-5)^{1/h})y_1/6$$

# Initial Value Problems: Linear Multistep Methods

## Convergence

A linear multistep method is **convergent** if whenever the initial values  $y_n$  are chosen such that  $\max_{0 \leq n \leq k} |e_n| \rightarrow 0$ , as  $h \rightarrow 0$ , then  $\max_{0 \leq n \leq N} |e_n| \rightarrow 0$ .

## Example

Does

$$y_{n+1} + 4y_n - 5y_{n-1} = h(4f_n + 2f_{n-1}).$$

this method converge?

Apply the method to the IVP  $y' = 0$ ,  $y(0) = 0$ . We have

$$y_{n+1} = -4y_n + 5y_{n-1}.$$

Solving the difference equation yields

$$y_n = (1 - (-5)^n)y_1/6.$$

If  $h = 1/N$ , then

$$e_N = y_N - y(1) = (1 - (-5)^N)y_1/6 = (1 - (-5)^{1/h})y_1/6$$

For  $y_1 = h$ , we have  $e_1 = h$ . We see that even as  $e_1 \rightarrow 0$  as  $h \rightarrow 0$ ,  $e_N \nrightarrow 0$ .

# Initial Value Problems: Linear Multistep Methods

## Convergence

A linear multistep method is **convergent** if whenever the initial values  $y_n$  are chosen such that  $\max_{0 \leq n \leq k} |e_n| \rightarrow 0$ , as  $h \rightarrow 0$ , then  $\max_{0 \leq n \leq N} |e_n| \rightarrow 0$ .

## Example

Does

$$y_{n+1} + 4y_n - 5y_{n-1} = h(4f_n + 2f_{n-1}).$$

this method converge?

Apply the method to the IVP  $y' = 0$ ,  $y(0) = 0$ . We have

$$y_{n+1} = -4y_n + 5y_{n-1}.$$

Solving the difference equation yields

$$y_n = (1 - (-5)^n)y_1/6.$$

If  $h = 1/N$ , then

$$e_N = y_N - y(1) = (1 - (-5)^N)y_1/6 = (1 - (-5)^{1/h})y_1/6$$

For  $y_1 = h$ , we have  $e_1 = h$ . We see that even as  $e_1 \rightarrow 0$  as  $h \rightarrow 0$ ,  $e_N \nrightarrow 0$ . Thus, the method does not converge.

# Initial Value Problems: Linear Multistep Methods

## Convergence

A linear multistep method is **convergent** if whenever the initial values  $y_n$  are chosen such that  $\max_{0 \leq n \leq k} |e_n| \rightarrow 0$ , as  $h \rightarrow 0$ , then  $\max_{0 \leq n \leq N} |e_n| \rightarrow 0$ .

## Example

Does

$$y_{n+1} + 4y_n - 5y_{n-1} = h(4f_n + 2f_{n-1}).$$

this method converge?

Apply the method to the IVP  $y' = 0$ ,  $y(0) = 0$ . We have

$$y_{n+1} = -4y_n + 5y_{n-1}.$$

Solving the difference equation yields

$$y_n = (1 - (-5)^n)y_1/6.$$

If  $h = 1/N$ , then

$$e_N = y_N - y(1) = (1 - (-5)^N)y_1/6 = (1 - (-5)^{1/h})y_1/6$$

For  $y_1 = h$ , we have  $e_1 = h$ . We see that even as  $e_1 \rightarrow 0$  as  $h \rightarrow 0$ ,  $e_N \not\rightarrow 0$ . Thus, the method does not converge.

Note that if we take exact starting values  $y_0 = y_1 = 0$ , then  $y_n = 0$  for all  $n$ .

# Initial Value Problems: Linear Multistep Methods

## Convergence

A linear multistep method is **convergent** if whenever the initial values  $y_n$  are chosen such that  $\max_{0 \leq n \leq k} |e_n| \rightarrow 0$ , as  $h \rightarrow 0$ , then  $\max_{0 \leq n \leq N} |e_n| \rightarrow 0$ .

## Example

Does

$$y_{n+1} + 4y_n - 5y_{n-1} = h(4f_n + 2f_{n-1}).$$

this method converge?

Apply the method to the IVP  $y' = 0$ ,  $y(0) = 0$ . We have

$$y_{n+1} = -4y_n + 5y_{n-1}.$$

Solving the difference equation yields

$$y_n = (1 - (-5)^n)y_1/6.$$

If  $h = 1/N$ , then

$$e_N = y_N - y(1) = (1 - (-5)^N)y_1/6 = (1 - (-5)^{1/h})y_1/6$$

For  $y_1 = h$ , we have  $e_1 = h$ . We see that even as  $e_1 \rightarrow 0$  as  $h \rightarrow 0$ ,  $e_N \not\rightarrow 0$ . Thus, the method does not converge.

Note that if we take exact starting values  $y_0 = y_1 = 0$ , then  $y_n = 0$  for all  $n$ . Thus, a perturbation of size  $\varepsilon$  in the starting values leads to a difference of size roughly  $5^{1/h}\varepsilon$  in the discrete solution.



# Initial Value Problems: Linear Multistep Methods

## Convergence

A linear multistep method is **convergent** if whenever the initial values  $y_n$  are chosen such that  $\max_{0 \leq n \leq k} |e_n| \rightarrow 0$ , as  $h \rightarrow 0$ , then  $\max_{0 \leq n \leq N} |e_n| \rightarrow 0$ .

## Example

Does

$$y_{n+1} + 4y_n - 5y_{n-1} = h(4f_n + 2f_{n-1}).$$

this method converge?

Apply the method to the IVP  $y' = 0$ ,  $y(0) = 0$ . We have

$$y_{n+1} = -4y_n + 5y_{n-1}.$$

Solving the difference equation yields

$$y_n = (1 - (-5)^n)y_1/6.$$

If  $h = 1/N$ , then

$$e_N = y_N - y(1) = (1 - (-5)^N)y_1/6 = (1 - (-5)^{1/h})y_1/6$$

For  $y_1 = h$ , we have  $e_1 = h$ . We see that even as  $e_1 \rightarrow 0$  as  $h \rightarrow 0$ ,  $e_N \not\rightarrow 0$ . Thus, the method does not converge.

Note that if we take exact starting values  $y_0 = y_1 = 0$ , then  $y_n = 0$  for all  $n$ . Thus, a perturbation of size  $\varepsilon$  in the starting values leads to a difference of size roughly  $5^{1/h}\varepsilon$  in the discrete solution. The method is, therefore, not stable.

# *Numerical Analysis & Scientific Computing II*

## *Module 2*

# *Initial Value Problems*

*2.4 Implicit method*

*2.5 Stiffness*

**2.6 Linear Multistep Methods**  
**- Stability**



*Akash Anand*  
MATH, IIT KANPUR

# Initial Value Problems: Linear Multistep Methods

## Stability

A linear  $k + 1$  step method is **stable** if for any initial value problem with Lipschitz continuous  $f$  and of  $\varepsilon > 0$ , there exists  $\delta, h_0 > 0$  such that if  $h \leq h_0$  and two choices of starting values  $y_j$  and  $\hat{y}_j$  are chosen satisfying

$$\max_{0 \leq j \leq k} |y_j - \hat{y}_j| \leq \delta,$$

then the corresponding approximate solutions satisfy

$$\max_{0 \leq j \leq N} |y_j - \hat{y}_j| \leq \varepsilon.$$

# Initial Value Problems: Linear Multistep Methods

## Stability

A linear  $k + 1$  step method is **stable** if for any initial value problem with Lipschitz continuous  $f$  and of  $\varepsilon > 0$ , there exists  $\delta, h_0 > 0$  such that if  $h \leq h_0$  and two choices of starting values  $y_j$  and  $\hat{y}_j$  are chosen satisfying

$$\max_{0 \leq j \leq k} |y_j - \hat{y}_j| \leq \delta,$$

then the corresponding approximate solutions satisfy

$$\max_{0 \leq j \leq N} |y_j - \hat{y}_j| \leq \varepsilon.$$

If  $y' = 0$ , then the general linear multistep method becomes

$$y_{n+1} + \sum_{j=0}^k a_j y_{n-j} = 0, \quad n = k, k+1, \dots$$

This is an example of a homogeneous linear difference equation of order  $k + 1$  with constant coefficients.

# Initial Value Problems: Linear Multistep Methods

## Stability

A linear  $k + 1$  step method is **stable** if for any initial value problem with Lipschitz continuous  $f$  and of  $\varepsilon > 0$ , there exists  $\delta, h_0 > 0$  such that if  $h \leq h_0$  and two choices of starting values  $y_j$  and  $\hat{y}_j$  are chosen satisfying

$$\max_{0 \leq j \leq k} |y_j - \hat{y}_j| \leq \delta,$$

then the corresponding approximate solutions satisfy

$$\max_{0 \leq j \leq N} |y_j - \hat{y}_j| \leq \varepsilon.$$

If  $y' = 0$ , then the general linear multistep method becomes

$$y_{n+1} + \sum_{j=0}^k a_j y_{n-j} = 0, \quad n = k, k+1, \dots$$

This is an example of a homogeneous linear difference equation of order  $k + 1$  with constant coefficients.

If a method is stable, it should, in particular, be stable while computing the zero solution of the initial value problem  $y' = 0, y(0) = 0$ . To investigate this, we need to solve the homogeneous linear difference equation of order  $k + 1$ .

# Initial Value Problems: Linear Multistep Methods

## Stability

A linear  $k + 1$  step method is **stable** if for any initial value problem with Lipschitz continuous  $f$  and of  $\varepsilon > 0$ , there exists  $\delta, h_0 > 0$  such that if  $h \leq h_0$  and two choices of starting values  $y_j$  and  $\hat{y}_j$  are chosen satisfying

$$\max_{0 \leq j \leq k} |y_j - \hat{y}_j| \leq \delta,$$

then the corresponding approximate solutions satisfy

$$\max_{0 \leq j \leq N} |y_j - \hat{y}_j| \leq \varepsilon.$$

If  $y' = 0$ , then the general linear multistep method becomes

$$y_{n+1} + \sum_{j=0}^k a_j y_{n-j} = 0, \quad n = k, k+1, \dots$$

This is an example of a homogeneous linear difference equation of order  $k + 1$  with constant coefficients.

If a method is stable, it should, in particular, be stable while computing the zero solution of the initial value problem  $y' = 0, y(0) = 0$ . To investigate this, we need to solve the homogeneous linear difference equation of order  $k + 1$ .

To find the general solution, we first try for a solution of the form  $(\lambda^n)_{n=0}^{\infty}$ . Substituting this in the difference equation, we see that it is a solution if and only if  $\lambda$  is a root of the characteristic polynomial

$$\rho(t) = t^{k+1} + \sum_{j=0}^k a_j t^{k-j}.$$

# Initial Value Problems: Linear Multistep Methods

If  $y' = 0$ , then the general linear multistep method becomes

$$y_{n+1} + \sum_{j=0}^k a_j y_{n-j} = 0, \quad n = k, k+1, \dots$$

This is an example of a homogeneous linear difference equation of order  $k+1$  with constant coefficients.

If a method is stable, it should, in particular, be stable while computing the zero solution of the initial value problem  $y' = 0, y(0) = 0$ . To investigate this, we need to solve the homogeneous linear difference equation of order  $k+1$ .

To find the general solution, we first try for a solution of the form  $(\lambda^n)_{n=0}^{\infty}$ . Substituting this in the difference equation, we see that it is a solution if and only if  $\lambda$  is a root of the characteristic polynomial

$$\rho(t) = t^{k+1} + \sum_{j=0}^k a_j t^{k-j}.$$

If there are  $k+1$  distinct roots  $\lambda_i, i = 0, \dots, k$ , then we have a full basis of  $k+1$  linearly independent solutions (note that the solutions form a vector space).

# Initial Value Problems: Linear Multistep Methods

*If  $y' = 0$ , then the general linear multistep method becomes*

$$y_{n+1} + \sum_{j=0}^k a_j y_{n-j} = 0, \quad n = k, k+1, \dots$$

*This is an example of a homogeneous linear difference equation of order  $k+1$  with constant coefficients.*

*If a method is stable, it should, in particular, be stable while computing the zero solution of the initial value problem  $y' = 0, y(0) = 0$ . To investigate this, we need to solve the homogeneous linear difference equation of order  $k+1$ .*

*To find the general solution, we first try for a solution of the form  $(\lambda^n)_{n=0}^\infty$ . Substituting this in the difference equation, we see that it is a solution if and only if  $\lambda$  is a root of the characteristic polynomial*

$$\rho(t) = t^{k+1} + \sum_{j=0}^k a_j t^{k-j}.$$

*If there are  $k+1$  distinct roots  $\lambda_i, i = 0, \dots, k$ , then we have a full basis of  $k+1$  linearly independent solutions (note that the solutions form a vector space).*

*In case of multiple roots, this does not give the complete set of solutions.*



# Initial Value Problems: Linear Multistep Methods

If  $y' = 0$ , then the general linear multistep method becomes

$$y_{n+1} + \sum_{j=0}^k a_j y_{n-j} = 0, \quad n = k, k+1, \dots$$

This is an example of a homogeneous linear difference equation of order  $k+1$  with constant coefficients.

If a method is stable, it should, in particular, be stable while computing the zero solution of the initial value problem  $y' = 0, y(0) = 0$ . To investigate this, we need to solve the homogeneous linear difference equation of order  $k+1$ .

To find the general solution, we first try for a solution of the form  $(\lambda^n)_{n=0}^\infty$ . Substituting this in the difference equation, we see that it is a solution if and only if  $\lambda$  is a root of the characteristic polynomial

$$\rho(t) = t^{k+1} + \sum_{j=0}^k a_j t^{k-j}.$$

If there are  $k+1$  distinct roots  $\lambda_i, i = 0, \dots, k$ , then we have a full basis of  $k+1$  linearly independent solutions (note that the solutions form a vector space).

In case of multiple roots, this does not give the complete set of solutions.

If  $\lambda$  is a double root, then  $(\lambda^n)_{n=0}^\infty$  and  $(n\lambda^n)_{n=0}^\infty$  both are solutions.

# Initial Value Problems: Linear Multistep Methods

If  $y' = 0$ , then the general linear multistep method becomes

$$y_{n+1} + \sum_{j=0}^k a_j y_{n-j} = 0, \quad n = k, k+1, \dots$$

This is an example of a homogeneous linear difference equation of order  $k+1$  with constant coefficients.

If a method is stable, it should, in particular, be stable while computing the zero solution of the initial value problem  $y' = 0, y(0) = 0$ . To investigate this, we need to solve the homogeneous linear difference equation of order  $k+1$ .

To find the general solution, we first try for a solution of the form  $(\lambda^n)_{n=0}^\infty$ . Substituting this in the difference equation, we see that it is a solution if and only if  $\lambda$  is a root of the characteristic polynomial

$$\rho(t) = t^{k+1} + \sum_{j=0}^k a_j t^{k-j}.$$

If there are  $k+1$  distinct roots  $\lambda_i, i = 0, \dots, k$ , then we have a full basis of  $k+1$  linearly independent solutions (note that the solutions form a vector space).

In case of multiple roots, this does not give the complete set of solutions.

If  $\lambda$  is a double root, then  $(\lambda^n)_{n=0}^\infty$  and  $(n\lambda^n)_{n=0}^\infty$  both are solutions. Similarly,  $\lambda$  is a root of multiplicity  $M > 2$ , then  $(n^m \lambda^n)_{n=0}^\infty, m = 0, 1, \dots, M-1$ , also satisfy the difference equation.

# Initial Value Problems: Linear Multistep Methods

If  $y' = 0$ , then the general linear multistep method becomes

$$y_{n+1} + \sum_{j=0}^k a_j y_{n-j} = 0, \quad n = k, k+1, \dots$$

This is an example of a homogeneous linear difference equation of order  $k+1$  with constant coefficients.

If a method is stable, it should, in particular, be stable while computing the zero solution of the initial value problem  $y' = 0, y(0) = 0$ . To investigate this, we need to solve the homogeneous linear difference equation of order  $k+1$ .

To find the general solution, we first try for a solution of the form  $(\lambda^n)_{n=0}^\infty$ . Substituting this in the difference equation, we see that it is a solution if and only if  $\lambda$  is a root of the characteristic polynomial

$$\rho(t) = t^{k+1} + \sum_{j=0}^k a_j t^{k-j}.$$

If there are  $k+1$  distinct roots  $\lambda_i, i = 0, \dots, k$ , then we have a full basis of  $k+1$  linearly independent solutions (note that the solutions form a vector space).

In case of multiple roots, this does not give the complete set of solutions.

If  $\lambda$  is a double root, then  $(\lambda^n)_{n=0}^\infty$  and  $(n\lambda^n)_{n=0}^\infty$  both are solutions. Similarly,  $\lambda$  is a root of multiplicity  $M > 2$ , then  $(n^m \lambda^n)_{n=0}^\infty, m = 0, 1, \dots, M-1$ , also satisfy the difference equation.

Thus, for  $\rho(t) = \prod_{j=1}^J (t - \lambda_j)^{M_j}$  where  $\sum_{j=1}^J M_j = k+1$ , the general solution is  $y_n = \sum_{j=1}^J \sum_{m=0}^{M_j-1} c_{jm} n^m \lambda_j^n$ .