

Numerical Analysis & Scientific Computing II

Module 2

Initial Value Problems

2.0 First-order system of ODE

2.1 Well-posedness

2.2 Stability

2.3 Euler's method



Akash Anand
MATH, IIT KANPUR

Numerical Analysis & Scientific Computing II

Module 2

Initial Value Problems

2.0 First-order system of ODE

2.1 Well-posedness

2.2 Stability

2.3 Euler's method



Initial Value Problems: First order system of ODE



If we learn to solve first order system of ODEs, then we can also solve a high order ODE provided it is explicit.

Initial Value Problems: First order system of ODE

If we learn to solve first order system of ODEs, then we can also solve a high order ODE provided it is explicit.

Recall that a k th order ODE is said to be explicit if it can be written in the form

$$u^{(k)} = f\left(t, u, u', u'', \dots, u^{(k-1)}\right)$$

where $f : \mathbb{R}^{kn+1} \rightarrow \mathbb{R}^n$.

Initial Value Problems: First order system of ODE

If we learn to solve first order system of ODEs, then we can also solve a high order ODE provided it is explicit.

Recall that a k th order ODE is said to be explicit if it can be written in the form

$$u^{(k)} = f \left(t, u, u', u'', \dots, u^{(k-1)} \right)$$

where $f : \mathbb{R}^{kn+1} \rightarrow \mathbb{R}^n$.

Introduce new variables

$$y_1(t) = u(t), y_2(t) = u'(t), \dots, y_k(t) = u^{(k-1)}(t)$$

so that the original k th order system becomes a system of kn first order equations

$$y' = \begin{bmatrix} y_1' \\ \vdots \\ y_{k-1}' \\ y_k' \end{bmatrix} = \begin{bmatrix} y_2 \\ \vdots \\ y_k \\ f(t, y_1, y_2, \dots, y_k) \end{bmatrix}$$

Numerical Analysis & Scientific Computing II

Module 2

Initial Value Problems

2.0 First-order system of ODE

2.1 Well-posedness

2.2 Stability

2.3 Euler's method



Akash Anand
MATH, IIT KANPUR

Initial Value Problems: Well-posedness



We want to study the methods for numerical solution of a first order system of ordinary differential equations with initial conditions,

$$y' = f(t, y), \quad y(t_0) = y_0,$$

where $y: [a, b] \rightarrow \mathbb{R}^n$ and $f: \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n$.

Initial Value Problems: Well-posedness



We want to study the methods for numerical solution of a first order system of ordinary differential equations with initial conditions,

$$y' = f(t, y), \quad y(t_0) = y_0,$$

where $y: [a, b] \rightarrow \mathbb{R}^n$ and $f: \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n$.

Before we study the numerical solution, we need to investigate the well-posedness of the problem, that is, we study if the problem has following three properties:

- (i) existence of a solution (**existence**),
- (ii) uniqueness of the solution (**uniqueness**), and
- (iii) continuous dependence of the solution of the data (**conditioning**).

Initial Value Problems: Well-posedness



We want to study the methods for numerical solution of a first order system of ordinary differential equations with initial conditions,

$$y' = f(t, y), \quad y(t_0) = y_0,$$

where $y: [a, b] \rightarrow \mathbb{R}^n$ and $f: \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n$.

Before we study the numerical solution, we need to investigate the well-posedness of the problem, that is, we study if the problem has following three properties:

- (i) existence of a solution (**existence**),
- (ii) uniqueness of the solution (**uniqueness**), and
- (iii) continuous dependence of the solution of the data (**conditioning**).

This investigation, however, is the main subject matter of the course on Ordinary Differential Equations (ODE) and we, in this course, will only recall the relevant discussions.

Initial Value Problems: Well-posedness



Initial Value Problems: Well-posedness



Let $D = [a, b] \times \Omega \subseteq \mathbb{R}^{n+1}$ be a closed and bounded set.

Suppose that $f: \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n$ is a **Lipschitz continuous function in y** on D , that is, there is a constant L such that for any $t \in [a, b]$ and for any y and $\hat{y} \in \Omega$,

$$\|f(t, \hat{y}) - f(t, y)\| \leq L \|\hat{y} - y\|.$$

Initial Value Problems: Well-posedness



Let $D = [a, b] \times \Omega \subseteq \mathbb{R}^{n+1}$ be a closed and bounded set.

Suppose that $f: \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n$ is a **Lipschitz continuous function in y** on D , that is, there is a constant L such that for any $t \in [a, b]$ and for any y and $\hat{y} \in \Omega$,

$$\|f(t, \hat{y}) - f(t, y)\| \leq L \|\hat{y} - y\|.$$

Recall, from your ODE course, that for such functions, the following Initial Value Problem (IVP)

$$y' = f(t, y), \quad y(t_0) = y_0,$$

has a unique solution in a subinterval of $[a, b]$ containing y_0 .

Initial Value Problems: Well-posedness



Let $D = [a, b] \times \Omega \subseteq \mathbb{R}^{n+1}$ be a closed and bounded set.

Suppose that $f: \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n$ is a **Lipschitz continuous function in y** on D , that is, there is a constant L such that for any $t \in [a, b]$ and for any y and $\hat{y} \in \Omega$,

$$\|f(t, \hat{y}) - f(t, y)\| \leq L \|\hat{y} - y\|.$$

Recall, from your ODE course, that for such functions, the following Initial Value Problem (IVP)

$$y' = f(t, y), \quad y(t_0) = y_0,$$

has a unique solution in a subinterval of $[a, b]$ containing y_0 .

Example: If f is differentiable, then f is Lipschitz continuous with

$$L = \max_{(t,y) \in D} \|f'(t, y)\|,$$

where f' is the $n \times n$ Jacobian matrix of f with respect to y , $[f'(t, y)]_{ij} = \frac{\partial f_i(t, y)}{\partial y_j}$.

Initial Value Problems: Well-posedness



Let $\hat{y}(t)$ be the solution of the following IVP with a perturbed initial condition

$$y' = f(t, y), \quad y(t_0) = \hat{y}_0.$$

Initial Value Problems: Well-posedness



Let $\hat{y}(t)$ be the solution of the following IVP with a perturbed initial condition

$$y' = f(t, y), \quad y(t_0) = \hat{y}_0.$$

Then, it can be shown that for any $t \geq t_0$,

$$\|\hat{y}(t) - y(t)\| \leq e^{L(t-t_0)} \|\hat{y}_0 - y_0\|.$$

Initial Value Problems: Well-posedness



Let $\hat{y}(t)$ be the solution of the following IVP with a perturbed initial condition

$$y' = f(t, y), \quad y(t_0) = \hat{y}_0.$$

Then, it can be shown that for any $t \geq t_0$,

$$\|\hat{y}(t) - y(t)\| \leq e^{L(t-t_0)} \|\hat{y}_0 - y_0\|.$$

If the function is also perturbed so that we have $\hat{y}' = \hat{f}(t, \hat{y})$, then ...

Initial Value Problems: Well-posedness

Let $\hat{y}(t)$ be the solution of the following IVP with a perturbed initial condition

$$y' = f(t, y), \quad y(t_0) = \hat{y}_0.$$

Then, it can be shown that for any $t \geq t_0$,

$$\|\hat{y}(t) - y(t)\| \leq e^{L(t-t_0)} \|\hat{y}_0 - y_0\|.$$

If the function is also perturbed so that we have $\hat{y}' = \hat{f}(t, \hat{y})$, then for $t \geq t_0$,

$$\|\hat{y}(t) - y(t)\| \leq e^{L(t-t_0)} \|\hat{y}_0 - y_0\| + \frac{e^{L(t-t_0)} - 1}{L} \|\hat{f} - f\|,$$

where

$$\|\hat{f} - f\| = \max_{(t,y) \in D} \|\hat{f}(t, y) - f(t, y)\|.$$

These perturbation bounds show that the unique solution to the IVP is a continuous function of the problem data, and hence the problem is well-posed.

Numerical Analysis & Scientific Computing II

Module 2

Initial Value Problems

2.0 First-order system of ODE

2.1 Well-posedness

2.2 Stability

2.3 Euler's method



Akash Anand
MATH, IIT KANPUR

Initial Value Problems: Stability



*We will see that the stability of the solution plays a significant role in the propagation of error, introduced by a numerical method, over time; they can either get **amplified** or **diminished** depending on the stability.*



We will see that the stability of the solution plays a significant role in the propagation of error, introduced by a numerical method, over time; they can either get **amplified** or **diminished** depending on the stability.

Also recall that a solution of the ODE $y' = f(t, y)$ is said to be **stable** if for every $\epsilon > 0$, there exists a $\delta > 0$ such that if $\hat{y}(t)$ satisfied the ODE and $\|\hat{y}(t_0) - y(t_0)\| \leq \delta$, then $\|\hat{y}(t) - y(t)\| \leq \epsilon$ for all $t \geq t_0$.



We will see that the stability of the solution plays a significant role in the propagation of error, introduced by a numerical method, over time; they can either get **amplified** or **diminished** depending on the stability.

Also recall that a solution of the ODE $y' = f(t, y)$ is said to be **stable** if for every $\epsilon > 0$, there exists a $\delta > 0$ such that if $\hat{y}(t)$ satisfied the ODE and $\|\hat{y}(t_0) - y(t_0)\| \leq \delta$, then $\|\hat{y}(t) - y(t)\| \leq \epsilon$ for all $t \geq t_0$.

Thus, for a **stable solution**, if the initial value is perturbed, then the perturbed solution remains close to the original solution, which **rules out the exponential divergence of perturbed solution** allowed by the perturbation bound

$$\|\hat{y}(t) - y(t)\| \leq e^{L(t-t_0)} \|\hat{y}_0 - y_0\|.$$



We will see that the stability of the solution plays a significant role in the propagation of error, introduced by a numerical method, over time; they can either get **amplified** or **diminished** depending on the stability.

Also recall that a solution of the ODE $y' = f(t, y)$ is said to be **stable** if for every $\epsilon > 0$, there exists a $\delta > 0$ such that if $\hat{y}(t)$ satisfied the ODE and $\|\hat{y}(t_0) - y(t_0)\| \leq \delta$, then $\|\hat{y}(t) - y(t)\| \leq \epsilon$ for all $t \geq t_0$.

Thus, for a **stable solution**, if the initial value is perturbed, then the perturbed solution remains close to the original solution, which **rules out the exponential divergence of perturbed solution** allowed by the perturbation bound

$$\|\hat{y}(t) - y(t)\| \leq e^{L(t-t_0)} \|\hat{y}_0 - y_0\|.$$

A stable solution is said to be **asymptotically stable** if $\|\hat{y}(t) - y(t)\| \rightarrow 0$ as $t \rightarrow \infty$. This stronger form means that the original and perturbed solution not only remain close to each other, but they converge toward each other over time.

Initial Value Problems: Stability



Examples:

(i) Solutions of $y' = b$, for a given b , are ...

Initial Value Problems: Stability



Examples:

(i) Solutions of $y' = b$, for a given b , are stable but not asymptotically stable.

Initial Value Problems: Stability



Examples:


- (i) Solutions of $y' = b$, for a given b , are stable but not asymptotically stable.
- (ii) Consider the IVP $y' = \lambda y$, for a given constant λ , $y(0) = y_0$. The solution is $y(t) = y_0 e^{\lambda t}$.

Initial Value Problems: Stability



Examples:

- (i) Solutions of $y' = b$, for a given b , are stable but not asymptotically stable.
- (ii) Consider the IVP $y' = \lambda y$, for a given constant λ , $y(0) = y_0$. The solution is $y(t) = y_0 e^{\lambda t}$.




Initial Value Problems: Stability

Examples:

(i) Solutions of $y' = b$, for a given b , are stable but not asymptotically stable.

(ii) Consider the IVP $y' = \lambda y$, for a given constant λ , $y(0) = y_0$. The solution is $y(t) = y_0 e^{\lambda t}$.

If $\text{Re}(\lambda) > 0$, then all non-zero solutions grow exponentially so that any two solutions diverge away, and hence every solution is unstable.





Examples:

(i) Solutions of $y' = b$, for a given b , are stable but not asymptotically stable.

(ii) Consider the IVP $y' = \lambda y$, for a given constant λ , $y(0) = y_0$. The solution is $y(t) = y_0 e^{\lambda t}$.

If $\text{Re}(\lambda) > 0$, then all non-zero solutions grow exponentially so that any two solutions diverge away, and hence every solution is unstable.



Initial Value Problems: Stability



Examples:

(i) Solutions of $y' = b$, for a given b , are stable but not asymptotically stable.

(ii) Consider the IVP $y' = \lambda y$, for a given constant λ , $y(0) = y_0$. The solution is $y(t) = y_0 e^{\lambda t}$.

If $\text{Re}(\lambda) > 0$, then all non-zero solutions grow exponentially so that any two solutions diverge away, and hence every solution is unstable.

If $\text{Re}(\lambda) < 0$, then all non-zero solutions decay exponentially so that any two solutions converge toward each other, and hence, every solution is not only stable but asymptotically stable.



Initial Value Problems: Stability



Examples:

(i) Solutions of $y' = b$, for a given b , are stable but not asymptotically stable.

(ii) Consider the IVP $y' = \lambda y$, for a given constant λ , $y(0) = y_0$. The solution is $y(t) = y_0 e^{\lambda t}$.

If $\operatorname{Re}(\lambda) > 0$, then all non-zero solutions grow exponentially so that any two solutions diverge away, and hence every solution is unstable.

If $\operatorname{Re}(\lambda) < 0$, then all non-zero solutions decay exponentially so that any two solutions converge toward each other, and hence, every solution is not only stable but asymptotically stable.

If $\operatorname{Re}(\lambda) = 0$, then ...

Initial Value Problems: Stability



Examples:

(i) Solutions of $y' = b$, for a given b , are stable but not asymptotically stable.

(ii) Consider the IVP $y' = \lambda y$, for a given constant λ , $y(0) = y_0$. The solution is $y(t) = y_0 e^{\lambda t}$.

If $\operatorname{Re}(\lambda) > 0$, then all non-zero solutions grow exponentially so that any two solutions diverge away, and hence every solution is unstable.

If $\operatorname{Re}(\lambda) < 0$, then all non-zero solutions decay exponentially so that any two solutions converge toward each other, and hence, every solution is not only stable but asymptotically stable.

If $\operatorname{Re}(\lambda) = 0$, then ... (**Exercise**)

Initial Value Problems: Stability

Examples:

(i) Solutions of $y' = b$, for a given b , are stable but not asymptotically stable.

(ii) Consider the IVP $y' = \lambda y$, for a given constant λ , $y(0) = y_0$. The solution is $y(t) = y_0 e^{\lambda t}$.

If $\operatorname{Re}(\lambda) > 0$, then all non-zero solutions grow exponentially so that any two solutions diverge away, and hence every solution is unstable.

If $\operatorname{Re}(\lambda) < 0$, then all non-zero solutions decay exponentially so that any two solutions converge toward each other, and hence, every solution is not only stable but asymptotically stable.

If $\operatorname{Re}(\lambda) = 0$, then ... (**Exercise**)

(iii) A linear homogeneous system of ODEs with constant coefficients has the form

$$y' = Ay,$$

where A is an $n \times n$ matrix. Suppose we have the initial condition $y(0) = y_0$. Discuss the stability of the solutions if

(a) A is diagonalizable, and

(b) A is not diagonalizable.

(**Exercise**)

Numerical Analysis & Scientific Computing II

Module 2

Initial Value Problems

2.0 First-order system of ODE

2.1 Well-posedness

2.2 Stability

2.3 Euler's method



Akash Anand
MATH, IIT KANPUR

Initial Value Problems: Euler's method



Starting at time t_0 with the given initial value y_0 , we would like to track the solution trajectory governed by the ODE

$$y' = f(t, y)$$

Initial Value Problems: Euler's method



Starting at time t_0 with the given initial value y_0 , we would like to track the solution trajectory governed by the ODE

$$y' = f(t, y)$$

The fundamental theorem of calculus tells us that

$$y(T) = y(t_0) + \int_{t_0}^T y'(s) ds$$

Initial Value Problems: Euler's method



Starting at time t_0 with the given initial value y_0 , we would like to track the solution trajectory governed by the ODE

$$y' = f(t, y)$$

The fundamental theorem of calculus tells us that

$$y(T) = y(t_0) + \int_{t_0}^T y'(s) ds = y(t_0) + \int_{t_0}^T f(s, y(s)) ds$$

Initial Value Problems: Euler's method



Starting at time t_0 with the given initial value y_0 , we would like to track the solution trajectory governed by the ODE

$$y' = f(t, y)$$

The fundamental theorem of calculus tells us that

$$y(T) = y(t_0) + \int_{t_0}^T y'(s) ds = y(t_0) + \int_{t_0}^T f(s, y(s)) ds$$

The simplest approximation for the integral is by using the left end-point rule,

Initial Value Problems: Euler's method

Starting at time t_0 with the given initial value y_0 , we would like to track the solution trajectory governed by the ODE


$$y' = f(t, y)$$

The fundamental theorem of calculus tells us that

$$y(T) = y(t_0) + \int_{t_0}^T y'(s) ds = y(t_0) + \int_{t_0}^T f(s, y(s)) ds$$

The simplest approximation for the integral is by using the left end-point rule, that is,

$$\int_a^b g(s) ds \approx (b - a)g(a)$$



Initial Value Problems: Euler's method

Starting at time t_0 with the given initial value y_0 , we would like to track the solution trajectory governed by the ODE

$$y' = f(t, y)$$

The fundamental theorem of calculus tells us that


$$y(T) = y(t_0) + \int_{t_0}^T y'(s) ds = y(t_0) + \int_{t_0}^T f(s, y(s)) ds$$

The simplest approximation for the integral is by using the left end-point rule, that is,

$$\int_a^b g(s) ds \approx (b - a)g(a)$$

yielding the method

$$y(T) = y(t_0) + (T - t_0)f(t_0, y_0)$$



Initial Value Problems: Euler's method

Starting at time t_0 with the given initial value y_0 , we would like to track the solution trajectory governed by the ODE

$$y' = f(t, y)$$

The fundamental theorem of calculus tells us that

$$y(T) = y(t_0) + \int_{t_0}^T y'(s) ds = y(t_0) + \int_{t_0}^T f(s, y(s)) ds$$


The simplest approximation for the integral is by using the left end-point rule, that is,

$$\int_a^b g(s) ds \approx (b - a)g(a)$$

yielding the method

$$y(T) = y(t_0) + (T - t_0)f(t_0, y_0)$$

To have more control over the error, we can also do this in multiple steps as follows:



Initial Value Problems: Euler's method

Starting at time t_0 with the given initial value y_0 , we would like to track the solution trajectory governed by the ODE

$$y' = f(t, y)$$

The fundamental theorem of calculus tells us that

$$y(T) = y(t_0) + \int_{t_0}^T y'(s) ds = y(t_0) + \int_{t_0}^T f(s, y(s)) ds$$


The simplest approximation for the integral is by using the left end-point rule, that is,

$$\int_a^b g(s) ds \approx (b - a)g(a)$$

yielding the method

$$y(T) = y(t_0) + (T - t_0)f(t_0, y_0)$$

To have more control over the error, we can also do this in multiple steps as follows:



Initial Value Problems: Euler's method

Starting at time t_0 with the given initial value y_0 , we would like to track the solution trajectory governed by the ODE

$$y' = f(t, y)$$

The fundamental theorem of calculus tells us that

$$y(t_{k+1}) = y(t_k) + \int_{t_k}^{t_{k+1}} y'(s) ds = y(t_k) + \int_{t_k}^{t_{k+1}} f(s, y(s)) ds$$


The simplest approximation for the integral is by using the left end-point rule, that is,

$$\int_a^b g(s) ds \approx (b - a)g(a)$$

yielding the method

$$y(T) = y(t_0) + (T - t_0)f(t_0, y_0)$$

To have more control over the error, we can also do this in multiple steps as follows:



Initial Value Problems: Euler's method

Starting at time t_0 with the given initial value y_0 , we would like to track the solution trajectory governed by the ODE

$$y' = f(t, y)$$

The fundamental theorem of calculus tells us that

$$y(t_{k+1}) = y(t_k) + \int_{t_k}^{t_{k+1}} y'(s) ds = y(t_k) + \int_{t_k}^{t_{k+1}} f(s, y(s)) ds$$


The simplest approximation for the integral is by using the left end-point rule, that is,

$$\int_a^b g(s) ds \approx (b - a)g(a)$$

yielding the method

$$y(T) = y(t_0) + (T - t_0)f(t_0, y_0)$$

To have more control over the error, we can also do this in multiple steps as follows:



Initial Value Problems: Euler's method

Starting at time t_0 with the given initial value y_0 , we would like to track the solution trajectory governed by the ODE

$$y' = f(t, y)$$

The fundamental theorem of calculus tells us that

$$y(t_{k+1}) = y(t_k) + \int_{t_k}^{t_{k+1}} y'(s) ds = y(t_k) + \int_{t_k}^{t_{k+1}} f(s, y(s)) ds$$


The simplest approximation for the integral is by using the left end-point rule, that is,

$$\int_a^b g(s) ds \approx (b - a)g(a)$$

yielding the method

$$y(T) = y(t_0) + (T - t_0)f(t_0, y_0)$$

To have more control over the error, we can also do this in multiple steps as follows:



Initial Value Problems: Euler's method

Starting at time t_0 with the given initial value y_0 , we would like to track the solution trajectory governed by the ODE

$$y' = f(t, y)$$

The fundamental theorem of calculus tells us that

$$y(t_{k+1}) = y(t_k) + \int_{t_k}^{t_{k+1}} y'(s) ds = y(t_k) + \int_{t_k}^{t_{k+1}} f(s, y(s)) ds$$


The simplest approximation for the integral is by using the left end-point rule, that is,

$$\int_a^b g(s) ds \approx (b - a)g(a)$$

yielding the method

$$y(T) = y(t_0) + (T - t_0)f(t_0, y_0)$$

To have more control over the error, we can also do this in multiple steps as follows:



Initial Value Problems: Euler's method

Starting at time t_0 with the given initial value y_0 , we would like to track the solution trajectory governed by the ODE

$$y' = f(t, y)$$

The fundamental theorem of calculus tells us that

$$y(t_{k+1}) = y(t_k) + \int_{t_k}^{t_{k+1}} y'(s) ds = y(t_k) + \int_{t_k}^{t_{k+1}} f(s, y(s)) ds$$


The simplest approximation for the integral is by using the left end-point rule, that is,

$$\int_a^b g(s) ds \approx (b - a)g(a)$$


yielding the method

$$y(T) = y(t_0) + (T - t_0)f(t_0, y_0)$$

To have more control over the error, we can also do this in multiple steps as follows:



$$y_{k+1} = y_k + (t_{k+1} - t_k)f(t_k, y_k)$$



Initial Value Problems: Euler's method

Starting at time t_0 with the given initial value y_0 , we would like to track the solution trajectory governed by the ODE

$$y' = f(t, y)$$

The fundamental theorem of calculus tells us that

$$y(t_{k+1}) = y(t_k) + \int_{t_k}^{t_{k+1}} y'(s) ds = y(t_k) + \int_{t_k}^{t_{k+1}} f(s, y(s)) ds$$

The simplest approximation for the integral is by using the left end-point rule, that is,

$$\int_a^b g(s) ds \approx (b - a)g(a)$$

yielding the method

$$y(T) = y(t_0) + (T - t_0)f(t_0, y_0)$$

To have more control over the error, we can also do this in multiple steps as follows:

