

Initial Value Problems: Linear Multistep Methods

Theorem

The linear multistep method is convergent if and only if it is consistent and stable.

Remark

This theorem states that it can be determined whether or not a linear multistep method is convergent simply by checking some algebraic conditions concerning its characteristic polynomials.

Proof. (sketch)

Convergence \Rightarrow Consistency

Apply the method to $y' = 0, y(0) = 1$ and $y' = 1, y(0) = 0$ for verifying satisfiability of the consistency conditions.

Convergence \Rightarrow Stability

Consistency and Stability \Rightarrow Convergence

Initial Value Problems: Linear Multistep Methods

Theorem

The linear multistep method is convergent if and only if it is consistent and stable.

Remark

This theorem states that it can be determined whether or not a linear multistep method is convergent simply by checking some algebraic conditions concerning its characteristic polynomials.

Proof. (sketch)

Convergence \Rightarrow Consistency

Apply the method to $y' = 0, y(0) = 1$ and $y' = 1, y(0) = 0$ for verifying satisfiability of the consistency conditions.

Convergence \Rightarrow Stability

Apply the method to $y' = 0, y(0) = 0$ for verifying satisfiability of the root condition.

Consistency and Stability \Rightarrow Convergence

Initial Value Problems: Linear Multistep Methods

Theorem

The linear multistep method is convergent if and only if it is consistent and stable.

Remark

This theorem states that it can be determined whether or not a linear multistep method is convergent simply by checking some algebraic conditions concerning its characteristic polynomials.

Proof. (sketch)

Convergence \Rightarrow Consistency

Apply the method to $y' = 0, y(0) = 1$ and $y' = 1, y(0) = 0$ for verifying satisfiability of the consistency conditions.

Convergence \Rightarrow Stability

Apply the method to $y' = 0, y(0) = 0$ for verifying satisfiability of the root condition.

Consistency and Stability \Rightarrow Convergence

Recall that

$$\begin{bmatrix} y_{n-k+1} \\ y_{n-k+2} \\ \vdots \\ y_n \\ y_{n+1} \end{bmatrix} = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ -a_k & -a_{k-1} & -a_{k-2} & \cdots & -a_0 \end{bmatrix} \begin{bmatrix} y_{n-k} \\ y_{n-k+1} \\ \vdots \\ y_{n-1} \\ y_n \end{bmatrix} + \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ h(b_{-1}f_{n+1} + b_0f_n + \cdots + b_nf_0) \end{bmatrix}$$

Initial Value Problems: Linear Multistep Methods

We define

$$A = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ -a_k & -a_{k-1} & -a_{k-2} & \cdots & -a_0 \end{bmatrix}, \quad E_n = \begin{bmatrix} y_{n-k} - y(t_{n-k}) \\ y_{n-k+1} - y(t_{n-k+1}) \\ \vdots \\ y_{n-1} - y(t_{n-1}) \\ y_n - y(t_n) \end{bmatrix},$$

where $y(t)$ is the exact solution that satisfies a similar difference equation

$$\begin{bmatrix} y(t_{n-k+1}) \\ y(t_{n-k+2}) \\ \vdots \\ y(t_n) \\ y(t_{n+1}) \end{bmatrix} = A \begin{bmatrix} y(t_{n-k}) \\ y(t_{n-k+1}) \\ \vdots \\ y(t_{n-1}) \\ y(t_n) \end{bmatrix} + \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ h(b_{-1}f(t_{n+1}, y(t_{n+1})) + b_0f(t_n, y(t_n)) + \cdots + b_nf(t_0, y(t_0))) - \ell_{n+1}(y, h) \end{bmatrix}.$$

Initial Value Problems: Linear Multistep Methods

We define

$$A = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ -a_k & -a_{k-1} & -a_{k-2} & \cdots & -a_0 \end{bmatrix}, \quad E_n = \begin{bmatrix} y_{n-k} - y(t_{n-k}) \\ y_{n-k+1} - y(t_{n-k+1}) \\ \vdots \\ y_{n-1} - y(t_{n-1}) \\ y_n - y(t_n) \end{bmatrix},$$

where $y(t)$ is the exact solution that satisfies a similar difference equation

$$\begin{bmatrix} y(t_{n-k+1}) \\ y(t_{n-k+2}) \\ \vdots \\ y(t_n) \\ y(t_{n+1}) \end{bmatrix} = A \begin{bmatrix} y(t_{n-k}) \\ y(t_{n-k+1}) \\ \vdots \\ y(t_{n-1}) \\ y(t_n) \end{bmatrix} + \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ h(b_{-1}f(t_{n+1}, y(t_{n+1})) + b_0f(t_n, y(t_n)) + \cdots + b_nf(t_0, y(t_0))) - \ell_{n+1}(y, h) \end{bmatrix}.$$

Then, we have $E_{n+1} = AE_n + Q_n$ where

$$Q_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ h(b_{-1}(f_{n+1} - f(t_{n+1}, y(t_{n+1}))) + b_0(f_n - f(t_n, y(t_n)))) + \cdots + b_n(f_0 - f(t_0, y(t_0)))) + \ell_{n+1}(y, h) \end{bmatrix}.$$

Initial Value Problems: Linear Multistep Methods

We define

$$A = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ -a_k & -a_{k-1} & -a_{k-2} & \cdots & -a_0 \end{bmatrix}, \quad E_n = \begin{bmatrix} y_{n-k} - y(t_{n-k}) \\ y_{n-k+1} - y(t_{n-k+1}) \\ \vdots \\ y_{n-1} - y(t_{n-1}) \\ y_n - y(t_n) \end{bmatrix},$$

where $y(t)$ is the exact solution that satisfies a similar difference equation

Then, we have $E_{n+1} = AE_n + Q_n$ where

$$Q_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ h(b_{-1}(f_{n+1} - f(t_{n+1}, y(t_{n+1}))) + b_0(f_n - f(t_n, y(t_n))) + \cdots + b_n(f_0 - f(t_0, y(t_0)))) + \ell_{n+1}(y, h) \end{bmatrix}.$$

So, there is a constant C such that, for $h \leq (2C\|b\|_1 L)^{-1}$, we have

$$\|E_{k+n}\| \leq C\|E_k\| + C \sum_{j=0}^{n-1} \|Q_{k+j}\| \leq 2C\|E_k\| + 4hC\|b\|_1 L \sum_{j=0}^{n-1} \|E_{k+j}\| + 2nC \max_{0 \leq j < n} |\ell_{k+j+1}(y, h)|.$$

Initial Value Problems: Linear Multistep Methods

We define

$$A = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ -a_k & -a_{k-1} & -a_{k-2} & \cdots & -a_0 \end{bmatrix}, \quad E_n = \begin{bmatrix} y_{n-k} - y(t_{n-k}) \\ y_{n-k+1} - y(t_{n-k+1}) \\ \vdots \\ y_{n-1} - y(t_{n-1}) \\ y_n - y(t_n) \end{bmatrix},$$

where $y(t)$ is the exact solution that satisfies a similar difference equation

Then, we have $E_{n+1} = AE_n + Q_n$ where

$$Q_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ h(b_{-1}(f_{n+1} - f(t_{n+1}, y(t_{n+1}))) + b_0(f_n - f(t_n, y(t_n)))) + \cdots + b_n(f_0 - f(t_0, y(t_0))) + \ell_{n+1}(y, h) \end{bmatrix}.$$

So, there is a constant C such that, for $h \leq (2C\|b\|_1 L)^{-1}$, we have

$$\|E_{k+n}\| \leq C\|E_k\| + C \sum_{j=0}^{n-1} \|Q_{k+j}\| \leq 2C\|E_k\| + 4hC\|b\|_1 L \sum_{j=0}^{n-1} \|E_{k+j}\| + 2nC \max_{0 \leq j < n} |\ell_{k+j+1}(y, h)|.$$

Therefore,

$$\|E_{k+n}\| \leq 2C \left(\|E_k\| + (T - t_0) \max_{0 \leq j < N} \left| \frac{\ell_j(y, h)}{h} \right| \right) e^{4(T-t_0)C\|b\|_1 L} \dots$$

Initial Value Problems: Linear Multistep Methods



Remark

The highest order attainable by a k -step method is $2k$.

Initial Value Problems: Linear Multistep Methods



Remark

The highest order attainable by a k -step method is $2k$. However, such a method is not stable for any $k > 1$.

Initial Value Problems: Linear Multistep Methods



Remark

The highest order attainable by a k -step method is $2k$. However, such a method is not stable for any $k > 1$.

Theorem

The highest order of a stable k -step method is $k + 1$ if k is odd and $k + 2$ if k is even.

Initial Value Problems: Linear Multistep Methods

Remark

The highest order attainable by a k -step method is $2k$. However, such a method is not stable for any $k > 1$.

Theorem

The highest order of a stable k -step method is $k + 1$ if k is odd and $k + 2$ if k is even.

Remark

The Adams method are linear multistep methods with best possible stability properties, namely, the first characteristic polynomial $\rho(t) = t^{k+1} - t^k$ has all its roots at the origin except for the mandatory root at 1.