

Numerical Analysis & Scientific Computing II

Module 2

Initial Value Problems

2.1 Well-posedness

2.2 Stability

2.3 Euler's method

- Derivations



Akash Anand
MATH, IIT KANPUR

Initial Value Problems: Euler's method



Derivation using Taylor series



Derivation using Taylor series

Consider the Taylor series

$$y(t_{k+1}) = y(t_k) + (t_{k+1} - t_k)y'(t_k) + \frac{(t_{k+1} - t_k)^2}{2}y''(t_k) + \dots$$



Derivation using Taylor series

Consider the Taylor series

$$y(t_{k+1}) = y(t_k) + (t_{k+1} - t_k)y'(t_k) + \frac{(t_{k+1} - t_k)^2}{2}y''(t_k) + \dots$$

The Euler's method results from taking $y_k = y(t_k)$, $y'(t_k) = f(t_k, y_k)$ and dropping terms of second and higher orders!

Initial Value Problems: Euler's method

Derivation using Taylor series

Consider the Taylor series

$$y(t_{k+1}) = y(t_k) + (t_{k+1} - t_k)y'(t_k) + \frac{(t_{k+1} - t_k)^2}{2}y''(t_k) + \dots$$

The Euler's method results from taking $y_k = y(t_k)$, $y'(t_k) = f(t_k, y_k)$ and dropping terms of second and higher orders!

Derivation using finite difference approximation

Initial Value Problems: Euler's method

Derivation using Taylor series

Consider the Taylor series

$$y(t_{k+1}) = y(t_k) + (t_{k+1} - t_k)y'(t_k) + \frac{(t_{k+1} - t_k)^2}{2}y''(t_k) + \dots$$

The Euler's method results from taking $y_k = y(t_k)$, $y'(t_k) = f(t_k, y_k)$ and dropping terms of second and higher orders!

Derivation using finite difference approximation

Replacing the $y'(t)$ in the ODE $y' = f(t, y)$ by a first order forward difference approximation, we obtain an algebraic equation

$$\frac{y_{k+1} - y_k}{t_{k+1} - t_k} = f(t_k, y_k)$$

that yields the Euler's method.

Initial Value Problems: Euler's method



Derivation using polynomial interpolation

Initial Value Problems: Euler's method



Derivation using polynomial interpolation

One point Hermite polynomial $p(t)$ that matches the function and derivative data at $t = t_k$, that is,

$$p(t_k) = y(t_k) (= y_k), \quad p'(t) = y'(t_k) (= f(t_k, y_k)),$$

Initial Value Problems: Euler's method

Derivation using polynomial interpolation

One point Hermite polynomial $p(t)$ that matches the function and derivative data at $t = t_k$, that is,

$$p(t_k) = y(t_k) (= y_k), \quad p'(t) = y'(t_k) (= f(t_k, y_k)),$$

is given by

$$p(t) = y_k + (t - t_k)f(t_k, y_k)$$

Initial Value Problems: Euler's method



Derivation using polynomial interpolation

One point Hermite polynomial $p(t)$ that matches the function and derivative data at $t = t_k$, that is,

$$p(t_k) = y(t_k) (= y_k), \quad p'(t) = y'(t_k) (= f(t_k, y_k)),$$

is given by

$$p(t) = y_k + (t - t_k)f(t_k, y_k)$$

The Euler's method results from evaluating this polynomial at $t = t_{k+1}$.

Initial Value Problems: Euler's method

Derivation using polynomial interpolation

One point Hermite polynomial $p(t)$ that matches the function and derivative data at $t = t_k$, that is,

$$p(t_k) = y(t_k) (= y_k), \quad p'(t) = y'(t_k) (= f(t_k, y_k)),$$

is given by

$$p(t) = y_k + (t - t_k)f(t_k, y_k)$$

The Euler's method results from evaluating this polynomial at $t = t_{k+1}$.

Derivation using the method of undetermined coefficients

Initial Value Problems: Euler's method

Derivation using polynomial interpolation

One point Hermite polynomial $p(t)$ that matches the function and derivative data at $t = t_k$, that is,

$$p(t_k) = y(t_k) (= y_k), \quad p'(t) = y'(t_k) (= f(t_k, y_k)),$$

is given by

$$p(t) = y_k + (t - t_k)f(t_k, y_k)$$

The Euler's method results from evaluating this polynomial at $t = t_{k+1}$.

Derivation using the method of undetermined coefficients

At $t = t_k$, we know the values y_k and y'_k , and based on these values we want to predict the value y_{k+1} .

Initial Value Problems: Euler's method

Derivation using polynomial interpolation

One point Hermite polynomial $p(t)$ that matches the function and derivative data at $t = t_k$, that is,

$$p(t_k) = y(t_k) (= y_k), \quad p'(t) = y'(t_k) (= f(t_k, y_k)),$$

is given by

$$p(t) = y_k + (t - t_k)f(t_k, y_k)$$

The Euler's method results from evaluating this polynomial at $t = t_{k+1}$.

Derivation using the method of undetermined coefficients

At $t = t_k$, we know the values y_k and y'_k , and based on these values we want to predict the value y_{k+1} . A predictor based on their linear combination has the form

$$y_{k+1} = \alpha y_k + \beta y'_k$$

where α and β are coefficients to be determined.

Initial Value Problems: Euler's method

Derivation using polynomial interpolation

One point Hermite polynomial $p(t)$ that matches the function and derivative data at $t = t_k$, that is,

$$p(t_k) = y(t_k) (= y_k), \quad p'(t) = y'(t_k) (= f(t_k, y_k)),$$

is given by

$$p(t) = y_k + (t - t_k)f(t_k, y_k)$$

The Euler's method results from evaluating this polynomial at $t = t_{k+1}$.

Derivation using the method of undetermined coefficients

At $t = t_k$, we know the values y_k and y'_k , and based on these values we want to predict the value y_{k+1} . A predictor based on their linear combination has the form

$$y_{k+1} = \alpha y_k + \beta y'_k$$

where α and β are coefficients to be determined. To find the coefficients, we enforce that the predictor should be exact for first two monomials, that is, for $y(t) = e$ and $y(t) = te$, where e is the vector whose components are all ones.

Initial Value Problems: Euler's method

Derivation using polynomial interpolation

One point Hermite polynomial $p(t)$ that matches the function and derivative data at $t = t_k$, that is,

$$p(t_k) = y(t_k) (= y_k), \quad p'(t) = y'(t_k) (= f(t_k, y_k)),$$

is given by

$$p(t) = y_k + (t - t_k)f(t_k, y_k)$$

The Euler's method results from evaluating this polynomial at $t = t_{k+1}$.

Derivation using the method of undetermined coefficients

At $t = t_k$, we know the values y_k and y'_k , and based on these values we want to predict the value y_{k+1} . A predictor based on their linear combination has the form

$$y_{k+1} = \alpha y_k + \beta y'_k$$

where α and β are coefficients to be determined. To find the coefficients, we enforce that the predictor should be exact for first two monomials, that is, for $y(t) = e$ and $y(t) = te$, where e is the vector whose components are all ones. Therefore, we have

$$e = \alpha e + \beta 0, \quad t_{k+1}e = \alpha t_k e + \beta e$$

Initial Value Problems: Euler's method

Derivation using polynomial interpolation

One point Hermite polynomial $p(t)$ that matches the function and derivative data at $t = t_k$, that is,

$$p(t_k) = y(t_k) (= y_k), \quad p'(t) = y'(t_k) (= f(t_k, y_k)),$$

is given by

$$p(t) = y_k + (t - t_k)f(t_k, y_k)$$

The Euler's method results from evaluating this polynomial at $t = t_{k+1}$.

Derivation using the method of undetermined coefficients

At $t = t_k$, we know the values y_k and y'_k , and based on these values we want to predict the value y_{k+1} . A predictor based on their linear combination has the form

$$y_{k+1} = \alpha y_k + \beta y'_k$$

where α and β are coefficients to be determined. To find the coefficients, we enforce that the predictor should be exact for first two monomials, that is, for $y(t) = e$ and $y(t) = te$, where e is the vector whose components are all ones. Therefore, we have

$$e = \alpha e + \beta 0, \quad t_{k+1}e = \alpha t_k e + \beta e$$

implying $\alpha = 1$ and $\beta = t_{k+1} - t_k$ resulting in the Euler's method.

Numerical Analysis & Scientific Computing II

Module 2

Initial Value Problems

2.1 Well-posedness

2.2 Stability

2.3 Euler's method

**- Errors and error
propagation**




Akash Anand
MATH, IIT KANPUR



Example

Let us solve $y' = y$, $y(0) = 1$ using the Euler's method taking the uniform step size $h = h_k = t_{k+1} - t_k = 0.5$.


$$y_1 = 1 + h, \quad y_2 = y_1 + hy_1 = (1 + h)^2, \quad y_3 = y_2 + hy_2 = (1 + h)^3, \dots, \quad y_k = (1 + h)^k, \dots$$



Example

Let us solve $y' = y$, $y(0) = 1$ using the Euler's method taking the uniform step size $h = h_k = t_{k+1} - t_k = 0.5$.

$$y_1 = 1 + h, \quad y_2 = y_1 + hy_1 = (1 + h)^2, \quad y_3 = y_2 + hy_2 = (1 + h)^3, \dots, \quad y_k = (1 + h)^k, \dots$$




Initial Value Problems: Euler's method

Example

Let us solve $y' = y$, $y(0) = 1$ using the Euler's method taking the uniform step size $h = h_k = t_{k+1} - t_k = 0.5$.


$$y_1 = 1 + h, \quad y_2 = y_1 + hy_1 = (1 + h)^2, \quad y_3 = y_2 + hy_2 = (1 + h)^3, \dots, \quad y_k = (1 + h)^k, \dots$$



Example

Let us solve $y' = y$, $y(0) = 1$ using the Euler's method taking the uniform step size $h = h_k = t_{k+1} - t_k = 0.5$.

$$y_1 = 1 + h, \quad y_2 = y_1 + hy_1 = (1 + h)^2, \quad y_3 = y_2 + hy_2 = (1 + h)^3, \dots, \quad y_k = (1 + h)^k, \dots$$




Initial Value Problems: Euler's method



Example

Let us solve $y' = y$, $y(0) = 1$ using the Euler's method taking the uniform step size $h = h_k = t_{k+1} - t_k = 0.5$.

$$y_1 = 1 + h, \quad y_2 = y_1 + hy_1 = (1 + h)^2, \quad y_3 = y_2 + hy_2 = (1 + h)^3, \dots, \quad y_k = (1 + h)^k, \dots$$




Initial Value Problems: Euler's method



Example

Let us solve $y' = y$, $y(0) = 1$ using the Euler's method taking the uniform step size $h = h_k = t_{k+1} - t_k = 0.5$.

$$y_1 = 1 + h, \quad y_2 = y_1 + hy_1 = (1 + h)^2, \quad y_3 = y_2 + hy_2 = (1 + h)^3, \dots, \quad y_k = (1 + h)^k, \dots$$




Initial Value Problems: Euler's method

Example

Let us solve $y' = y$, $y(0) = 1$ using the Euler's method taking the uniform step size $h = h_k = t_{k+1} - t_k = 0.5$.

$$y_1 = 1 + h, \quad y_2 = y_1 + hy_1 = (1 + h)^2, \quad y_3 = y_2 + hy_2 = (1 + h)^3, \dots, \quad y_k = (1 + h)^k, \dots$$

An error is introduced at each step of the method – *one step error* or *single step error*.



Initial Value Problems: Euler's method


Example

Let us solve $y' = y$, $y(0) = 1$ using the Euler's method taking the uniform step size $h = h_k = t_{k+1} - t_k = 0.5$.

$$y_1 = 1 + h, \quad y_2 = y_1 + hy_1 = (1 + h)^2, \quad y_3 = y_2 + hy_2 = (1 + h)^3, \dots, \quad y_k = (1 + h)^k, \dots$$

An error is introduced at each step of the method – *one step error* or *single step error*.

These local errors accumulate over time to produce a *global error*.



Initial Value Problems: Euler's method

Example


Let us solve $y' = y$, $y(0) = 1$ using the Euler's method taking the uniform step size $h = h_k = t_{k+1} - t_k = 0.5$.

$$y_1 = 1 + h, \quad y_2 = y_1 + hy_1 = (1 + h)^2, \quad y_3 = y_2 + hy_2 = (1 + h)^3, \dots, \quad y_k = (1 + h)^k, \dots$$

An error is introduced at each step of the method – **one step error** or **single step error**.

These local errors accumulate over time to produce a **global error**.

Note that how local errors accumulate is not always clear and it depends on the underlying solution.



Initial Value Problems: Euler's method

Example


Let us solve $y' = y$, $y(0) = 1$ using the Euler's method taking the uniform step size $h = h_k = t_{k+1} - t_k = 0.5$.

$$y_1 = 1 + h, \quad y_2 = y_1 + hy_1 = (1 + h)^2, \quad y_3 = y_2 + hy_2 = (1 + h)^3, \dots, \quad y_k = (1 + h)^k, \dots$$

An error is introduced at each step of the method – **one step error** or **single step error**.

These local errors accumulate over time to produce a **global error**.

Note that how local errors accumulate is not always clear and it depends on the underlying solution.



With each step, we hop from one solution of the ODE to another.

Initial Value Problems: Euler's method

Example


Let us solve $y' = y$, $y(0) = 1$ using the Euler's method taking the uniform step size $h = h_k = t_{k+1} - t_k = 0.5$.

$$y_1 = 1 + h, \quad y_2 = y_1 + hy_1 = (1 + h)^2, \quad y_3 = y_2 + hy_2 = (1 + h)^3, \dots, \quad y_k = (1 + h)^k, \dots$$

An error is introduced at each step of the method – **one step error** or **single step error**.

These local errors accumulate over time to produce a **global error**.

Note that how local errors accumulate is not always clear and it depends on the underlying solution.



With each step, we hop from one solution of the ODE to another.

Local errors get amplified if the solutions to the ODE are unstable.

Initial Value Problems: Euler's method



Example

Let us solve $y' = y$, $y(0) = 1$ using the Euler's method taking the uniform step size $h = h_k = t_{k+1} - t_k = 0.5$.

$$y_1 = 1 + h, \quad y_2 = y_1 + hy_1 = (1 + h)^2, \quad y_3 = y_2 + hy_2 = (1 + h)^3, \dots, \quad y_k = (1 + h)^k, \dots$$



With each step, we hop from one solution of the ODE to another.

Local errors get amplified if the solutions to the ODE are unstable.

Initial Value Problems: Euler's method



Example

Let us solve $y' = -y$, $y(0) = 1$ using the Euler's method taking the uniform step size $h = 0.5$.



Example

Let us solve $y' = -y$, $y(0) = 1$ using the Euler's method taking the uniform step size $h = 0.5$.

$$y_1 = 1 - h, \quad y_2 = y_1 - hy_1 = (1 - h)^2, \quad y_3 = y_2 - hy_2 = (1 - h)^3, \dots, \quad y_k = (1 - h)^k, \dots$$


Initial Value Problems: Euler's method



Example

Let us solve $y' = -y$, $y(0) = 1$ using the Euler's method taking the uniform step size $h = 0.5$.

$$y_1 = 1 - h, \quad y_2 = y_1 - hy_1 = (1 - h)^2, \quad y_3 = y_2 - hy_2 = (1 - h)^3, \dots, \quad y_k = (1 - h)^k, \dots$$




Initial Value Problems: Euler's method



Example

Let us solve $y' = -y$, $y(0) = 1$ using the Euler's method taking the uniform step size $h = 0.5$.

$$y_1 = 1 - h, \quad y_2 = y_1 - hy_1 = (1 - h)^2, \quad y_3 = y_2 - hy_2 = (1 - h)^3, \dots, \quad y_k = (1 - h)^k, \dots$$



For an equation with stable solutions, the errors in the numerical solution do not grow, and for equations with asymptotically stable solutions, the errors diminish with time.

Initial Value Problems: Accuracy and Stability



Sources of error

Rounding error –

... due to truncation of data (e.g., real numbers requiring infinite space is represented using a finite amount of space); finite precision of the floating point arithmetic.

Truncation error –

... due to truncation of infinite mathematical processes to finite processes or algorithms (e.g., derivative replaced by a finite difference approximation, or integration replaced by a quadrature).

Initial Value Problems: Accuracy and Stability



Sources of error

Rounding error –

... due to truncation of data (e.g., real numbers requiring infinite space is represented using a finite amount of space); finite precision of the floating point arithmetic.

Truncation error –

... due to truncation of infinite mathematical processes to finite processes or algorithms (e.g., derivative replaced by a finite difference approximation, or integration replaced by a quadrature).

Remarks

Errors propagate from one step to the next. We have seen how single step error in the Euler's method can get amplified over multiple steps.

Initial Value Problems: Accuracy and Stability

Sources of error

Rounding error –

... due to truncation of data (e.g., real numbers requiring infinite space is represented using a finite amount of space); finite precision of the floating point arithmetic.

Truncation error –

... due to truncation of infinite mathematical processes to finite processes or algorithms (e.g., derivative replaced by a finite difference approximation, or integration replaced by a quadrature).

Remarks

Errors propagate from one step to the next. We have seen how single step error in the Euler's method can get amplified over multiple steps.

In most practical situations, truncation error dominates other sources and, therefore, in analysis of numerical methods, we will focus exclusively on truncation error.