

Numerical Analysis & Scientific Computing II

Lesson 4

Numerical Solution of PDE

4.3 Hyperbolic PDE

- Finite Difference Methods



Numerical Methods for PDE: Hyperbolic PDE



Now let's look at the difference methods for advection equation. For simplicity, we will investigate the model problem of the advection equation.

Numerical Methods for PDE: Hyperbolic PDE



Now let's look at the difference methods for advection equation. For simplicity, we will investigate the model problem of the advection equation.

To begin with, let's study the forward-forward, forward-centered and forward-backward difference methods.



Now let's look at the difference methods for advection equation. For simplicity, we will investigate the model problem of the advection equation.

To begin with, let's study the forward-forward, forward-centered and forward-backward difference methods. These three methods are -

forward-forward:

$$\frac{u_n^{j+1} - u_n^j}{k} + c \frac{u_{n+1}^j - u_n^j}{h} = 0$$

Numerical Methods for PDE: Hyperbolic PDE

Now let's look at the difference methods for advection equation. For simplicity, we will investigate the model problem of the advection equation.

To begin with, let's study the forward-forward, forward-centered and forward-backward difference methods. These three methods are -

forward-forward:

$$\frac{u_n^{j+1} - u_n^j}{k} + c \frac{u_{n+1}^j - u_n^j}{h} = 0$$

forward-backward:

$$\frac{u_n^{j+1} - u_n^j}{k} + c \frac{u_n^j - u_{n-1}^j}{h} = 0$$

Numerical Methods for PDE: Hyperbolic PDE

Now let's look at the difference methods for advection equation. For simplicity, we will investigate the model problem of the advection equation.

To begin with, let's study the forward-forward, forward-centered and forward-backward difference methods. These three methods are -

forward-forward:

$$\frac{u_n^{j+1} - u_n^j}{k} + c \frac{u_{n+1}^j - u_n^j}{h} = 0$$

forward-backward:

$$\frac{u_n^{j+1} - u_n^j}{k} + c \frac{u_n^j - u_{n-1}^j}{h} = 0$$

forward-centered:

$$\frac{u_n^{j+1} - u_n^j}{k} + c \frac{u_{n+1}^j - u_{n-1}^j}{2h} = 0$$

Numerical Methods for PDE: Hyperbolic PDE

Now let's look at the difference methods for advection equation. For simplicity, we will investigate the model problem of the advection equation.

To begin with, let's study the forward-forward, forward-centered and forward-backward difference methods. These three methods are -

forward-forward:

$$\frac{u_n^{j+1} - u_n^j}{k} + c \frac{u_{n+1}^j - u_n^j}{h} = 0$$

forward-backward:

$$\frac{u_n^{j+1} - u_n^j}{k} + c \frac{u_n^j - u_{n-1}^j}{h} = 0$$

forward-centered:

$$\frac{u_n^{j+1} - u_n^j}{k} + c \frac{u_{n+1}^j - u_{n-1}^j}{2h} = 0$$

If we set $\lambda = ck/h$, these can be written as

forward-forward:

forward-backward:

forward-centered:

Numerical Methods for PDE: Hyperbolic PDE

Now let's look at the difference methods for advection equation. For simplicity, we will investigate the model problem of the advection equation.

To begin with, let's study the forward-forward, forward-centered and forward-backward difference methods. These three methods are -

forward-forward:

$$\frac{u_n^{j+1} - u_n^j}{k} + c \frac{u_{n+1}^j - u_n^j}{h} = 0$$

forward-backward:

$$\frac{u_n^{j+1} - u_n^j}{k} + c \frac{u_n^j - u_{n-1}^j}{h} = 0$$

forward-centered:

$$\frac{u_n^{j+1} - u_n^j}{k} + c \frac{u_{n+1}^j - u_{n-1}^j}{2h} = 0$$

If we set $\lambda = ck/h$, these can be written as

forward-forward: $u_n^{j+1} = -\lambda u_{n+1}^j + (1 + \lambda)u_n^j$

forward-backward: $u_n^{j+1} = (1 - \lambda)u_n^j + \lambda u_{n-1}^j$

forward-centered: $u_n^{j+1} = -\frac{\lambda}{2}u_{n+1}^j + u_n^j + \frac{\lambda}{2}u_{n-1}^j$

Numerical Methods for PDE: Hyperbolic PDE

If we set $\lambda = ck/h$, these can be written as

forward-forward: $u_n^{j+1} = -\lambda u_{n+1}^j + (1 + \lambda)u_n^j$

forward-backward: $u_n^{j+1} = (1 - \lambda)u_n^j + \lambda u_{n-1}^j$

forward-centered: $u_n^{j+1} = -\frac{\lambda}{2}u_{n+1}^j + u_n^j + \frac{\lambda}{2}u_{n-1}^j$

How do these method perform? Are they equally good?

Numerical Methods for PDE: Hyperbolic PDE

If we set $\lambda = ck/h$, these can be written as

forward-forward: $u_n^{j+1} = -\lambda u_{n+1}^j + (1 + \lambda)u_n^j$

forward-backward: $u_n^{j+1} = (1 - \lambda)u_n^j + \lambda u_{n-1}^j$

forward-centered: $u_n^{j+1} = -\frac{\lambda}{2}u_{n+1}^j + u_n^j + \frac{\lambda}{2}u_{n-1}^j$

How do these method perform? Are they equally good?

The local truncation error is $O(k + h)$ for the first two methods and $O(k + h^2)$ for the third method.

Numerical Methods for PDE: Hyperbolic PDE

If we set $\lambda = ck/h$, these can be written as

forward-forward: $u_n^{j+1} = -\lambda u_{n+1}^j + (1 + \lambda)u_n^j$

forward-backward: $u_n^{j+1} = (1 - \lambda)u_n^j + \lambda u_{n-1}^j$

forward-centered: $u_n^{j+1} = -\frac{\lambda}{2}u_{n+1}^j + u_n^j + \frac{\lambda}{2}u_{n-1}^j$

How do these method perform? Are they equally good?

The local truncation error is $O(k + h)$ for the first two methods and $O(k + h^2)$ for the third method.

It is easy to establish non-convergence of forward-forward difference method even without looking at the stability.

Numerical Methods for PDE: Hyperbolic PDE

If we set $\lambda = ck/h$, these can be written as

forward-forward: $u_n^{j+1} = -\lambda u_{n+1}^j + (1 + \lambda)u_n^j$

forward-backward: $u_n^{j+1} = (1 - \lambda)u_n^j + \lambda u_{n-1}^j$

forward-centered: $u_n^{j+1} = -\frac{\lambda}{2}u_{n+1}^j + u_n^j + \frac{\lambda}{2}u_{n-1}^j$

How do these method perform? Are they equally good?

The local truncation error is $O(k + h)$ for the first two methods and $O(k + h^2)$ for the third method.

It is easy to establish non-convergence of forward-forward difference method even without looking at the stability. It is easy to see that the value at a mesh-point $x = nh$ at a time $t = jk$ depends only on the initial data at the mesh points in the interval $[x, x + ct/\lambda]$.

Numerical Methods for PDE: Hyperbolic PDE

If we set $\lambda = ck/h$, these can be written as

forward-forward: $u_n^{j+1} = -\lambda u_{n+1}^j + (1 + \lambda)u_n^j$


forward-backward: $u_n^{j+1} = (1 - \lambda)u_n^j + \lambda u_{n-1}^j$

forward-centered: $u_n^{j+1} = -\frac{\lambda}{2}u_{n+1}^j + u_n^j + \frac{\lambda}{2}u_{n-1}^j$

How do these methods perform? Are they equally good?

The local truncation error is $O(k + h)$ for the first two methods and $O(k + h^2)$ for the third method.

It is easy to establish non-convergence of forward-forward difference method even without looking at the stability. It is easy to see that the value at a mesh-point $x = nh$ at a time $t = jk$ depends only on the initial data at the mesh points in the interval $[x, x + ct/\lambda]$.



Numerical Methods for PDE: Hyperbolic PDE

If we set $\lambda = ck/h$, these can be written as

forward-forward: $u_n^{j+1} = -\lambda u_{n+1}^j + (1 + \lambda)u_n^j$


forward-backward: $u_n^{j+1} = (1 - \lambda)u_n^j + \lambda u_{n-1}^j$

forward-centered: $u_n^{j+1} = -\frac{\lambda}{2}u_{n+1}^j + u_n^j + \frac{\lambda}{2}u_{n-1}^j$

How do these methods perform? Are they equally good?

The local truncation error is $O(k + h)$ for the first two methods and $O(k + h^2)$ for the third method.

It is easy to establish non-convergence of forward-forward difference method even without looking at the stability. It is easy to see that the value at a mesh-point $x = nh$ at a time $t = jk$ depends only on the initial data at the mesh points in the interval $[x, x + ct/\lambda]$. This interval does not contain the point $x - ct$ for any positive value of λ .



Numerical Methods for PDE: Hyperbolic PDE

If we set $\lambda = ck/h$, these can be written as

forward-forward: $u_n^{j+1} = -\lambda u_{n+1}^j + (1 + \lambda)u_n^j$


forward-backward: $u_n^{j+1} = (1 - \lambda)u_n^j + \lambda u_{n-1}^j$

forward-centered: $u_n^{j+1} = -\frac{\lambda}{2}u_{n+1}^j + u_n^j + \frac{\lambda}{2}u_{n-1}^j$

How do these methods perform? Are they equally good?

The local truncation error is $O(k + h)$ for the first two methods and $O(k + h^2)$ for the third method.

It is easy to establish non-convergence of forward-forward difference method even without looking at the stability. It is easy to see that the value at a mesh-point $x = nh$ at a time $t = jk$ depends only on the initial data at the mesh points in the interval $[x, x + ct/\lambda]$. This interval does not contain the point $x - ct$ for any positive value of λ . Hence if we choose an initial data which is equal to 1 at $x - ct$ but identically zero on $[x, \infty)$, the true solution of the advection will satisfy $u(x, t) = 1$ but the numerical solution will be zero, and will remain zero as $h, k \rightarrow 0$.



Numerical Methods for PDE: Hyperbolic PDE

If we set $\lambda = ck/h$, these can be written as

forward-forward:
$$u_n^{j+1} = -\lambda u_{n+1}^j + (1 + \lambda)u_n^j$$

forward-backward:
$$u_n^{j+1} = (1 - \lambda)u_n^j + \lambda u_{n-1}^j$$


forward-centered:
$$u_n^{j+1} = -\frac{\lambda}{2}u_{n+1}^j + u_n^j + \frac{\lambda}{2}u_{n-1}^j$$

How do these method perform? Are they equally good?

The local truncation error is $O(k + h)$ for the first two methods and $O(k + h^2)$ for the third method.

It is easy to establish non-convergence of forward-forward difference method even without looking at the stability. It is easy to see that the value at a mesh-point $x = nh$ at a time $t = jk$ depends only on the initial data at the mesh points in the interval $[x, x + ct/\lambda]$. This interval does not contain the point $x - ct$ for any positive value of λ . Hence if we choose an initial data which is equal to 1 at $x - ct$ but identically zero on $[x, \infty)$, the true solution of the advection will satisfy $u(x, t) = 1$ but the numerical solution will be zero, and will remain zero as $h, k \rightarrow 0$.

In short, the **numerical domain of dependence** fails to include **true domain of dependence** which is necessary for a convergent method.



Numerical Methods for PDE: Hyperbolic PDE

If we set $\lambda = ck/h$, these can be written as

forward-forward: $u_n^{j+1} = -\lambda u_{n+1}^j + (1 + \lambda)u_n^j$

forward-backward: $u_n^{j+1} = (1 - \lambda)u_n^j + \lambda u_{n-1}^j$

forward-centered: $u_n^{j+1} = -\frac{\lambda}{2}u_{n+1}^j + u_n^j + \frac{\lambda}{2}u_{n-1}^j$


How do these method perform? Are they equally good?

The local truncation error is $O(k + h)$ for the first two methods and $O(k + h^2)$ for the third method.

It is easy to establish non-convergence of forward-forward difference method even without looking at the stability. It is easy to see that the value at a mesh-point $x = nh$ at a time $t = jk$ depends only on the initial data at the mesh points in the interval $[x, x + ct/\lambda]$. This interval does not contain the point $x - ct$ for any positive value of λ . Hence if we choose an initial data which is equal to 1 at $x - ct$ but identically zero on $[x, \infty)$, the true solution of the advection will satisfy $u(x, t) = 1$ but the numerical solution will be zero, and will remain zero as $h, k \rightarrow 0$.

In short, the **numerical domain of dependence** fails to include **true domain of dependence** which is necessary for a convergent method.

This necessary condition, which fails for forward-forward difference method, is called the **Courant-Friedrichs-Levy condition**, or **CFL condition**.



Numerical Methods for PDE: Hyperbolic PDE

If we set $\lambda = ck/h$, these can be written as

forward-forward: $u_n^{j+1} = -\lambda u_{n+1}^j + (1 + \lambda)u_n^j$


forward-backward: $u_n^{j+1} = (1 - \lambda)u_n^j + \lambda u_{n-1}^j$

forward-centered: $u_n^{j+1} = -\frac{\lambda}{2}u_{n+1}^j + u_n^j + \frac{\lambda}{2}u_{n-1}^j$

How do these method perform? Are they equally good?

The local truncation error is $O(k + h)$ for the first two methods and $O(k + h^2)$ for the third method.

If we apply the CFL condition to the forward-backward method, we see that the method can only be conditionally convergent.



Numerical Methods for PDE: Hyperbolic PDE

If we set $\lambda = ck/h$, these can be written as

forward-forward: $u_n^{j+1} = -\lambda u_{n+1}^j + (1 + \lambda)u_n^j$

forward-backward: $u_n^{j+1} = (1 - \lambda)u_n^j + \lambda u_{n-1}^j$

forward-centered: $u_n^{j+1} = -\frac{\lambda}{2}u_{n+1}^j + u_n^j + \frac{\lambda}{2}u_{n-1}^j$

How do these method perform? Are they equally good?


The local truncation error is $O(k + h)$ for the first two methods and $O(k + h^2)$ for the third method.

If we apply the CFL condition to the forward-backward method, we see that the method can only be conditionally convergent. Indeed, for convergence, we need


$$x - ct \in [x - ct/\lambda, x]$$

that is,

$$x \geq x - ct \geq x - ct/\lambda$$



<i>forward-forward:</i>	$u_n^{j+1} = -\lambda u_{n+1}^j + (1 + \lambda)u_n^j$
<i>forward-backward:</i>	$u_n^{j+1} = (1 - \lambda)u_n^j + \lambda u_{n-1}^j$
<i>forward-centered:</i>	$u_n^{j+1} = -\frac{\lambda}{2}u_{n+1}^j + u_n^j + \frac{\lambda}{2}u_{n-1}^j$



The local truncation error is $O(k + h)$ for the first two methods and $O(k + h^2)$ for the third method.

If we apply the CFL condition to the forward-backward method, we see that the method can only be conditionally convergent. Indeed, for convergence, we need

$$x - ct \in [x - ct/\lambda, x]$$

that is,

$$x \geq x - ct \geq x - ct/\lambda$$

and therefore, $0 < \lambda \leq 1$, is a necessary condition for convergence.

Numerical Methods for PDE: Hyperbolic PDE

If we set $\lambda = ck/h$, these can be written as

forward-forward: $u_n^{j+1} = -\lambda u_{n+1}^j + (1 + \lambda)u_n^j$

forward-backward: $u_n^{j+1} = (1 - \lambda)u_n^j + \lambda u_{n-1}^j$

forward-centered: $u_n^{j+1} = -\frac{\lambda}{2}u_{n+1}^j + u_n^j + \frac{\lambda}{2}u_{n-1}^j$

How do these method perform? Are they equally good?

The local truncation error is $O(k + h)$ for the first two methods and $O(k + h^2)$ for the third method.


If we apply the CFL condition to the forward-backward method, we see that the method can only be conditionally convergent. Indeed, for convergence, we need

$$x - ct \in [x - ct/\lambda, x]$$

that is,

$$x \geq x - ct \geq x - ct/\lambda$$

and therefore, $0 < \lambda \leq 1$, is a necessary condition for convergence. We will soon see that the method is stable and convergent if and only if this condition is met.



Numerical Methods for PDE: Hyperbolic PDE

If we set $\lambda = ck/h$, these can be written as

forward-forward: $u_n^{j+1} = -\lambda u_{n+1}^j + (1 + \lambda)u_n^j$

forward-backward: $u_n^{j+1} = (1 - \lambda)u_n^j + \lambda u_{n-1}^j$

forward-centered: $u_n^{j+1} = -\frac{\lambda}{2}u_{n+1}^j + u_n^j + \frac{\lambda}{2}u_{n-1}^j$

How do these method perform? Are they equally good?

The local truncation error is $O(k + h)$ for the first two methods and $O(k + h^2)$ for the third method.

If we apply the CFL condition to the forward-backward method, we see that the method can only be conditionally convergent. Indeed, for convergence, we need


$$x - ct \in [x - ct/\lambda, x]$$

that is,

$$x \geq x - ct \geq x - ct/\lambda$$

and therefore, $0 < \lambda \leq 1$, is a necessary condition for convergence. We will soon see that the method is stable and convergent if and only if this condition is met.

In general, however, CFL is not sufficient for convergence. It turns out that, for forward-central scheme, while the CFL condition is $|\lambda| \leq 1$, the method is **unconditionally unstable**.



Numerical Analysis & Scientific Computing II

Lesson 4

Numerical Solution of PDE

4.3 Hyperbolic PDE

- *Finite Difference Methods*
- ***Stability Analysis***



Numerical Methods for PDE: Hyperbolic PDE



Let's use the Fourier analysis (von Neumann stability analysis) to study the stability of these methods.



Numerical Methods for PDE: Hyperbolic PDE

Let's use the Fourier analysis (von Neumann stability analysis) to study the stability of these methods.

For simplicity, let's consider a 1-periodic problem rather than a boundary value problem:

$$\begin{aligned}\frac{\partial u}{\partial t} + c \frac{\partial u}{\partial x} &= 0, & x \in \mathbb{R}, t > 0, \\ u(x, 0) &= u_0(x), & x \in \mathbb{R}, \\ u(x + 1, t) &= u(x, t), & x \in \mathbb{R}, t > 0.\end{aligned}$$

Numerical Methods for PDE: Hyperbolic PDE

Let's use the Fourier analysis (von Neumann stability analysis) to study the stability of these methods.

For simplicity, let's consider a 1-periodic problem rather than a boundary value problem:

$$\begin{aligned}\frac{\partial u}{\partial t} + c \frac{\partial u}{\partial x} &= 0, & x \in \mathbb{R}, t > 0, \\ u(x, 0) &= u_0(x), & x \in \mathbb{R}, \\ u(x+1, t) &= u(x, t), & x \in \mathbb{R}, t > 0.\end{aligned}$$

For fixed $N > 0$, let $h = 1/N$, we take the spatial mesh $h\mathbb{Z} = \{nh : n \in \mathbb{Z}\}$ and we seek a solution $u_n^j \approx u(nh, jk)$ which satisfies the **periodicity condition**

$$u_{n+N}^j = u_n^j, \quad n \in \mathbb{Z}.$$



Let's use the Fourier analysis (von Neumann stability analysis) to study the stability of these methods.

For simplicity, let's consider a 1-periodic problem rather than a boundary value problem:

$$\begin{aligned}\frac{\partial u}{\partial t} + c \frac{\partial u}{\partial x} &= 0, & x \in \mathbb{R}, t > 0, \\ u(x, 0) &= u_0(x), & x \in \mathbb{R}, \\ u(x+1, t) &= u(x, t), & x \in \mathbb{R}, t > 0.\end{aligned}$$

For fixed $N > 0$, let $h = 1/N$, we take the spatial mesh $h\mathbb{Z} = \{nh : n \in \mathbb{Z}\}$ and we seek a solution $u_n^j \approx u(nh, jk)$ which satisfies the **periodicity condition**

$$u_{n+N}^j = u_n^j, \quad n \in \mathbb{Z}.$$

Let L_h^{per} denote the space of complex valued 1-periodic functions on $h\mathbb{Z}$, that is,

$$L_h^{per} = \{v : h\mathbb{Z} \rightarrow \mathbb{C} \mid v((n+N)h) = v(nh+1) = v(nh)\}.$$

Numerical Methods for PDE: Hyperbolic PDE

Let's use the Fourier analysis (von Neumann stability analysis) to study the stability of these methods.

For simplicity, let's consider a 1-periodic problem rather than a boundary value problem:

$$\begin{aligned}\frac{\partial u}{\partial t} + c \frac{\partial u}{\partial x} &= 0, & x \in \mathbb{R}, t > 0, \\ u(x, 0) &= u_0(x), & x \in \mathbb{R}, \\ u(x + 1, t) &= u(x, t), & x \in \mathbb{R}, t > 0.\end{aligned}$$

For fixed $N > 0$, let $h = 1/N$, we take the spatial mesh $h\mathbb{Z} = \{nh : n \in \mathbb{Z}\}$ and we seek a solution $u_n^j \approx u(nh, jk)$ which satisfies the **periodicity condition**

$$u_{n+N}^j = u_n^j, \quad n \in \mathbb{Z}.$$

Let L_h^{per} denote the space of complex valued 1-periodic functions on $h\mathbb{Z}$, that is,

$$L_h^{per} = \{v : h\mathbb{Z} \rightarrow \mathbb{C} \mid v((n+N)h) = v(nh+1) = v(nh)\}.$$

As a basis for this space, we choose

$$\psi_m = \exp(2\pi i m x), \quad x \in h\mathbb{Z}, \quad m = 0, 1, \dots, N-1.$$

Numerical Methods for PDE: Hyperbolic PDE

Let's use the Fourier analysis (von Neumann stability analysis) to study the stability of these methods.

For simplicity, let's consider a 1-periodic problem rather than a boundary value problem:

$$\begin{aligned}\frac{\partial u}{\partial t} + c \frac{\partial u}{\partial x} &= 0, & x \in \mathbb{R}, t > 0, \\ u(x, 0) &= u_0(x), & x \in \mathbb{R}, \\ u(x + 1, t) &= u(x, t), & x \in \mathbb{R}, t > 0.\end{aligned}$$

For fixed $N > 0$, let $h = 1/N$, we take the spatial mesh $h\mathbb{Z} = \{nh : n \in \mathbb{Z}\}$ and we seek a solution $u_n^j \approx u(nh, jk)$ which satisfies the **periodicity condition**

$$u_{n+N}^j = u_n^j, \quad n \in \mathbb{Z}.$$

Let L_h^{per} denote the space of complex valued 1-periodic functions on $h\mathbb{Z}$, that is,

$$L_h^{per} = \{v : h\mathbb{Z} \rightarrow \mathbb{C} \mid v((n+N)h) = v(nh+1) = v(nh)\}.$$

As a basis for this space, we choose

$$\psi_m = \exp(2\pi i m x), \quad x \in h\mathbb{Z}, \quad m = 0, 1, \dots, N-1.$$

The ψ_m are orthogonal with respect to the inner product (**exercise**)

$$\langle \phi, \psi \rangle_h = h \sum_{n=0}^{N-1} \phi(nh) \overline{\psi(nh)}.$$



Numerical Methods for PDE: Hyperbolic PDE

Note that ψ_m is an eigenvector for the forward difference operator D_h^+ , the backward difference operator D_h^- and the centered difference operator D_h . For example,

$$D_h^- \psi_m(x) = \frac{\psi_m(x) - \psi_m(x - h)}{h} = \frac{1 - e^{-2\pi i m h}}{h} \psi_m(x).$$

Numerical Methods for PDE: Hyperbolic PDE

Note that ψ_m is an eigenvector for the forward difference operator D_h^+ , the backward difference operator D_h^- and the centered difference operator D_h . For example,

$$D_h^- \psi_m(x) = \frac{\psi_m(x) - \psi_m(x-h)}{h} = \frac{1 - e^{-2\pi i m h}}{h} \psi_m(x).$$

It is also an eigenvector for the forward and backward shift operators S_h^+ and S_h^- defined as

$$S_h^+ v(x) = v(x+h), \quad S_h^- v(x) = v(x-h),$$

where

$$\begin{aligned} S_h^+ \psi_m(x) &= e^{2\pi i m(x+h)} = e^{2\pi i m h} \psi_m(x), \\ S_h^- \psi_m(x) &= e^{2\pi i m(x-h)} = e^{-2\pi i m h} \psi_m(x). \end{aligned}$$

Numerical Methods for PDE: Hyperbolic PDE

Note that ψ_m is an eigenvector for the forward difference operator D_h^+ , the backward difference operator D_h^- and the centered difference operator D_h . For example,

$$D_h^- \psi_m(x) = \frac{\psi_m(x) - \psi_m(x-h)}{h} = \frac{1 - e^{-2\pi i m h}}{h} \psi_m(x).$$

It is also an eigenvector for the forward and backward shift operators S_h^+ and S_h^- defined as

$$S_h^+ v(x) = v(x+h), \quad S_h^- v(x) = v(x-h),$$

where

$$\begin{aligned} S_h^+ \psi_m(x) &= e^{2\pi i m(x+h)} = e^{2\pi i m h} \psi_m(x), \\ S_h^- \psi_m(x) &= e^{2\pi i m(x-h)} = e^{-2\pi i m h} \psi_m(x). \end{aligned}$$

Now, consider forward-backward difference method in the operator form:

$$u^{j+1} = (1-\lambda)u^j + \lambda S_h^- u^j = ((1-\lambda)I + \lambda S_h^-)u^j.$$

Numerical Methods for PDE: Hyperbolic PDE

Note that ψ_m is an eigenvector for the forward difference operator D_h^+ , the backward difference operator D_h^- and the centered difference operator D_h . For example,

$$D_h^- \psi_m(x) = \frac{\psi_m(x) - \psi_m(x-h)}{h} = \frac{1 - e^{-2\pi i m h}}{h} \psi_m(x).$$

It is also an eigenvector for the forward and backward shift operators S_h^+ and S_h^- defined as

$$S_h^+ v(x) = v(x+h), \quad S_h^- v(x) = v(x-h),$$

where

$$\begin{aligned} S_h^+ \psi_m(x) &= e^{2\pi i m(x+h)} = e^{2\pi i m h} \psi_m(x), \\ S_h^- \psi_m(x) &= e^{2\pi i m(x-h)} = e^{-2\pi i m h} \psi_m(x). \end{aligned}$$

Now, consider forward-backward difference method in the operator form:

$$u^{j+1} = (1 - \lambda)u^j + \lambda S_h^- u^j = ((1 - \lambda)I + \lambda S_h^-)u^j.$$

Thus, for stability of the method, we need that the eigenvalues are less than or equal to one in magnitude, that is

$$|1 - \lambda + \lambda e^{-2\pi i m h}| \leq 1.$$

Numerical Methods for PDE: Hyperbolic PDE

Note that ψ_m is an eigenvector for the forward difference operator D_h^+ , the backward difference operator D_h^- and the centered difference operator D_h . For example,

$$D_h^- \psi_m(x) = \frac{\psi_m(x) - \psi_m(x-h)}{h} = \frac{1 - e^{-2\pi i m h}}{h} \psi_m(x).$$

It is also an eigenvector for the forward and backward shift operators S_h^+ and S_h^- defined as

$$S_h^+ v(x) = v(x+h), \quad S_h^- v(x) = v(x-h),$$

where

$$\begin{aligned} S_h^+ \psi_m(x) &= e^{2\pi i m(x+h)} = e^{2\pi i m h} \psi_m(x), \\ S_h^- \psi_m(x) &= e^{2\pi i m(x-h)} = e^{-2\pi i m h} \psi_m(x). \end{aligned}$$

Now, consider forward-backward difference method in the operator form:

$$u^{j+1} = (1 - \lambda)u^j + \lambda S_h^- u^j = ((1 - \lambda)I + \lambda S_h^-)u^j.$$

Thus, for stability of the method, we need that the eigenvalues are less than or equal to one in magnitude, that is

$$|1 - \lambda + \lambda e^{-2\pi i m h}| \leq 1.$$

As $1 - \lambda + \lambda e^{-2\pi i m h}$ describes a circle centered at $1 - \lambda$ of radius $|\lambda|$, we see that the method is stable if and only if $0 \leq \lambda \leq 1$.