

Initial Value Problems: Linear Multistep Methods

Definition

A linear multistep method satisfies the **root condition** if

- (1) all roots of the first characteristic polynomial has modulus less than or equal to 1, and
- (2) all roots of modulus 1 are simple.

Theorem

The linear multistep method

$$\sum_{j=-1}^k a_j y_{n-j} = h \sum_{j=-1}^k b_j f(t_{n-j}, y_{n-j})$$

is stable only if it satisfies the root condition. If the method satisfies the root condition (and f is Lipschitz continuous), then it is stable.

Proof.

Note that

$$\begin{bmatrix} y_{n-k+1} \\ y_{n-k+2} \\ \vdots \\ y_n \\ y_{n+1} \end{bmatrix} = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ -a_k & -a_{k-1} & -a_{k-2} & \cdots & -a_0 \end{bmatrix} \begin{bmatrix} y_{n-k} \\ y_{n-k+1} \\ \vdots \\ y_{n-1} \\ y_n \end{bmatrix} + \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ h(b_{-1}f_{n+1} + b_0f_n + \cdots + b_kf_{n-k}) \end{bmatrix}$$

Initial Value Problems: Linear Multistep Methods

Proof.

Note that

$$\begin{bmatrix} y_{n-k+1} \\ y_{n-k+2} \\ \vdots \\ y_n \\ y_{n+1} \end{bmatrix} = \begin{bmatrix} 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & 1 \\ -a_k & -a_{k-1} & -a_{k-2} & \dots & -a_0 \end{bmatrix} \begin{bmatrix} y_{n-k} \\ y_{n-k+1} \\ \vdots \\ y_{n-1} \\ y_n \end{bmatrix} + \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ h(b_{-1}f_{n+1} + b_0f_n + \dots + b_kf_{n-k}) \end{bmatrix}$$

For solutions y_j and \hat{y}_j , let

$$E_n = \begin{bmatrix} y_{n-k} - \hat{y}_{n-k} \\ y_{n-k+1} - \hat{y}_{n-k+1} \\ \vdots \\ y_{n-1} - \hat{y}_{n-1} \\ y_n - \hat{y}_n \end{bmatrix}$$

Initial Value Problems: Linear Multistep Methods

Proof.

Note that

$$\begin{bmatrix} y_{n-k+1} \\ y_{n-k+2} \\ \vdots \\ y_n \\ y_{n+1} \end{bmatrix} = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ -a_k & -a_{k-1} & -a_{k-2} & \cdots & -a_0 \end{bmatrix} \begin{bmatrix} y_{n-k} \\ y_{n-k+1} \\ \vdots \\ y_{n-1} \\ y_n \end{bmatrix} + \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ h(b_{-1}f_{n+1} + b_0f_n + \cdots + b_kf_{n-k}) \end{bmatrix}$$

For solutions y_j and \hat{y}_j , let

$$E_n = \begin{bmatrix} y_{n-k} - \hat{y}_{n-k} \\ y_{n-k+1} - \hat{y}_{n-k+1} \\ \vdots \\ y_{n-1} - \hat{y}_{n-1} \\ y_n - \hat{y}_n \end{bmatrix}, \quad A = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ -a_k & -a_{k-1} & -a_{k-2} & \cdots & -a_0 \end{bmatrix}$$

and

$$Q_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ h(b_{-1}(f(t_{n+1}, y_{n+1}) - f(t_{n+1}, \hat{y}_{n+1})) + \cdots + b_k(f(t_{n-k}, y_{n-k}) - f(t_{n-k}, \hat{y}_{n-k}))) \end{bmatrix}$$

to get

$$E_{n+1} = AE_n + Q_n$$

Initial Value Problems: Linear Multistep Methods

Proof.

For solutions y_j and \hat{y}_j , let

$$E_n = \begin{bmatrix} y_{n-k} - \hat{y}_{n-k} \\ y_{n-k+1} - \hat{y}_{n-k+1} \\ \vdots \\ y_{n-1} - \hat{y}_{n-1} \\ y_n - \hat{y}_n \end{bmatrix}, \quad A = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ -a_k & -a_{k-1} & -a_{k-2} & \cdots & -a_0 \end{bmatrix}$$

and

$$Q_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ h(b_{-1}(f(t_{n+1}, y_{n+1}) - f(t_{n+1}, \hat{y}_{n+1})) + \cdots + b_k(f(t_{n-k}, y_{n-k}) - f(t_{n-k}, \hat{y}_{n-k}))) \end{bmatrix}$$

to get

$$E_{n+1} = AE_n + Q_n.$$

Note that $|\lambda I - A| = \rho(\lambda)$. (Why?)

Initial Value Problems: Linear Multistep Methods

Proof.

For solutions y_j and \hat{y}_j , let

$$E_n = \begin{bmatrix} y_{n-k} - \hat{y}_{n-k} \\ y_{n-k+1} - \hat{y}_{n-k+1} \\ \vdots \\ y_{n-1} - \hat{y}_{n-1} \\ y_n - \hat{y}_n \end{bmatrix}, \quad A = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ -a_k & -a_{k-1} & -a_{k-2} & \cdots & -a_0 \end{bmatrix}$$

and

$$Q_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ h(b_{-1}(f(t_{n+1}, y_{n+1}) - f(t_{n+1}, \hat{y}_{n+1})) + \cdots + b_k(f(t_{n-k}, y_{n-k}) - f(t_{n-k}, \hat{y}_{n-k}))) \end{bmatrix}$$

to get

$$E_{n+1} = AE_n + Q_n.$$

Thus, we have

$$E_{k+n} = A^n E_k + \sum_{j=0}^{n-1} A^{n-1-j} Q_{k+j}.$$

Initial Value Problems: Linear Multistep Methods

Proof.

For solutions y_j and \hat{y}_j , let

$$E_n = \begin{bmatrix} y_{n-k} - \hat{y}_{n-k} \\ y_{n-k+1} - \hat{y}_{n-k+1} \\ \vdots \\ y_{n-1} - \hat{y}_{n-1} \\ y_n - \hat{y}_n \end{bmatrix}, \quad A = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ -a_k & -a_{k-1} & -a_{k-2} & \cdots & -a_0 \end{bmatrix}$$

and

$$Q_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ h(b_{-1}(f(t_{n+1}, y_{n+1}) - f(t_{n+1}, \hat{y}_{n+1})) + \cdots + b_k(f(t_{n-k}, y_{n-k}) - f(t_{n-k}, \hat{y}_{n-k}))) \end{bmatrix}$$

to get

$$E_{k+n} = A^n E_k + \sum_{j=0}^{n-1} A^{n-1-j} Q_{k+j}.$$

Initial Value Problems: Linear Multistep Methods

Proof.

For solutions y_j and \hat{y}_j , let

$$E_n = \begin{bmatrix} y_{n-k} - \hat{y}_{n-k} \\ y_{n-k+1} - \hat{y}_{n-k+1} \\ \vdots \\ y_{n-1} - \hat{y}_{n-1} \\ y_n - \hat{y}_n \end{bmatrix}, \quad A = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ -a_k & -a_{k-1} & -a_{k-2} & \cdots & -a_0 \end{bmatrix}$$

and

$$Q_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ h(b_{-1}(f(t_{n+1}, y_{n+1}) - f(t_{n+1}, \hat{y}_{n+1})) + \cdots + b_k(f(t_{n-k}, y_{n-k}) - f(t_{n-k}, \hat{y}_{n-k}))) \end{bmatrix}$$

to get

$$E_{k+n} = A^n E_k + \sum_{j=0}^{n-1} A^{n-1-j} Q_{k+j}.$$

Thus, using ℓ_∞ norm for vectors and the fact that there is a constant C so that $\|A^m\| \leq C$, for all m , we have

$$\|E_{k+n}\| \leq C\|E_k\| + C \sum_{j=0}^{n-1} \|Q_{k+j}\|$$

Initial Value Problems: Linear Multistep Methods

Proof.

For solutions y_j and \hat{y}_j , let

$$E_n = \begin{bmatrix} y_{n-k} - \hat{y}_{n-k} \\ y_{n-k+1} - \hat{y}_{n-k+1} \\ \vdots \\ y_{n-1} - \hat{y}_{n-1} \\ y_n - \hat{y}_n \end{bmatrix}, \quad A = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ -a_k & -a_{k-1} & -a_{k-2} & \cdots & -a_0 \end{bmatrix}$$

and

$$Q_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ h(b_{-1}(f(t_{n+1}, y_{n+1}) - f(t_{n+1}, \hat{y}_{n+1})) + \cdots + b_k(f(t_{n-k}, y_{n-k}) - f(t_{n-k}, \hat{y}_{n-k}))) \end{bmatrix}$$

to get

$$E_{k+n} = A^n E_k + \sum_{j=0}^{n-1} A^{n-1-j} Q_{k+j}.$$

Thus, using ℓ_∞ norm for vectors and the fact that there is a constant C so that $\|A^m\| \leq C$, for all m , we have

$$\|E_{k+n}\| \leq C\|E_k\| + C \sum_{j=0}^{n-1} \|Q_{k+j}\| \leq C\|E_k\| + hC\|b\|_1 L \sum_{j=0}^{n-1} (\|E_{k+j}\| + \|E_{k+j+1}\|)$$

Initial Value Problems: Linear Multistep Methods

Proof.

For solutions y_j and \hat{y}_j , let

$$E_n = \begin{bmatrix} y_{n-k} - \hat{y}_{n-k} \\ y_{n-k+1} - \hat{y}_{n-k+1} \\ \vdots \\ y_{n-1} - \hat{y}_{n-1} \\ y_n - \hat{y}_n \end{bmatrix}, \quad A = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ -a_k & -a_{k-1} & -a_{k-2} & \cdots & -a_0 \end{bmatrix}$$

and

$$Q_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ h(b_{-1}(f(t_{n+1}, y_{n+1}) - f(t_{n+1}, \hat{y}_{n+1})) + \cdots + b_k(f(t_{n-k}, y_{n-k}) - f(t_{n-k}, \hat{y}_{n-k}))) \end{bmatrix}$$

to get

$$\|E_{k+n}\| \leq C\|E_k\| + C \sum_{j=0}^{n-1} \|Q_{k+j}\| \leq C\|E_k\| + hC\|b\|_1 L \sum_{j=0}^{n-1} (\|E_{k+j}\| + \|E_{k+j+1}\|)$$

Initial Value Problems: Linear Multistep Methods

Proof.

For solutions y_j and \hat{y}_j , let

$$E_n = \begin{bmatrix} y_{n-k} - \hat{y}_{n-k} \\ y_{n-k+1} - \hat{y}_{n-k+1} \\ \vdots \\ y_{n-1} - \hat{y}_{n-1} \\ y_n - \hat{y}_n \end{bmatrix}, \quad A = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ -a_k & -a_{k-1} & -a_{k-2} & \cdots & -a_0 \end{bmatrix}$$

and

$$Q_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ h(b_{-1}(f(t_{n+1}, y_{n+1}) - f(t_{n+1}, \hat{y}_{n+1})) + \cdots + b_k(f(t_{n-k}, y_{n-k}) - f(t_{n-k}, \hat{y}_{n-k}))) \end{bmatrix}$$

to get

$$\begin{aligned} \|E_{k+n}\| &\leq C\|E_k\| + C \sum_{j=0}^{n-1} \|Q_{k+j}\| \leq C\|E_k\| + hC\|b\|_1 L \sum_{j=0}^{n-1} (\|E_{k+j}\| + \|E_{k+j+1}\|) \\ \|E_{k+n}\| &\leq C\|E_k\| + 2hC\|b\|_1 L \sum_{j=1}^{n-1} \|E_{k+j}\| + hC\|b\|_1 L\|E_k\| + hC\|b\|_1 L\|E_{k+n}\| \end{aligned}$$

Initial Value Problems: Linear Multistep Methods

Proof.

For solutions y_j and \hat{y}_j , let

$$E_n = \begin{bmatrix} y_{n-k} - \hat{y}_{n-k} \\ y_{n-k+1} - \hat{y}_{n-k+1} \\ \vdots \\ y_{n-1} - \hat{y}_{n-1} \\ y_n - \hat{y}_n \end{bmatrix}, \quad A = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ -a_k & -a_{k-1} & -a_{k-2} & \cdots & -a_0 \end{bmatrix}$$

and

$$Q_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ h(b_{-1}(f(t_{n+1}, y_{n+1}) - f(t_{n+1}, \hat{y}_{n+1})) + \cdots + b_k(f(t_{n-k}, y_{n-k}) - f(t_{n-k}, \hat{y}_{n-k}))) \end{bmatrix}$$

to get

$$\begin{aligned} \|E_{k+n}\| &\leq C\|E_k\| + C \sum_{j=0}^{n-1} \|Q_{k+j}\| \leq C\|E_k\| + hC\|b\|_1 L \sum_{j=0}^{n-1} (\|E_{k+j}\| + \|E_{k+j+1}\|) \\ \|E_{k+n}\| &\leq C\|E_k\| + 2hC\|b\|_1 L \sum_{j=1}^{n-1} \|E_{k+j}\| + hC\|b\|_1 L \|E_k\| + hC\|b\|_1 L \|E_{k+n}\| \\ (1 - hC\|b\|_1 L) \|E_{k+n}\| &\leq C\|E_k\| + 2hC\|b\|_1 L \sum_{j=0}^{n-1} \|E_{k+j}\| \end{aligned}$$

Initial Value Problems: Linear Multistep Methods

Proof.

For solutions y_j and \hat{y}_j , let

$$E_n = \begin{bmatrix} y_{n-k} - \hat{y}_{n-k} \\ y_{n-k+1} - \hat{y}_{n-k+1} \\ \vdots \\ y_{n-1} - \hat{y}_{n-1} \\ y_n - \hat{y}_n \end{bmatrix}, \quad A = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ -a_k & -a_{k-1} & -a_{k-2} & \cdots & -a_0 \end{bmatrix}$$

and

$$Q_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ h(b_{-1}(f(t_{n+1}, y_{n+1}) - f(t_{n+1}, \hat{y}_{n+1})) + \cdots + b_k(f(t_{n-k}, y_{n-k}) - f(t_{n-k}, \hat{y}_{n-k}))) \end{bmatrix}$$

to get

$$(1 - hC\|b\|_1 L)\|E_{k+n}\| \leq C\|E_k\| + 2hC\|b\|_1 L \sum_{j=0}^{n-1} \|E_{k+j}\|.$$

Initial Value Problems: Linear Multistep Methods

Proof.

For solutions y_j and \hat{y}_j , let

$$E_n = \begin{bmatrix} y_{n-k} - \hat{y}_{n-k} \\ y_{n-k+1} - \hat{y}_{n-k+1} \\ \vdots \\ y_{n-1} - \hat{y}_{n-1} \\ y_n - \hat{y}_n \end{bmatrix}, \quad A = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ -a_k & -a_{k-1} & -a_{k-2} & \cdots & -a_0 \end{bmatrix}$$

and

$$Q_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ h(b_{-1}(f(t_{n+1}, y_{n+1}) - f(t_{n+1}, \hat{y}_{n+1})) + \cdots + b_k(f(t_{n-k}, y_{n-k}) - f(t_{n-k}, \hat{y}_{n-k}))) \end{bmatrix}$$

to get

$$(1 - hC\|b\|_1 L)\|E_{k+n}\| \leq C\|E_k\| + 2hC\|b\|_1 L \sum_{j=0}^{n-1} \|E_{k+j}\|.$$

Thus, for $h \leq (2C\|b\|_1 L)^{-1}$, we have

$$\|E_{k+n}\| \leq \frac{C\|E_k\|}{(1 - hC\|b\|_1 L)} + \frac{2hC\|b\|_1 L (\sum_{j=0}^{n-1} \|E_{k+j}\|)}{(1 - hC\|b\|_1 L)}$$

Initial Value Problems: Linear Multistep Methods

Proof.

For solutions y_j and \hat{y}_j , let

$$E_n = \begin{bmatrix} y_{n-k} - \hat{y}_{n-k} \\ y_{n-k+1} - \hat{y}_{n-k+1} \\ \vdots \\ y_{n-1} - \hat{y}_{n-1} \\ y_n - \hat{y}_n \end{bmatrix}, \quad A = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ -a_k & -a_{k-1} & -a_{k-2} & \cdots & -a_0 \end{bmatrix}$$

and

$$Q_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ h(b_{-1}(f(t_{n+1}, y_{n+1}) - f(t_{n+1}, \hat{y}_{n+1})) + \cdots + b_k(f(t_{n-k}, y_{n-k}) - f(t_{n-k}, \hat{y}_{n-k}))) \end{bmatrix}$$

to get

$$(1 - hC\|b\|_1 L)\|E_{k+n}\| \leq C\|E_k\| + 2hC\|b\|_1 L \sum_{j=0}^{n-1} \|E_{k+j}\|.$$

Thus, for $h \leq (2C\|b\|_1 L)^{-1}$, we have

$$\|E_{k+n}\| \leq \frac{C\|E_k\|}{(1 - hC\|b\|_1 L)} + \frac{2hC\|b\|_1 L (\sum_{j=0}^{n-1} \|E_{k+j}\|)}{(1 - hC\|b\|_1 L)} \leq 2C\|E_k\| + 4hC\|b\|_1 L \left(\sum_{j=0}^{n-1} \|E_{k+j}\| \right)$$

Initial Value Problems: Linear Multistep Methods

Proof.

For solutions y_j and \hat{y}_j , let

$$E_n = \begin{bmatrix} y_{n-k} - \hat{y}_{n-k} \\ y_{n-k+1} - \hat{y}_{n-k+1} \\ \vdots \\ y_{n-1} - \hat{y}_{n-1} \\ y_n - \hat{y}_n \end{bmatrix}, \quad A = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ -a_k & -a_{k-1} & -a_{k-2} & \cdots & -a_0 \end{bmatrix}$$

and

$$Q_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ h(b_{-1}(f(t_{n+1}, y_{n+1}) - f(t_{n+1}, \hat{y}_{n+1})) + \cdots + b_k(f(t_{n-k}, y_{n-k}) - f(t_{n-k}, \hat{y}_{n-k}))) \end{bmatrix}$$

Thus, for $h \leq (2C\|b\|_1 L)^{-1}$, we have

$$\|E_{k+n}\| \leq 2C\|E_k\| + 4hC\|b\|_1 L \left(\sum_{j=0}^{n-1} \|E_{k+j}\| \right).$$

Proof.

For solutions y_j and \hat{y}_j , let

$$E_n = \begin{bmatrix} y_{n-k} - \hat{y}_{n-k} \\ y_{n-k+1} - \hat{y}_{n-k+1} \\ \vdots \\ y_{n-1} - \hat{y}_{n-1} \\ y_n - \hat{y}_n \end{bmatrix}, \quad A = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ -a_k & -a_{k-1} & -a_{k-2} & \cdots & -a_0 \end{bmatrix}$$

and

$$Q_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ h(b_{-1}(f(t_{n+1}, y_{n+1}) - f(t_{n+1}, \hat{y}_{n+1})) + \cdots + b_k(f(t_{n-k}, y_{n-k}) - f(t_{n-k}, \hat{y}_{n-k}))) \end{bmatrix}$$

Thus, for $h \leq (2C\|b\|_1 L)^{-1}$, we have

$$\|E_{k+n}\| \leq 2C\|E_k\| + 4hC\|b\|_1 L \left(\sum_{j=0}^{n-1} \|E_{k+j}\| \right).$$

$$\|E_{k+n}\| \leq 2C\|E_k\|(1 + 4hC\|b\|_1 L)^n$$

Initial Value Problems: Linear Multistep Methods

Proof.

For solutions y_j and \hat{y}_j , let

$$E_n = \begin{bmatrix} y_{n-k} - \hat{y}_{n-k} \\ y_{n-k+1} - \hat{y}_{n-k+1} \\ \vdots \\ y_{n-1} - \hat{y}_{n-1} \\ y_n - \hat{y}_n \end{bmatrix}, \quad A = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ -a_k & -a_{k-1} & -a_{k-2} & \cdots & -a_0 \end{bmatrix}$$

and

$$Q_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ h(b_{-1}(f(t_{n+1}, y_{n+1}) - f(t_{n+1}, \hat{y}_{n+1})) + \cdots + b_k(f(t_{n-k}, y_{n-k}) - f(t_{n-k}, \hat{y}_{n-k}))) \end{bmatrix}$$

Thus, for $h \leq (2C\|b\|_1 L)^{-1}$, we have

$$\|E_{k+n}\| \leq 2C\|E_k\| + 4hC\|b\|_1 L \left(\sum_{j=0}^{n-1} \|E_{k+j}\| \right).$$

$$\|E_{k+n}\| \leq 2C\|E_k\|(1 + 4hC\|b\|_1 L)^n \leq 2C\|E_k\|(1 + 4hC\|b\|_1 L)^N$$

Proof.

For solutions y_j and \hat{y}_j , let

$$E_n = \begin{bmatrix} y_{n-k} - \hat{y}_{n-k} \\ y_{n-k+1} - \hat{y}_{n-k+1} \\ \vdots \\ y_{n-1} - \hat{y}_{n-1} \\ y_n - \hat{y}_n \end{bmatrix}, \quad A = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ -a_k & -a_{k-1} & -a_{k-2} & \cdots & -a_0 \end{bmatrix}$$

and

$$Q_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ h(b_{-1}(f(t_{n+1}, y_{n+1}) - f(t_{n+1}, \hat{y}_{n+1})) + \cdots + b_k(f(t_{n-k}, y_{n-k}) - f(t_{n-k}, \hat{y}_{n-k}))) \end{bmatrix}$$

Thus, for $h \leq (2C\|b\|_1 L)^{-1}$, we have

$$\|E_{k+n}\| \leq 2C\|E_k\| + 4hC\|b\|_1 L \left(\sum_{j=0}^{n-1} \|E_{k+j}\| \right).$$

$$\|E_{k+n}\| \leq 2C\|E_k\|(1 + 4hC\|b\|_1 L)^n \leq 2C\|E_k\|(1 + 4hC\|b\|_1 L)^N \leq 2C\|E_k\|e^{4NhC\|b\|_1 L}$$

Initial Value Problems: Linear Multistep Methods

Proof.

For solutions y_j and \hat{y}_j , let

$$E_n = \begin{bmatrix} y_{n-k} - \hat{y}_{n-k} \\ y_{n-k+1} - \hat{y}_{n-k+1} \\ \vdots \\ y_{n-1} - \hat{y}_{n-1} \\ y_n - \hat{y}_n \end{bmatrix}, \quad A = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ -a_k & -a_{k-1} & -a_{k-2} & \cdots & -a_0 \end{bmatrix}$$

and

$$Q_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ h(b_{-1}(f(t_{n+1}, y_{n+1}) - f(t_{n+1}, \hat{y}_{n+1})) + \cdots + b_k(f(t_{n-k}, y_{n-k}) - f(t_{n-k}, \hat{y}_{n-k}))) \end{bmatrix}$$

Thus, for $h \leq (2C\|b\|_1 L)^{-1}$, we have

$$\|E_{k+n}\| \leq 2C\|E_k\| + 4hC\|b\|_1 L \left(\sum_{j=0}^{n-1} \|E_{k+j}\| \right).$$

$$\|E_{k+n}\| \leq 2C\|E_k\|(1 + 4hC\|b\|_1 L)^n \leq 2C\|E_k\|(1 + 4hC\|b\|_1 L)^N \leq 2C\|E_k\|e^{4NhC\|b\|_1 L} \leq Ce^{4(T-t_0)C\|b\|_1 L}\|E_k\|.$$

Initial Value Problems: Linear Multistep Methods

Proof.

For solutions y_j and \hat{y}_j , let

$$E_n = \begin{bmatrix} y_{n-k} - \hat{y}_{n-k} \\ y_{n-k+1} - \hat{y}_{n-k+1} \\ \vdots \\ y_{n-1} - \hat{y}_{n-1} \\ y_n - \hat{y}_n \end{bmatrix}, \quad A = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ -a_k & -a_{k-1} & -a_{k-2} & \cdots & -a_0 \end{bmatrix}$$

and

$$Q_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ h(b_{-1}(f(t_{n+1}, y_{n+1}) - f(t_{n+1}, \hat{y}_{n+1})) + \cdots + b_k(f(t_{n-k}, y_{n-k}) - f(t_{n-k}, \hat{y}_{n-k}))) \end{bmatrix}$$

Thus, for $h \leq (2C\|b\|_1 L)^{-1}$, we have

$$\|E_{k+n}\| \leq 2C\|E_k\| + 4hC\|b\|_1 L \left(\sum_{j=0}^{n-1} \|E_{k+j}\| \right).$$

$$\|E_{k+n}\| \leq 2C\|E_k\|(1 + 4hC\|b\|_1 L)^n \leq 2C\|E_k\|(1 + 4hC\|b\|_1 L)^N \leq 2C\|E_k\|e^{4NhC\|b\|_1 L} \leq Ce^{4(T-t_0)C\|b\|_1 L}\|E_k\|.$$

So stability follows.

Numerical Analysis & Scientific Computing II

Module 2

Initial Value Problems

2.4 Implicit method

2.5 Stiffness

2.6 Linear Multistep Methods

- Consistency, stability and convergence



Akash Anand
MATH, IIT KANPUR

Initial Value Problems: Linear Multistep Methods



Theorem

The linear multistep method is convergent if and only if it is consistent and stable.

Initial Value Problems: Linear Multistep Methods



Theorem

The linear multistep method is convergent if and only if it is consistent and stable.

Remark

This theorem states that it can be determined whether or not a linear multistep method is convergent simply by checking some algebraic conditions concerning its characteristic polynomials.

Initial Value Problems: Linear Multistep Methods

Theorem

The linear multistep method is convergent if and only if it is consistent and stable.

Remark

This theorem states that it can be determined whether or not a linear multistep method is convergent simply by checking some algebraic conditions concerning its characteristic polynomials.

Proof. (sketch)

Convergence \Rightarrow Consistency

Convergence \Rightarrow Stability

Consistency and Stability \Rightarrow Convergence

Initial Value Problems: Linear Multistep Methods

Theorem

The linear multistep method is convergent if and only if it is consistent and stable.

Remark

This theorem states that it can be determined whether or not a linear multistep method is convergent simply by checking some algebraic conditions concerning its characteristic polynomials.

Proof. (sketch)

Convergence \Rightarrow Consistency

Apply the method to $y' = 0, y(0) = 1$ and $y' = 1, y(0) = 0$ for verifying satisfiability of the consistency conditions.

Convergence \Rightarrow Stability

Consistency and Stability \Rightarrow Convergence

Initial Value Problems: Linear Multistep Methods

Theorem

The linear multistep method is convergent if and only if it is consistent and stable.

Remark

This theorem states that it can be determined whether or not a linear multistep method is convergent simply by checking some algebraic conditions concerning its characteristic polynomials.

Proof. (sketch)

Convergence \Rightarrow Consistency

Apply the method to $y' = 0, y(0) = 1$ and $y' = 1, y(0) = 0$ for verifying satisfiability of the consistency conditions.

Convergence \Rightarrow Stability

Apply the method to $y' = 0, y(0) = 0$ for verifying satisfiability of the root condition.

Consistency and Stability \Rightarrow Convergence