
MTH 424 - PARTIAL DIFFERENTIAL EQUATION

IIT KANPUR

Instructor: Indranil Chowdhury

Odd Semester, 2024-25

Assignment Prep

1. Find the eigenvalues and corresponding eigenfunctions of the following problems:

(i) $-\frac{d^2\phi}{dx^2} = \lambda\phi$ in $(0, 1)$ with $\phi(0) = 0$ and $\phi(1) = 0$.

(ii) $-\frac{d^2\phi}{dx^2} = \lambda\phi$ in $(0, 1)$ with $\phi(0) = 0$ and $\frac{d\phi}{dx}(1) = 0$.

2. Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ be defined by

$$f(x, y) = \begin{cases} \frac{xy^2}{x^2+y^4} & \text{if } x \neq 0, \\ 0 & \text{if } x = 0. \end{cases}$$

Show that directional derivative of f at $(0, 0)$ exists for all direction (i.e. $D_v f(0, 0)$ exists for all $v \in \mathbb{R}^2$). Check whether f is continuous at $(0, 0)$.

3. **Support of a function:** A support of a function f is a closed set defined as the closure of the set $\{x : f(x) \neq 0\}$. We denote the set by $\text{supp}(f)$.

Compactly Supported function: A function is said to be compactly supported if $\text{supp}(f)$ is a compact set.

Consider the function $\rho : \mathbb{R}^2 \rightarrow \mathbb{R}$ defined by

$$\rho = \begin{cases} c \exp\left(\frac{1}{1-|x|^2}\right) & \text{if } |x| < 1, \\ 0 & \text{if } |x| \geq 1, \end{cases}$$

where the constant c is chosen such a way that $\int_{\mathbb{R}^2} \rho(x) dx = 1$. Verify $\text{supp}(\rho) = \overline{B_1(0)}$ and $\rho \in C^\infty(\mathbb{R}^2)$. Define ρ_ϵ as

$$\rho_\epsilon(x) = \frac{1}{\epsilon^2} \rho\left(\frac{x}{\epsilon}\right).$$

Show that $\rho_\epsilon \in C^\infty(\mathbb{R}^2)$, $\text{supp}(\rho_\epsilon) = \overline{B_\epsilon(0)}$ and $\int_{\mathbb{R}^2} \rho_\epsilon(x) dx = 1$.

(Similarly, ρ and ρ_ϵ can be defined in \mathbb{R}^n)

4. Find directional derivative of the function $f(x, y, z) = xyz$ in the direction of the vector $v = (5, -3, 2)$.
5. Let $f(x, y) = \tan(\frac{x^2}{y})$; $y \neq 0$. Verify the mixed derivatives f_{xy} and f_{yx} are equal.
6. Let R be the region lying inside $x^2 + y^2 = 2x$ and to the right $x = 1$. Use polar co-ordinate to compute the following integration:

$$\iint_R \frac{x dx dy}{x^2 + y^2}.$$

7. Let $g_1, g_2 : \mathbb{R}^2 \rightarrow \mathbb{R}$ be continuously differentiable and suppose that $\partial_x g_2 = \partial_y g_1$. Let

$$f(x, y) = \int_0^x g_1(t, 0) dt + \int_0^y g_2(x, t) dt$$

Show that $\partial_x f(x, y) = g_1(x, y)$.