

# Boundary Value Problems: Finite Difference Method

This yields a system of algebraic equations

$$\frac{u_{i+1} - 2u_i + u_{i-1}}{h^2} = f\left(t_i, u_i, \frac{u_{i+1} - u_{i-1}}{2h}\right), i = 1, \dots, n,$$

to be solved for the unknowns  $u_i, i = 1, \dots, n$ .

In the matrix form, we have

$$\frac{1}{h^2} \begin{bmatrix} -2 & 1 & 0 & & \cdots & 0 \\ 1 & -2 & 1 & & & \\ \vdots & \vdots & & \ddots & \vdots & \\ & & & & 1 & -2 & 1 \\ 0 & \dots & & & 0 & 1 & -2 \end{bmatrix} \begin{bmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{bmatrix} = \begin{bmatrix} f\left(t_1, u_1, \frac{u_2 - \alpha}{2h}\right) \\ f\left(t_2, u_2, \frac{u_3 - u_1}{2h}\right) \\ \vdots \\ f\left(t_n, u_n, \frac{\beta - u_n}{2h}\right) \end{bmatrix} - \begin{bmatrix} \frac{\alpha}{h^2} \\ \vdots \\ \frac{\beta}{h^2} \end{bmatrix}$$

which is denoted as

$$\frac{1}{h^2} Au = F(u) + g.$$

Thus, the Newton's method for solving the system of algebraic equations is given by

$$u^{(m+1)} = u^{(m)} - \left[ \frac{1}{h^2} A - F'(u^{(m)}) \right]^{-1} \left[ \frac{1}{h^2} Au^{(m)} - F(u^{(m)}) - g \right]$$

where the Jacobian matrix is given by  $[F(u)]_{ij} = [\partial f(t_i, u_i, (u_{i+1} - u_{i-1})/(2h))/\partial u_j]$ .

# Boundary Value Problems: Finite Difference Method

In the matrix form, we have

$$\frac{1}{h^2} Au = F(u) + g.$$

Thus, the Newton's method for solving the system of algebraic equations is given by

$$u^{(m+1)} = u^{(m)} - \left[ \frac{1}{h^2} A - F'(u^{(m)}) \right]^{-1} \left[ \frac{1}{h^2} Au^{(m)} - F(u^{(m)}) - g \right]$$

where the Jacobian matrix is given by  $[F'(u)]_{ij} = [\partial f(t_i, u_i, (u_{i+1} - u_{i-1})/(2h))/\partial u_j]$ .

In particular,

$$\begin{aligned} [F'(u)]_{ii} &= f_2 \left( t_i, u_i, \frac{u_{i+1} - u_{i-1}}{2h} \right), & 1 \leq i \leq n, \\ [F'(u)]_{i,i-1} &= -\frac{1}{2h} f_3 \left( t_i, u_i, \frac{u_{i+1} - u_{i-1}}{2h} \right), & 2 \leq i \leq n, \\ [F'(u)]_{i,i+1} &= \frac{1}{2h} f_3 \left( t_i, u_i, \frac{u_{i+1} - u_{i-1}}{2h} \right), & 1 \leq i \leq n-1, \end{aligned}$$

where all other entries of  $F'(u)$  are 0 and  $f_2(t, u, v)$ ,  $f_3(t, u, v)$  denote the partial derivatives of  $f$  with respect to  $u$  and  $v$  respectively.

# Boundary Value Problems: Shooting Method

## Example

Consider the two-point BVP

$$u'' = -u + \frac{2(u')^2}{u}, \quad -1 < t < 1,$$

$$u(-1) = u(1) = (e + e^{-1})^{-1}.$$

The iterative solution via Newton's method satisfies

$$u^{(m+1)} = u^{(m)} - \left[ \frac{1}{h^2} A - F'(u^{(m)}) \right]^{-1} \left[ \frac{1}{h^2} A u^{(m)} - F(u^{(m)}) - g \right],$$

where

$$[F'(u)]_{ii} =$$

$$[F'(u)]_{i,i-1} =$$

$$[F'(u)]_{i,i+1} =$$

and

$$g =$$

# Boundary Value Problems: Shooting Method

## Example

Consider the two-point BVP

$$u'' = -u + \frac{2(u')^2}{u}, \quad -1 < t < 1,$$

$$u(-1) = u(1) = (e + e^{-1})^{-1}.$$

The iterative solution via Newton's method satisfies

$$u^{(m+1)} = u^{(m)} - \left[ \frac{1}{h^2} A - F'(u^{(m)}) \right]^{-1} \left[ \frac{1}{h^2} A u^{(m)} - F(u^{(m)}) - g \right],$$

where

$$[F'(u)]_{ii} = -1 - 2 \left( \frac{u_{i+1} - u_{i-1}}{2hu_i} \right)^2, \quad [F'(u)]_{i,i-1} =$$

$$[F'(u)]_{i,i+1} =$$

and

$$g =$$

# Boundary Value Problems: Shooting Method

## Example

Consider the two-point BVP

$$u'' = -u + \frac{2(u')^2}{u}, \quad -1 < t < 1,$$

$$u(-1) = u(1) = (e + e^{-1})^{-1}.$$

The iterative solution via Newton's method satisfies

$$u^{(m+1)} = u^{(m)} - \left[ \frac{1}{h^2} A - F'(u^{(m)}) \right]^{-1} \left[ \frac{1}{h^2} A u^{(m)} - F(u^{(m)}) - g \right],$$

where

$$[F'(u)]_{ii} = -1 - 2 \left( \frac{u_{i+1} - u_{i-1}}{2hu_i} \right)^2, \quad [F'(u)]_{i,i-1} = -\frac{1}{2h} \left( 4 \frac{u_{i+1} - u_{i-1}}{2hu_i} \right),$$

$$[F'(u)]_{i,i+1} =$$

and

$$g =$$

# Boundary Value Problems: Shooting Method

## Example

Consider the two-point BVP

$$u'' = -u + \frac{2(u')^2}{u}, \quad -1 < t < 1,$$

$$u(-1) = u(1) = (e + e^{-1})^{-1}.$$

The iterative solution via Newton's method satisfies

$$u^{(m+1)} = u^{(m)} - \left[ \frac{1}{h^2} A - F'(u^{(m)}) \right]^{-1} \left[ \frac{1}{h^2} A u^{(m)} - F(u^{(m)}) - g \right],$$

where

$$[F'(u)]_{ii} = -1 - 2 \left( \frac{u_{i+1} - u_{i-1}}{2hu_i} \right)^2, \quad [F'(u)]_{i,i-1} = -\frac{1}{2h} \left( 4 \frac{u_{i+1} - u_{i-1}}{2hu_i} \right),$$

$$[F'(u)]_{i,i+1} = \frac{1}{2h} \left( 4 \frac{u_{i+1} - u_{i-1}}{2hu_i} \right),$$

and

$$g =$$

# Boundary Value Problems: Shooting Method

## Example

Consider the two-point BVP

$$u'' = -u + \frac{2(u')^2}{u}, \quad -1 < t < 1,$$

$$u(-1) = u(1) = (e + e^{-1})^{-1}.$$

The iterative solution via Newton's method satisfies

$$u^{(m+1)} = u^{(m)} - \left[ \frac{1}{h^2} A - F'(u^{(m)}) \right]^{-1} \left[ \frac{1}{h^2} A u^{(m)} - F(u^{(m)}) - g \right],$$

where

$$[F'(u)]_{ii} = -1 - 2 \left( \frac{u_{i+1} - u_{i-1}}{2hu_i} \right)^2, \quad [F'(u)]_{i,i-1} = -\frac{1}{2h} \left( 4 \frac{u_{i+1} - u_{i-1}}{2hu_i} \right),$$

$$[F'(u)]_{i,i+1} = \frac{1}{2h} \left( 4 \frac{u_{i+1} - u_{i-1}}{2hu_i} \right),$$

and

$$g = \frac{1}{h^2} \begin{bmatrix} (e + e^{-1})^{-1} \\ 0 \\ \vdots \\ 0 \\ (e + e^{-1})^{-1} \end{bmatrix}.$$

# *Numerical Analysis & Scientific Computing II*

## *Lesson 3*

# *Boundary Value Problems for ODEs*

*3.1 Well-posedness*

*3.2 Shooting Method*

**3.3 Finite Difference Method**

**- Error Analysis**



**Akash Anand**  
MATH, IIT KANPUR



# Boundary Value Problems: Finite Difference Method



*The error analysis in full generality is too complicated to be discussed in this course. We will, therefore, look at simpler situation where the right hand side is independent of  $u$  and  $u'$ , that is,  $f(t, u, u') = f(t)$ .*



# Boundary Value Problems: Finite Difference Method

*The error analysis in full generality is too complicated to be discussed in this course. We will, therefore, look at simpler situation where the right hand side is independent of  $u$  and  $u'$ , that is,  $f(t, u, u') = f(t)$ .*

*The model BVP, therefore, reads*

$$\begin{aligned} u'' &= f(t), & a < t < b, \\ u(a) &= \alpha, & u(b) = \beta. \end{aligned}$$

# Boundary Value Problems: Finite Difference Method

*The error analysis in full generality is too complicated to be discussed in this course. We will, therefore, look at simpler situation where the right hand side is independent of  $u$  and  $u'$ , that is,  $f(t, u, u') = f(t)$ .*

*The model BVP, therefore, reads*

$$\begin{aligned} u'' &= f(t), & a < t < b, \\ u(a) &= \alpha, & u(b) = \beta. \end{aligned}$$

*We define*

$$D_h^2 v(t) = \frac{v(t+h) - 2v(t) + v(t-h)}{h^2}.$$

*Then, the local (truncation) error*

$$\ell(t_i, u) = D_h^2 u(t_i) - f(t_i)$$

# Boundary Value Problems: Finite Difference Method

*The error analysis in full generality is too complicated to be discussed in this course. We will, therefore, look at simpler situation where the right hand side is independent of  $u$  and  $u'$ , that is,  $f(t, u, u') = f(t)$ .*

*The model BVP, therefore, reads*

$$\begin{aligned} u'' &= f(t), & a < t < b, \\ u(a) &= \alpha, & u(b) = \beta. \end{aligned}$$

*We define*

$$D_h^2 v(t) = \frac{v(t+h) - 2v(t) + v(t-h)}{h^2}.$$

*Then, the local (truncation) error*

$$\ell(t_i, u) = D_h^2 u(t_i) - f(t_i) = \frac{u(t_i+h) - 2u(t_i) + u(t_i-h)}{h^2} - u''(t_i)$$

# Boundary Value Problems: Finite Difference Method

The error analysis in full generality is too complicated to be discussed in this course. We will, therefore, look at simpler situation where the right hand side is independent of  $u$  and  $u'$ , that is,  $f(t, u, u') = f(t)$ .

The model BVP, therefore, reads

$$\begin{aligned} u'' &= f(t), & a < t < b, \\ u(a) &= \alpha, & u(b) = \beta. \end{aligned}$$

We define

$$D_h^2 v(t) = \frac{v(t+h) - 2v(t) + v(t-h)}{h^2}.$$

Then, the local (truncation) error

$$\ell(t_i, u) = D_h^2 u(t_i) - f(t_i) = \frac{u(t_i+h) - 2u(t_i) + u(t_i-h)}{h^2} - u''(t_i)$$

Now, if  $u \in C^4([a, b])$ , then

$$u(t_i+h) = u(t_i) + hu'(t_i) + \frac{h^2}{2}u''(t_i) + \frac{h^3}{6}u'''(t_i) + \frac{h^4}{24}u^{(4)}(\xi_1), \quad \xi_1 \in (t_i, t_i+h),$$

$$u(t_i-h) = u(t_i) - hu'(t_i) + \frac{h^2}{2}u''(t_i) - \frac{h^3}{6}u'''(t_i) + \frac{h^4}{24}u^{(4)}(\xi_2), \quad \xi_2 \in (t_i-h, t_i).$$

# Boundary Value Problems: Finite Difference Method

The error analysis in full generality is too complicated to be discussed in this course. We will, therefore, look at simpler situation where the right hand side is independent of  $u$  and  $u'$ , that is,  $f(t, u, u') = f(t)$ .

The model BVP, therefore, reads

$$\begin{aligned} u'' &= f(t), & a < t < b, \\ u(a) &= \alpha, & u(b) = \beta. \end{aligned}$$

We define

$$D_h^2 v(t) = \frac{v(t+h) - 2v(t) + v(t-h)}{h^2}.$$

Then, the local (truncation) error

$$\ell(t_i, u) = D_h^2 u(t_i) - f(t_i) = \frac{u(t_i+h) - 2u(t_i) + u(t_i-h)}{h^2} - u''(t_i)$$

Now, if  $u \in C^4([a, b])$ , then

$$u(t_i+h) = u(t_i) + hu'(t_i) + \frac{h^2}{2}u''(t_i) + \frac{h^3}{6}u'''(t_i) + \frac{h^4}{24}u^{(4)}(\xi_1), \quad \xi_1 \in (t_i, t_i+h),$$

$$u(t_i-h) = u(t_i) - hu'(t_i) + \frac{h^2}{2}u''(t_i) - \frac{h^3}{6}u'''(t_i) + \frac{h^4}{24}u^{(4)}(\xi_2), \quad \xi_2 \in (t_i-h, t_i).$$

Thus,

$$\ell(t_i, u) = \frac{h^2}{24} \left( u^{(4)}(\xi_1) + u^{(4)}(\xi_2) \right)$$

# Boundary Value Problems: Finite Difference Method

The error analysis in full generality is too complicated to be discussed in this course. We will, therefore, look at simpler situation where the right hand side is independent of  $u$  and  $u'$ , that is,  $f(t, u, u') = f(t)$ .

The model BVP, therefore, reads

$$\begin{aligned} u'' &= f(t), & a < t < b, \\ u(a) &= \alpha, & u(b) = \beta. \end{aligned}$$

We define

$$D_h^2 v(t) = \frac{v(t+h) - 2v(t) + v(t-h)}{h^2}.$$

Then, the local (truncation) error

$$\ell(t_i, u) = D_h^2 u(t_i) - f(t_i) = \frac{u(t_i+h) - 2u(t_i) + u(t_i-h)}{h^2} - u''(t_i)$$

Now, if  $u \in C^4([a, b])$ , then

$$u(t_i+h) = u(t_i) + hu'(t_i) + \frac{h^2}{2}u''(t_i) + \frac{h^3}{6}u'''(t_i) + \frac{h^4}{24}u^{(4)}(\xi_1), \quad \xi_1 \in (t_i, t_i+h),$$

$$u(t_i-h) = u(t_i) - hu'(t_i) + \frac{h^2}{2}u''(t_i) - \frac{h^3}{6}u'''(t_i) + \frac{h^4}{24}u^{(4)}(\xi_2), \quad \xi_2 \in (t_i-h, t_i).$$

Thus,

$$\ell(t_i, u) = \frac{h^2}{24} \left( u^{(4)}(\xi_1) + u^{(4)}(\xi_2) \right) = \frac{h^2}{12} u^{(4)}(\xi), \quad \xi \in (t_i-h, t_i+h).$$

# Boundary Value Problems: Finite Difference Method

## Theorem

If  $v \in C^2([a, b])$ , then

$$\lim_{h \rightarrow 0} \|D_h^2 v - v''\|_{\infty, h} = 0.$$

If  $v \in C^4([a, b])$ , then

$$\|D_h^2 v - v''\|_{\infty, h} \leq \frac{h^2}{12} \|v^{(4)}\|_{\infty, [a, b]}.$$



# Boundary Value Problems: Finite Difference Method

## Theorem

If  $v \in C^2([a, b])$ , then

$$\lim_{h \rightarrow 0} \|D_h^2 v - v''\|_{\infty, h} = 0.$$

If  $v \in C^4([a, b])$ , then

$$\|D_h^2 v - v''\|_{\infty, h} \leq \frac{h^2}{12} \|v^{(4)}\|_{\infty, [a, b]}.$$

## Proof:

We have already seen the proof for the second part. For the first part, we use

$$v(t_i + h) = v(t_i) + hv'(t_i) + \frac{h^2}{2} v''(\xi_1), \quad \xi_1 \in (t_i, t_i + h),$$

$$v(t_i - h) = v(t_i) - hv'(t_i) + \frac{h^2}{2} v''(\xi_2), \quad \xi_2 \in (t_i - h, t_i),$$

yielding

$$D_h^2 v(t_i) - v''(t_i) = \frac{v''(\xi_1) + v''(\xi_2)}{2} - v''(t_i) = v''(\xi) - v''(t_i), \quad \xi \in (t_i - h, t_i + h).$$

The result follows!



# Boundary Value Problems: Finite Difference Method

## *Theorem (Discrete Maximum Principle)*

Let  $v$  be a function on  $[a, b]$  satisfying  $D_h^2 v \geq 0$  on  $t_i, i = 1, \dots, n$ . Then  $\max_{1 \leq i \leq n} v(t_i) \leq \max\{v(t_0), v(t_{n+1})\}$ . Equality holds if and only if  $v$  is constant.

# Boundary Value Problems: Finite Difference Method

## Theorem (Discrete Maximum Principle)

Let  $v$  be a function on  $[a, b]$  satisfying  $D_h^2 v \geq 0$  on  $t_i, i = 1, \dots, n$ . Then  $\max_{1 \leq i \leq n} v(t_i) \leq \max\{v(t_0), v(t_{n+1})\}$ . Equality holds if and only if  $v$  is constant.

### Proof:

Suppose  $\max_{1 \leq i \leq n} v(t_i) \geq \max\{v(t_0), v(t_{n+1})\}$  and the max is achieved at  $t_{i_0}$ .

# Boundary Value Problems: Finite Difference Method

## Theorem (Discrete Maximum Principle)

Let  $v$  be a function on  $[a, b]$  satisfying  $D_h^2 v \geq 0$  on  $t_i, i = 1, \dots, n$ . Then  $\max_{1 \leq i \leq n} v(t_i) \leq \max\{v(t_0), v(t_{n+1})\}$ . Equality holds if and only if  $v$  is constant.

### Proof:

Suppose  $\max_{1 \leq i \leq n} v(t_i) \geq \max\{v(t_0), v(t_{n+1})\}$  and the max is achieved at  $t_{i_0}$ . Then,

$$2v(t_{i_0}) = v(t_{i_0+1}) + v(t_{i_0-1}) - h^2 D_h^2 v(t_{i_0})$$



# Boundary Value Problems: Finite Difference Method

## Theorem (Discrete Maximum Principle)

Let  $v$  be a function on  $[a, b]$  satisfying  $D_h^2 v \geq 0$  on  $t_i, i = 1, \dots, n$ . Then  $\max_{1 \leq i \leq n} v(t_i) \leq \max\{v(t_0), v(t_{n+1})\}$ . Equality holds if and only if  $v$  is constant.

### Proof:

Suppose  $\max_{1 \leq i \leq n} v(t_i) \geq \max\{v(t_0), v(t_{n+1})\}$  and the max is achieved at  $t_{i_0}$ . Then,

$$2v(t_{i_0}) = v(t_{i_0+1}) + v(t_{i_0-1}) - h^2 D_h^2 v(t_{i_0}) \leq v(t_{i_0+1}) + v(t_{i_0-1})$$

# Boundary Value Problems: Finite Difference Method

## Theorem (Discrete Maximum Principle)

Let  $v$  be a function on  $[a, b]$  satisfying  $D_h^2 v \geq 0$  on  $t_i, i = 1, \dots, n$ . Then  $\max_{1 \leq i \leq n} v(t_i) \leq \max\{v(t_0), v(t_{n+1})\}$ . Equality holds if and only if  $v$  is constant.

### Proof:

Suppose  $\max_{1 \leq i \leq n} v(t_i) \geq \max\{v(t_0), v(t_{n+1})\}$  and the max is achieved at  $t_{i_0}$ . Then,

$$2v(t_{i_0}) = v(t_{i_0+1}) + v(t_{i_0-1}) - h^2 D_h^2 v(t_{i_0}) \leq v(t_{i_0+1}) + v(t_{i_0-1}) \leq 2v(t_{i_0})$$

since  $v(t_i) \leq v(t_{i_0})$ .

# Boundary Value Problems: Finite Difference Method

## Theorem (Discrete Maximum Principle)

Let  $v$  be a function on  $[a, b]$  satisfying  $D_h^2 v \geq 0$  on  $t_i, i = 1, \dots, n$ . Then  $\max_{1 \leq i \leq n} v(t_i) \leq \max\{v(t_0), v(t_{n+1})\}$ . Equality holds if and only if  $v$  is constant.

### Proof:

Suppose  $\max_{1 \leq i \leq n} v(t_i) \geq \max\{v(t_0), v(t_{n+1})\}$  and the max is achieved at  $t_{i_0}$ . Then,

$$2v(t_{i_0}) = v(t_{i_0+1}) + v(t_{i_0-1}) - h^2 D_h^2 v(t_{i_0}) \leq v(t_{i_0+1}) + v(t_{i_0-1}) \leq 2v(t_{i_0})$$

since  $v(t_i) \leq v(t_{i_0})$ . Thus, equality holds throughout and  $v$  achieves its maximum at  $t_{i_0-1}$  and  $t_{i_0+1}$  as well.

# Boundary Value Problems: Finite Difference Method

## Theorem (Discrete Maximum Principle)

Let  $v$  be a function on  $[a, b]$  satisfying  $D_h^2 v \geq 0$  on  $t_i, i = 1, \dots, n$ . Then  $\max_{1 \leq i \leq n} v(t_i) \leq \max\{v(t_0), v(t_{n+1})\}$ . Equality holds if and only if  $v$  is constant.

### Proof:

Suppose  $\max_{1 \leq i \leq n} v(t_i) \geq \max\{v(t_0), v(t_{n+1})\}$  and the max is achieved at  $t_{i_0}$ . Then,

$$2v(t_{i_0}) = v(t_{i_0+1}) + v(t_{i_0-1}) - h^2 D_h^2 v(t_{i_0}) \leq v(t_{i_0+1}) + v(t_{i_0-1}) \leq 2v(t_{i_0})$$

since  $v(t_i) \leq v(t_{i_0})$ . Thus, equality holds throughout and  $v$  achieves its maximum at  $t_{i_0-1}$  and  $t_{i_0+1}$  as well. Applying the same arguments again to  $t_{i_0-1}$  and  $t_{i_0+1}$  (if in the interior), and then to their neighbors, etc., we conclude that  $v$  is constant.



# Boundary Value Problems: Finite Difference Method

## Theorem (Discrete Maximum Principle)

Let  $v$  be a function on  $[a, b]$  satisfying  $D_h^2 v \geq 0$  on  $t_i, i = 1, \dots, n$ . Then  $\max_{1 \leq i \leq n} v(t_i) \leq \max\{v(t_0), v(t_{n+1})\}$ . Equality holds if and only if  $v$  is constant.

### Proof:

Suppose  $\max_{1 \leq i \leq n} v(t_i) \geq \max\{v(t_0), v(t_{n+1})\}$  and the max is achieved at  $t_{i_0}$ . Then,

$$2v(t_{i_0}) = v(t_{i_0+1}) + v(t_{i_0-1}) - h^2 D_h^2 v(t_{i_0}) \leq v(t_{i_0+1}) + v(t_{i_0-1}) \leq 2v(t_{i_0})$$

since  $v(t_i) \leq v(t_{i_0})$ . Thus, equality holds throughout and  $v$  achieves its maximum at  $t_{i_0-1}$  and  $t_{i_0+1}$  as well. Applying the same arguments again to  $t_{i_0-1}$  and  $t_{i_0+1}$  (if in the interior), and then to their neighbors, etc., we conclude that  $v$  is constant.

### Remark

The analogous discrete minimum principle, obtained by reversing the inequalities and replacing max by min holds.

# Boundary Value Problems: Finite Difference Method

## Theorem (Discrete Maximum Principle)

Let  $v$  be a function on  $[a, b]$  satisfying  $D_h^2 v \geq 0$  on  $t_i, i = 1, \dots, n$ . Then  $\max_{1 \leq i \leq n} v(t_i) \leq \max\{v(t_0), v(t_{n+1})\}$ . Equality holds if and only if  $v$  is constant.

### Proof:

Suppose  $\max_{1 \leq i \leq n} v(t_i) \geq \max\{v(t_0), v(t_{n+1})\}$  and the max is achieved at  $t_{i_0}$ . Then,

$$2v(t_{i_0}) = v(t_{i_0+1}) + v(t_{i_0-1}) - h^2 D_h^2 v(t_{i_0}) \leq v(t_{i_0+1}) + v(t_{i_0-1}) \leq 2v(t_{i_0})$$

since  $v(t_i) \leq v(t_{i_0})$ . Thus, equality holds throughout and  $v$  achieves its maximum at  $t_{i_0-1}$  and  $t_{i_0+1}$  as well. Applying the same arguments again to  $t_{i_0-1}$  and  $t_{i_0+1}$  (if in the interior), and then to their neighbors, etc., we conclude that  $v$  is constant.

### Remark

The analogous discrete minimum principle, obtained by reversing the inequalities and replacing max by min holds.

### Theorem

There is a unique solution to the discrete BVP

$$\begin{aligned} D_h^2 u_h(t_i) &= f(t_i), \quad t_i, i = 1, \dots, n, \\ u_h(a) &= \alpha, \quad u_h(b) = \beta. \end{aligned}$$