

Numerical Analysis & Scientific Computing II

Lesson 3

Boundary Value Problems for ODEs

3.1 Well-posedness

3.2 Shooting Method



Akash Anand
MATH, IIT KANPUR

Boundary Value Problems: Shooting Method

Recall the following discussion in the context of solvability of two-point BVP

$$y' = f(t, y), \quad a < t < b,$$

with boundary conditions

$$g(y(a), y(b)) = 0.$$

We noted that if $y(t; x)$ denotes the solution to the IVP $y' = f(t, y)$, $y(a) = x$, $x \in \mathbb{R}^n$, then this solution is a solution to the BVP if

$$g(x, y(b; x)) = 0.$$

Thus, one approach to solve the BVP therefore is to find an x that solves the system of nonlinear algebraic equations $h(x) = 0$, where $h(x) = g(x, y(b; x))$.

Boundary Value Problems: Shooting Method

Recall the following discussion in the context of solvability of two-point BVP

$$y' = f(t, y), \quad a < t < b,$$

with boundary conditions


$$g(y(a), y(b)) = 0.$$

We noted that if $y(t; x)$ denotes the solution to the IVP $y' = f(t, y)$, $y(a) = x$, $x \in \mathbb{R}^n$, then this solution is a solution to the BVP if

$$g(x, y(b; x)) = 0.$$

Thus, one approach to solve the BVP therefore is to find an x that solves the system of nonlinear algebraic equations $h(x) = 0$, where $h(x) = g(x, y(b; x))$.

For example, consider the ODE $u'' = f(t, u, u')$ with boundary conditions $u(a) = \alpha$, $u(b) = \beta$.



Boundary Value Problems: Shooting Method

Recall the following discussion in the contest of solvability of two-point BVP

$$y' = f(t, y), \quad a < t < b,$$

with boundary conditions

$$g(y(a), y(b)) = 0.$$


We noted that if $y(t; x)$ denotes the solution to the IVP $y' = f(t, y)$, $y(a) = x$, $x \in \mathbb{R}^n$, then this solution is a solution to the BVP if

$$g(x, y(b; x)) = 0.$$

Thus, one approach to solve the BVP therefore is to find an x that solves the system of nonlinear algebraic equations $h(x) = 0$, where $h(x) = g(x, y(b; x))$.

For example, consider the ODE $u'' = f(t, u, u')$ with boundary conditions $u(a) = \alpha$, $u(b) = \beta$.

If we knew $u'(a)$, then ...



Boundary Value Problems: Shooting Method

Recall the following discussion in the context of solvability of two-point BVP

$$y' = f(t, y), \quad a < t < b,$$

with boundary conditions

$$g(y(a), y(b)) = 0.$$

We noted that if $y(t; x)$ denotes the solution to the IVP $y' = f(t, y)$, $y(a) = x$, $x \in \mathbb{R}^n$, then this solution is a solution to the BVP if


$$g(x, y(b; x)) = 0.$$

Thus, one approach to solve the BVP therefore is to find an x that solves the system of nonlinear algebraic equations $h(x) = 0$, where $h(x) = g(x, y(b; x))$.

For example, consider the ODE $u'' = f(t, u, u')$ with boundary conditions $u(a) = \alpha$, $u(b) = \beta$.

If we knew $u'(a)$, then we could have solved the IVP $u'' = f(t, u, u')$, $u(a) = \alpha$, $u'(a) = \alpha'$ (say), to obtain the solution that passes through (b, β) , that is, $y_1(t; \alpha, \alpha') = u(t)$, $y_2(t; \alpha, \alpha') = u'(t)$ satisfies $h([\alpha, \alpha']^T) = 0$, where

$$h(s) = \begin{bmatrix} y_1(a; \alpha, s) - \alpha \\ y_1(b; \alpha, s) - \beta \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$



Boundary Value Problems: Shooting Method

Recall the following discussion in the contest of solvability of two-point BVP

$$y' = f(t, y), \quad a < t < b,$$

with boundary conditions

$$g(y(a), y(b)) = 0.$$

We noted that if $y(t; x)$ denotes the solution to the IVP $y' = f(t, y)$, $y(a) = x$, $x \in \mathbb{R}^n$, then this solution is a solution to the BVP if


$$g(x, y(b; x)) = 0.$$

Thus, one approach to solve the BVP therefore is to find an x that solves the system of nonlinear algebraic equations $h(x) = 0$, where $h(x) = g(x, y(b; x))$.

For example, consider the ODE $u'' = f(t, u, u')$ with boundary conditions $u(a) = \alpha$, $u(b) = \beta$.

If we knew $u'(a)$, then we could have solved the IVP $u'' = f(t, u, u')$, $u(a) = \alpha$, $u'(a) = \alpha'$ (say), to obtain the solution that passes through (b, β) , that is, $y_1(t; \alpha, \alpha') = u(t)$, $y_2(t; \alpha, \alpha') = u'(t)$ satisfies $h([\alpha, \alpha']^T) = 0$, where

$$h(s) = \begin{bmatrix} y_1(a; \alpha, s) - \alpha \\ y_1(b; \alpha, s) - \beta \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}, \text{ that is, } h_2(\alpha, s) := y_1(b; \alpha, s) - \beta = 0$$



Boundary Value Problems: Shooting Method

Recall the following discussion in the contest of solvability of two-point BVP

$$y' = f(t, y), \quad a < t < b,$$

with boundary conditions

$$g(y(a), y(b)) = 0.$$

We noted that if $y(t; x)$ denotes the solution to the IVP $y' = f(t, y)$, $y(a) = x$, $x \in \mathbb{R}^n$, then this solution is a solution to the BVP if


$$g(x, y(b; x)) = 0.$$

Thus, one approach to solve the BVP therefore is to find an x that solves the system of nonlinear algebraic equations $h(x) = 0$, where $h(x) = g(x, y(b; x))$.

For example, consider the ODE $u'' = f(t, u, u')$ with boundary conditions $u(a) = \alpha$, $u(b) = \beta$.

If we knew $u'(a)$, then we could have solved the IVP $u'' = f(t, u, u')$, $u(a) = \alpha$, $u'(a) = \alpha'$ (say), to obtain the solution that passes through (b, β) , that is, $y_1(t; \alpha, \alpha') = u(t)$, $y_2(t; \alpha, \alpha') = u'(t)$ satisfies $h([\alpha, \alpha']^T) = 0$, where

$$h(s) = \begin{bmatrix} y_1(a; \alpha, s) - \alpha \\ y_1(b; \alpha, s) - \beta \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}, \text{ that is, } h_2(\alpha, s) := y_1(b; \alpha, s) - \beta = 0$$



Boundary Value Problems: Shooting Method

Recall the following discussion in the context of solvability of two-point BVP

$$y' = f(t, y), \quad a < t < b,$$

with boundary conditions

$$g(y(a), y(b)) = 0.$$

We noted that if $y(t; x)$ denotes the solution to the IVP $y' = f(t, y)$, $y(a) = x$, $x \in \mathbb{R}^n$, then this solution is a solution to the BVP if


$$g(x, y(b; x)) = 0.$$

Thus, one approach to solve the BVP therefore is to find an x that solves the system of nonlinear algebraic equations $h(x) = 0$, where $h(x) = g(x, y(b; x))$.

For example, consider the ODE $u'' = f(t, u, u')$ with boundary conditions $u(a) = \alpha$, $u(b) = \beta$.

If we knew $u'(a)$, then we could have solved the IVP $u'' = f(t, u, u')$, $u(a) = \alpha$, $u'(a) = \alpha'$ (say), to obtain the solution that passes through (b, β) , that is, $y_1(t; \alpha, \alpha') = u(t)$, $y_2(t; \alpha, \alpha') = u'(t)$ satisfies $h([\alpha, \alpha']^T) = 0$, where

$$h(s) = \begin{bmatrix} y_1(a; \alpha, s) - \alpha \\ y_1(b; \alpha, s) - \beta \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}, \text{ that is, } h_2(\alpha, s) := y_1(b; \alpha, s) - \beta = 0$$



Boundary Value Problems: Shooting Method

Recall the following discussion in the contest of solvability of two-point BVP

$$y' = f(t, y), \quad a < t < b,$$

with boundary conditions

$$g(y(a), y(b)) = 0.$$

We noted that if $y(t; x)$ denotes the solution to the IVP $y' = f(t, y)$, $y(a) = x$, $x \in \mathbb{R}^n$, then this solution is a solution to the BVP if


$$g(x, y(b; x)) = 0.$$

Thus, one approach to solve the BVP therefore is to find an x that solves the system of nonlinear algebraic equations $h(x) = 0$, where $h(x) = g(x, y(b; x))$.

For example, consider the ODE $u'' = f(t, u, u')$ with boundary conditions $u(a) = \alpha$, $u(b) = \beta$.

If we knew $u'(a)$, then we could have solved the IVP $u'' = f(t, u, u')$, $u(a) = \alpha$, $u'(a) = \alpha'$ (say), to obtain the solution that passes through (b, β) , that is, $y_1(t; \alpha, \alpha') = u(t)$, $y_2(t; \alpha, \alpha') = u'(t)$ satisfies $h([\alpha, \alpha']^T) = 0$, where

$$h(s) = \begin{bmatrix} y_1(a; \alpha, s) - \alpha \\ y_1(b; \alpha, s) - \beta \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}, \text{ that is, } h_2(\alpha, s) := y_1(b; \alpha, s) - \beta = 0$$



Boundary Value Problems: Shooting Method

Recall the following discussion in the contest of solvability of two-point BVP

$$y' = f(t, y), \quad a < t < b,$$

with boundary conditions

$$g(y(a), y(b)) = 0.$$

We noted that if $y(t; x)$ denotes the solution to the IVP $y' = f(t, y)$, $y(a) = x$, $x \in \mathbb{R}^n$, then this solution is a solution to the BVP if


$$g(x, y(b; x)) = 0.$$

Thus, one approach to solve the BVP therefore is to find an x that solves the system of nonlinear algebraic equations $h(x) = 0$, where $h(x) = g(x, y(b; x))$.

For example, consider the ODE $u'' = f(t, u, u')$ with boundary conditions $u(a) = \alpha$, $u(b) = \beta$.

If we knew $u'(a)$, then we could have solved the IVP $u'' = f(t, u, u')$, $u(a) = \alpha$, $u'(a) = \alpha'$ (say), to obtain the solution that passes through (b, β) , that is, $y_1(t; \alpha, \alpha') = u(t)$, $y_2(t; \alpha, \alpha') = u'(t)$ satisfies $h([\alpha, \alpha']^T) = 0$, where

$$h(s) = \begin{bmatrix} y_1(a; \alpha, s) - \alpha \\ y_1(b; \alpha, s) - \beta \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}, \text{ that is, } h_2(\alpha, s) := y_1(b; \alpha, s) - \beta = 0$$



Boundary Value Problems: Shooting Method

Recall the following discussion in the context of solvability of two-point BVP

$$y' = f(t, y), \quad a < t < b,$$

with boundary conditions

$$g(y(a), y(b)) = 0.$$

We noted that if $y(t; x)$ denotes the solution to the IVP $y' = f(t, y)$, $y(a) = x$, $x \in \mathbb{R}^n$, then this solution is a solution to the BVP if


$$g(x, y(b; x)) = 0.$$

Thus, one approach to solve the BVP therefore is to find an x that solves the system of nonlinear algebraic equations $h(x) = 0$, where $h(x) = g(x, y(b; x))$.

For example, consider the ODE $u'' = f(t, u, u')$ with boundary conditions $u(a) = \alpha$, $u(b) = \beta$.

If we knew $u'(a)$, then we could have solved the IVP $u'' = f(t, u, u')$, $u(a) = \alpha$, $u'(a) = \alpha'$ (say), to obtain the solution that passes through (b, β) , that is, $y_1(t; \alpha, \alpha') = u(t)$, $y_2(t; \alpha, \alpha') = u'(t)$ satisfies $h([\alpha, \alpha']^T) = 0$, where

$$h(s) = \begin{bmatrix} y_1(a; \alpha, s) - \alpha \\ y_1(b; \alpha, s) - \beta \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}, \text{ that is, } h_2(\alpha, s) := y_1(b; \alpha, s) - \beta = 0$$



Boundary Value Problems: Shooting Method

Recall the following discussion in the contest of solvability of two-point BVP

$$y' = f(t, y), \quad a < t < b,$$

with boundary conditions

$$g(y(a), y(b)) = 0.$$

We noted that if $y(t; x)$ denotes the solution to the IVP $y' = f(t, y)$, $y(a) = x$, $x \in \mathbb{R}^n$, then this solution is a solution to the BVP if


$$g(x, y(b; x)) = 0.$$

Thus, one approach to solve the BVP therefore is to find an x that solves the system of nonlinear algebraic equations $h(x) = 0$, where $h(x) = g(x, y(b; x))$.

For example, consider the ODE $u'' = f(t, u, u')$ with boundary conditions $u(a) = \alpha$, $u(b) = \beta$.

If we knew $u'(a)$, then we could have solved the IVP $u'' = f(t, u, u')$, $u(a) = \alpha$, $u'(a) = \alpha'$ (say), to obtain the solution that passes through (b, β) , that is, $y_1(t; \alpha, \alpha') = u(t)$, $y_2(t; \alpha, \alpha') = u'(t)$ satisfies $h([\alpha, \alpha']^T) = 0$, where

$$h(s) = \begin{bmatrix} y_1(a; \alpha, s) - \alpha \\ y_1(b; \alpha, s) - \beta \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}, \text{ that is, } h_2(\alpha, s) := y_1(b; \alpha, s) - \beta = 0$$



Boundary Value Problems: Shooting Method

Recall the following discussion in the contest of solvability of two-point BVP

$$y' = f(t, y), \quad a < t < b,$$

with boundary conditions

$$g(y(a), y(b)) = 0.$$

We noted that if $y(t; x)$ denotes the solution to the IVP $y' = f(t, y)$, $y(a) = x$, $x \in \mathbb{R}^n$, then this solution is a solution to the BVP if

$$g(x, y(b; x)) = 0.$$

Thus, one approach to solve the BVP therefore is to find an x that solves the system of nonlinear algebraic equations $h(x) = 0$, where $h(x) = g(x, y(b; x))$.

For example, consider the ODE $u'' = f(t, u, u')$ with boundary conditions $u(a) = \alpha$, $u(b) = \beta$.

If we knew $u'(a)$, then we could have solved the IVP $u'' = f(t, u, u')$, $u(a) = \alpha$, $u'(a) = \alpha'$ (say), to obtain the solution that passes through (b, β) , that is, $y_1(t; \alpha, \alpha') = u(t)$, $y_2(t; \alpha, \alpha') = u'(t)$ satisfies $h([\alpha, \alpha']^T) = 0$, where

$$h(s) = \begin{bmatrix} y_1(a; \alpha, s) - \alpha \\ y_1(b; \alpha, s) - \beta \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}, \text{ that is, } h_2(\alpha, s) := y_1(b; \alpha, s) - \beta = 0$$



Boundary Value Problems: Shooting Method

In particular, consider the two-point BVP

$$\begin{aligned} u'' &= f(t, u, u'), & a < t < b, \\ a_0 u(a) - a_1 u'(a) &= \alpha, & b_0 u(b) + b_1 u'(b) = \beta, \end{aligned}$$

where the function f is assumed to satisfy the following Lipschitz conditions:

$$\begin{aligned} |f(t, u_1, v) - f(t, u_2, v)| &\leq K|u_1 - u_2|, \\ |f(t, u, v_1) - f(t, u, v_2)| &\leq K|v_1 - v_2|, \end{aligned}$$

for all points $(t, u_i, v), (t, u, v_j) \in R := [a, b] \times \mathbb{R} \times \mathbb{R}$. In addition, assume that on R , f satisfies

$$f_u(t, u, v) = \frac{\partial f(t, u, v)}{\partial u} > 0, \quad |f_v(t, u, v)| = \left| \frac{\partial f(t, u, v)}{\partial v} \right| \leq M,$$

for some $M > 0$. For the boundary conditions, assume

$$a_0 a_1 \geq 0, \quad b_0 b_1 \geq 0, \quad |a_0| + |a_1| \neq 0, \quad |b_0| + |b_1| \neq 0, \quad |a_0| + |b_0| \neq 0.$$

Boundary Value Problems: Shooting Method

In particular, consider the two-point BVP

$$\begin{aligned} u'' &= f(t, u, u'), & a < t < b, \\ a_0 u(a) - a_1 u'(a) &= \alpha, & b_0 u(b) + b_1 u'(b) = \beta, \end{aligned}$$

where the function f is assumed to satisfy the following Lipschitz conditions:

$$\begin{aligned} |f(t, u_1, v) - f(t, u_2, v)| &\leq K|u_1 - u_2|, \\ |f(t, u, v_1) - f(t, u, v_2)| &\leq K|v_1 - v_2|, \end{aligned}$$

for all points $(t, u_i, v), (t, u, v_j) \in R := [a, b] \times \mathbb{R} \times \mathbb{R}$. In addition, assume that on R , f satisfies

$$f_u(t, u, v) = \frac{\partial f(t, u, v)}{\partial u} > 0, \quad |f_v(t, u, v)| = \left| \frac{\partial f(t, u, v)}{\partial v} \right| \leq M,$$

for some $M > 0$. For the boundary conditions, assume

$$a_0 a_1 \geq 0, \quad b_0 b_1 \geq 0, \quad |a_0| + |a_1| \neq 0, \quad |b_0| + |b_1| \neq 0, \quad |a_0| + |b_0| \neq 0.$$

Theorem

The BVP, under these assumptions, has a unique solution.

Boundary Value Problems: Shooting Method

In particular, consider the two-point BVP

$$\begin{aligned} u'' &= f(t, u, u'), & a < t < b, \\ a_0 u(a) - a_1 u'(a) &= \alpha, & b_0 u(b) + b_1 u'(b) = \beta, \end{aligned}$$

where the function f is assumed to satisfy the following Lipschitz conditions:

$$\begin{aligned} |f(t, u_1, v) - f(t, u_2, v)| &\leq K|u_1 - u_2|, \\ |f(t, u, v_1) - f(t, u, v_2)| &\leq K|v_1 - v_2|, \end{aligned}$$

for all points $(t, u_i, v), (t, u, v_j) \in R := [a, b] \times \mathbb{R} \times \mathbb{R}$. In addition, assume that on R , f satisfies

$$f_u(t, u, v) = \frac{\partial f(t, u, v)}{\partial u} > 0, \quad |f_v(t, u, v)| = \left| \frac{\partial f(t, u, v)}{\partial v} \right| \leq M,$$

for some $M > 0$. For the boundary conditions, assume

$$a_0 a_1 \geq 0, \quad b_0 b_1 \geq 0, \quad |a_0| + |a_1| \neq 0, \quad |b_0| + |b_1| \neq 0, \quad |a_0| + |b_0| \neq 0.$$

Theorem

The BVP, under these assumptions, has a unique solution.

How do we solve the BVP using the shooting method?

Boundary Value Problems: Shooting Method



Consider the two-point BVP

$$\begin{aligned} u'' &= f(t, u, u'), & a < t < b, \\ a_0 u(a) - a_1 u'(a) &= \alpha, & b_0 u(b) + b_1 u'(b) = \beta. \end{aligned}$$

Boundary Value Problems: Shooting Method

Consider the two-point BVP

$$\begin{aligned} u'' &= f(t, u, u'), & a < t < b, \\ a_0 u(a) - a_1 u'(a) &= \alpha, & b_0 u(b) + b_1 u'(b) = \beta. \end{aligned}$$

Let $y(t; s)$ be the solution to the IVP

$$\begin{aligned} y'' &= f(t, y, y'), & a < t < b, \\ y(a) &= a_1 s - c_1 \alpha, & y'(a) = a_0 s - c_0 \alpha \end{aligned}$$

depending on the parameter s , where c_0, c_1 are arbitrary constants satisfying $a_1 c_0 - a_0 c_1 = 1$. Note that $a_0 y(a; s) - a_1 y'(a; s) = \alpha$.

Boundary Value Problems: Shooting Method

Consider the two-point BVP

$$\begin{aligned} u'' &= f(t, u, u'), & a < t < b, \\ a_0 u(a) - a_1 u'(a) &= \alpha, & b_0 u(b) + b_1 u'(b) &= \beta. \end{aligned}$$

Let $y(t; s)$ be the solution to the IVP

$$\begin{aligned} y'' &= f(t, y, y'), & a < t < b, \\ y(a) &= a_1 s - c_1 \alpha, & y'(a) &= a_0 s - c_0 \alpha \end{aligned}$$

depending on the parameter s , where c_0, c_1 are arbitrary constants satisfying $a_1 c_0 - a_0 c_1 = 1$. Note that $a_0 y(a; s) - a_1 y'(a; s) = \alpha$.

For $y(t; s)$ to be the solution to the BVP, it must satisfy

$$h(s) := b_0 y(b; s) + b_1 y'(b; s) - \beta = 0.$$

Boundary Value Problems: Shooting Method

Consider the two-point BVP

$$\begin{aligned} u'' &= f(t, u, u'), & a < t < b, \\ a_0 u(a) - a_1 u'(a) &= \alpha, & b_0 u(b) + b_1 u'(b) = \beta. \end{aligned}$$

Let $y(t; s)$ be the solution to the IVP

$$\begin{aligned} y'' &= f(t, y, y'), & a < t < b, \\ y(a) &= a_1 s - c_1 \alpha, & y'(a) = a_0 s - c_0 \alpha \end{aligned}$$

depending on the parameter s , where c_0, c_1 are arbitrary constants satisfying $a_1 c_0 - a_0 c_1 = 1$. Note that $a_0 y(a; s) - a_1 y'(a; s) = \alpha$.

For $y(t; s)$ to be the solution to the BVP, it must satisfy

$$h(s) := b_0 y(b; s) + b_1 y'(b; s) - \beta = 0.$$

For solution of this equation using the Newton's method, we have

$$s_{m+1} = s_m - \frac{h(s_m)}{h'(s_m)}, \quad m = 0, 1, 2, \dots$$

Boundary Value Problems: Shooting Method

Consider the two-point BVP

$$\begin{aligned} u'' &= f(t, u, u'), & a < t < b, \\ a_0 u(a) - a_1 u'(a) &= \alpha, & b_0 u(b) + b_1 u'(b) &= \beta. \end{aligned}$$

Let $y(t; s)$ be the solution to the IVP

$$\begin{aligned} y'' &= f(t, y, y'), & a < t < b, \\ y(a) &= a_1 s - c_1 \alpha, & y'(a) &= a_0 s - c_0 \alpha \end{aligned}$$

depending on the parameter s , where c_0, c_1 are arbitrary constants satisfying $a_1 c_0 - a_0 c_1 = 1$. Note that $a_0 y(a; s) - a_1 y'(a; s) = \alpha$.

For $y(t; s)$ to be the solution to the BVP, it must satisfy

$$h(s) := b_0 y(b; s) + b_1 y'(b; s) - \beta = 0.$$

For solution of this equation using the Newton's method, we have

$$s_{m+1} = s_m - \frac{h(s_m)}{h'(s_m)}, \quad m = 0, 1, 2, \dots$$

We need to find the derivative h' . Note that

$$h'(s) = b_0 z_s(b) + b_1 z'_s(b)$$

where

$$z_s(t) = \frac{\partial y(t; s)}{\partial s}.$$

Boundary Value Problems: Shooting Method

Consider the two-point BVP

$$\begin{aligned} u'' &= f(t, u, u'), & a < t < b, \\ a_0 u(a) - a_1 u'(a) &= \alpha, & b_0 u(b) + b_1 u'(b) = \beta. \end{aligned}$$

Let $y(t; s)$ be the solution to the IVP

$$\begin{aligned} y'' &= f(t, y, y'), & a < t < b, \\ y(a) &= a_1 s - c_1 \alpha, & y'(a) = a_0 s - c_0 \alpha \end{aligned}$$

depending on the parameter s , where c_0, c_1 are arbitrary constants satisfying $a_1 c_0 - a_0 c_1 = 1$.

...

We need to find the derivative h' . Note that

$$h'(s) = b_0 z_s(b) + b_1 z'_s(b)$$

where

$$z_s(t) = \frac{\partial y(t; s)}{\partial s}.$$

To find z_s , we differentiate the equation $y''(t; s) = f(t, y(t; s), y'(t; s))$ with respect to s .

Boundary Value Problems: Shooting Method

Consider the two-point BVP

$$\begin{aligned} u'' &= f(t, u, u'), & a < t < b, \\ a_0 u(a) - a_1 u'(a) &= \alpha, & b_0 u(b) + b_1 u'(b) = \beta. \end{aligned}$$

Let $y(t; s)$ be the solution to the IVP

$$\begin{aligned} y'' &= f(t, y, y'), & a < t < b, \\ y(a) &= a_1 s - c_1 \alpha, & y'(a) = a_0 s - c_0 \alpha \end{aligned}$$

depending on the parameter s , where c_0, c_1 are arbitrary constants satisfying $a_1 c_0 - a_0 c_1 = 1$.

...

We need to find the derivative h' . Note that

$$h'(s) = b_0 z_s(b) + b_1 z'_s(b)$$

where

$$z_s(t) = \frac{\partial y(t; s)}{\partial s}.$$

To find z_s , we differentiate the equation $y''(t; s) = f(t, y(t; s), y'(t; s))$ with respect to s . Then, z_s satisfies the IVP

$$z_s''(t) = f_2(t, y(t; s), y'(t; s)) z_s(t) + f_3(t, y(t; s), y'(t; s)) z'_s(t), \quad z_s(a) = a_1, \quad z'_s(a) = a_0.$$

The functions f_2 and f_3 denote the partial derivatives of $f(t, u, v)$ with respect to u and v respectively.