

The Differential of f

by Sophia



WHAT'S COVERED

In this lesson, you will express linear approximations in terms of differentials. Specifically, this lesson will cover:

1. Defining the Differential of f
2. Calculating the Differential of f

1. Defining the Differential of f


From when we first discussed rates of change and the derivative, recall the following quantities:

- Δx = change in x (horizontal change)
- Δy = change in y (vertical change)

When $y = f(x)$, Δy can be replaced with Δf to show that this is the change in function f .

Another goal in linear approximation is to find the change in f for a corresponding change in x .

Consider the image below:



Let Δx = the horizontal change in x -values. This was " $x - a$ " in the linear approximation formula.

Let Δf = the actual change in f when moving from x to $x + \Delta x$. Then, $\Delta f = f(x + \Delta x) - f(x)$.

Now, let A = the approximate change in f along the tangent line, which can be found as follows:

- Slope = $\frac{\text{rise}}{\text{run}} = f'(x) = \frac{A}{\Delta x}$
- Then, solving for A , we get $A = f'(x) \cdot \Delta x$.

Since A is approximating Δf , we can also say that $\Delta f \approx f'(x) \cdot \Delta x$.

This means that the change in f (when moving from x to $x + \Delta x$) can be approximated by multiplying the slope $f'(x)$ by Δx , the change in x .

This leads to the definition of the **differential of f** .



FORMULA

Differential of f

$df = f'(x)dx$ for any choice of x and any real number dx .

When $y = f(x)$, we can also write $dy = f'(x)dx$.



HINT

The differential uses the derivative at an x -value to give the approximate change in f when x changes to $x + \Delta x$.

When approximating the change in y , we use $dx = \Delta x$, but dy is an approximation of Δy .

2. Calculating the Differential of f

➡ EXAMPLE Given $f(x) = 4x^2 + 7x$, find the differential df .

Since $f'(x) = 8x + 7$, the differential is $df = (8x + 7)dx$.

➡ EXAMPLE Given $y = \ln(x^2 + 3)$, find the differential dy .

Since $\frac{dy}{dx} = \frac{1}{x^2 + 3} \cdot 2x = \frac{2x}{x^2 + 3}$, the differential is $dy = \frac{2x}{x^2 + 3} dx$.



TRY IT

Let $f(x) = x^2 \sin(2x)$.

Find the differential df .

+

$$df = (2x \sin(2x) + 2x^2 \cos(2x))dx$$



WATCH

Here is a video in which we find the differential dy of $y = e^{-4x} \cos(7x)$.



SUMMARY

In this lesson, you learned how to **define and calculate the differential of f** , which is an approximation for the change in f when x changes by dx units.

SOURCE: THIS WORK IS ADAPTED FROM CHAPTER 2 OF *CONTEMPORARY CALCULUS* BY DALE HOFFMAN.



FORMULAS TO KNOW

Differential of f

$df = f'(x)dx$ for any choice of x and any real number dx .

When $y = f(x)$, we can also write $dy = f'(x)dx$.