

Differential Geometry I

Lecture notes

Andriy Haydys

November 20, 2021

This is a draft. If you spot a mistake, please e-mail me: [andriy\(DOT\)haydys@ulb.be](mailto:andriy(DOT)haydys@ulb.be).

Contents

1	Introduction	2
2	Smooth manifolds	4
2.1	Basic definitions and examples	4
2.2	Smooth maps	8
2.3	The fundamental theorem of algebra	11
2.4	Tangent spaces	13
2.5	Cut off and bump functions	19
2.6	The differential of a smooth map	20
3	Submanifolds and partitions of unity	23
3.1	Submanifolds	23
3.2	Immersions and embeddings	29
3.3	Partitions of unity	32
4	The tangent bundle and the group of diffeomorphisms	35
4.1	Some elements of linear algebra	35
4.2	The tangent and cotangent bundles	36

Chapter 1

Introduction

A substantial part of mathematics is related to solving equations of various types. Given any equation, we may try to analyze this by studying the following sequence of questions:

1. Does there exist a solution (a root)?
2. If the answer to the previous question is affirmative, how many solutions does the equation have?
3. If there are finitely many solutions, can we find all of them?

For example, the reader learned at school the properties of the quadratic equation $ax^2 + bx + c = 0$. In this case the above questions are easy to settle and the answers are well known to the reader.

Sometimes an equation may have an infinite number of solutions. If there are only countably many roots, the last question from the list above still makes sense. For example, all solutions of the equation $\sin x = 0$ are given by a simple formula: $x_n = \pi n$, $n \in \mathbb{Z}$.

In many cases, however, equations have uncountably many solutions so that asking to find all solutions is not really meaningful. Instead, it turns out to be more interesting to replace Question 3 by the following one:

- 3'. What are the properties of the set of *all solutions*?

Which particular properties we are interested in may depend on the context. The property most relevant to the content of this course is concerned with the local structure of the set of all solutions.

Let us consider an example. The equation

$$x_1^2 + x_2^2 + x_3^2 = 1, \tag{1.1}$$

where $x_1, x_2, x_3 \in \mathbb{R}$, clearly has uncountably many solutions.

Denote $S^2 := \{x = (x_1, x_2, x_3) \in \mathbb{R}^3 \mid x_1^2 + x_2^2 + x_3^2 = 1\}$, that is S^2 is the set of all solutions of (1.1). Of course, S^2 is the sphere of radius 1, however let us pretend for a moment that we do not know this. As a subset of \mathbb{R}^3 , S^2 is a topological space. It turns out that this topological space has a very particular property, which we consider in some detail next.

The familiar stereographic projection from the north pole $N := (0, 0, 1)$ is given by

$$\varphi_N: S^2 \setminus \{N\} \rightarrow \mathbb{R}^2, \quad \varphi_N(x) = \left(\frac{x_1}{1 - x_3}, \frac{x_2}{1 - x_3} \right).$$

This is in fact a homeomorphism with the inverse

$$\varphi_N^{-1}(y) = \frac{1}{1 + y_1^2 + y_2^2} (2y_1, 2y_2, -1 + y_1^2 + y_2^2), \quad y = (y_1, y_2) \in \mathbb{R}^2.$$

We can also define a stereographic projection from the south pole $S := (0, 0, -1)$ by

$$\varphi_S: S^2 \setminus \{S\} \rightarrow \mathbb{R}^2, \quad \varphi_S(x) = \left(\frac{x_1}{1+x_3}, \frac{x_2}{1+x_3} \right),$$

which is also a homeomorphism.

Since any point on the sphere lies either in $S^2 \setminus \{N\}$ or $S^2 \setminus \{S\}$ (or both), any point on the sphere has a neighbourhood, which is homeomorphic to an open subset of \mathbb{R}^n (of course, $n = 2$ in our particular example and the open subset is \mathbb{R}^2 itself). This property leads to the notion of *a manifold*, which will play a central rôle in the course. We will see below, that this property is *not* specific to Equation (1.1). On the contrary, for any smooth map $F: \mathbb{R}^k \rightarrow \mathbb{R}^\ell$ and almost any $c \in \mathbb{R}^\ell$ the set of all solutions to the equation $F(x) = c$ is a manifold. That is, there is a huge pull of examples of manifolds and many objects of particular interest in mathematics turn out to be manifolds.

Coming back to our example, we compute:

$$\varphi_S \circ \varphi_N^{-1}(y) = \left(\frac{y_1}{|y|^2}, \frac{y_2}{|y|^2} \right). \quad (1.2)$$

Hence, $\varphi_S \circ \varphi_N^{-1}$ is *smooth* on an open subset $\mathbb{R}^2 \setminus \{0\}$ and a similar computation yields that this is also true for $\varphi_N \circ \varphi_S^{-1}$. This property can be used to study smooth functions on the sphere directly without reference to the ambient space. More importantly, in more general situations where the ambient Euclidean space may be simply absent, an analogue of this property allows one to apply familiar tools of analysis to functions defined on more sophisticated objects than just subsets of an Euclidean space. In some sense, this constitutes the core of differential geometry.

Summing up, the aim of these notes is to transfer familiar tools of mathematical analysis to a more geometric setting where the underlying domain of a function (map) is not just an open subset of \mathbb{R}^n , but rather a manifold. The benefits of doing so are ubiquitous, but explaining this in some detail requires a bit of work. It is my hope to convey that the notion of a manifold is useful and well worth studying further.

Chapter 2

Smooth manifolds

2.1 Basic definitions and examples

Recall that a topological space M is called Hausdorff, if for any two distinct points $m_1, m_2 \in M$ there are neighbourhoods $U_1 \ni m_1$ and $U_2 \ni m_2$ such that $U_1 \cap U_2 = \emptyset$. If the topology of M admits a countable base, then M is said to be second countable. For example, \mathbb{R}^k is both Hausdorff and second countable.

Definition 2.1. A Hausdorff second countable topological space M is called a *topological manifold* of dimension k , if M is locally homeomorphic to \mathbb{R}^k .

To explain, this means that any point $m \in M$ admits a neighbourhood U and a homeomorphism $\varphi: U \rightarrow V$, where V is an open subset of \mathbb{R}^k . The pair (U, φ) (or, sometimes just U) is called a chart on M near m .

Notice that the requirements that a manifold is Hausdorff and second countable are to a great extent of technical nature, whereas being locally homeomorphic to \mathbb{R}^k is a crucial property of manifolds.

Clearly, \mathbb{R}^k and in fact any open subset of \mathbb{R}^k are examples of topological manifolds of dimension k . As we have established in the introduction, 2-spheres are manifolds of dimension two. Similar arguments yield in fact that the k -sphere

$$S^k := \left\{ (x_1, \dots, x_{k+1}) \in \mathbb{R}^{k+1} \mid \sum_{j=1}^{k+1} x_j^2 = 1 \right\}$$

is a k -manifold.

Somewhat special is the case of dimension zero. Since \mathbb{R}^0 is by definition a single point, the above definition requires that each point of M has a neighborhood consisting only of this point. In other words, M is a countable discrete space.

Definition 2.2. A collection of charts $\mathcal{U} = \{(U_\alpha, \phi_\alpha) \mid \alpha \in A\}$ is called a C^0 -atlas, if $\bigcup_{\alpha \in A} U_\alpha = M$, that is if any point of M is contained in some chart. Here A is an arbitrary index set.

For example, \mathbb{R}^k admits a C^0 -atlas consisting of a single chart (\mathbb{R}^k, id) . In the introduction we have constructed a C^0 -atlas on the 2-sphere consisting of two charts. However, there is no C^0 -atlas on S^2 consisting of a single chart, since S^2 is not homeomorphic to an open subset of \mathbb{R}^2 (why?).

Given a C^0 -atlas \mathcal{U} , pick any two charts $(U_\alpha, \varphi_\alpha)$ and (U_β, φ_β) such that $U_\alpha \cap U_\beta \neq \emptyset$. The map

$$\theta_{\alpha\beta} := \varphi_\alpha \circ \varphi_\beta^{-1}: \varphi_\beta(U_\alpha \cap U_\beta) \rightarrow \varphi_\alpha(U_\alpha \cap U_\beta), \quad (2.3)$$

which is a homeomorphism between two open subsets of \mathbb{R}^k , is called a *coordinate transformation*¹. Notice that $\theta_{\beta\alpha}$ is the inverse map to $\theta_{\alpha\beta}$. In particular, $\theta_{\alpha\beta}$ is a homeomorphism between $\varphi_\beta(U_\alpha \cap U_\beta)$ and $\varphi_\alpha(U_\alpha \cap U_\beta)$.

It is a common practice to suppress the domain and the target of $\theta_{\alpha\beta}$ writing simply $\theta_{\alpha\beta} = \varphi_\alpha \circ \varphi_\beta^{-1}$. While this may be confusing at first, the advantage is that this allows us to suppress less important details so that the most essential features are clearer. If in doubt, the reader should write the domain and target explicitly.

Definition 2.4. A C^0 -atlas \mathcal{U} is called *smooth*, if all coordinate transformation maps $\theta_{\alpha\beta}$, $\alpha, \beta \in A$, are smooth.

Remark 2.5. Equally well, we can say that \mathcal{U} is a C^ℓ -atlas, if all coordinate transformation maps belong to $C^\ell(\mathbb{R}^n; \mathbb{R}^n)$ (keep in mind that these are defined on open subsets of \mathbb{R}^n only) for some fixed natural number ℓ . The theory does not depend much on the choice of ℓ as long as ℓ is not too small. In practice $\ell \geq 3$ would suffice in most of the cases, however to avoid non-essential details it is convenient to put $\ell = \infty$ from the very beginning.

Two charts (U, φ) and any (V, ψ) not necessarily from the same atlas are said to be *smoothly compatible* if the maps

$$\varphi \circ \psi^{-1} \quad \text{and} \quad \psi \circ \varphi^{-1} \quad (2.6)$$

are smooth, compare with (2.3). We consider two atlases \mathcal{U} and \mathcal{V} as “essentially equal”, if all charts from \mathcal{U} are smoothly compatible with all charts in \mathcal{V} . More formally, we have the following definition.

Definition 2.7. Two atlases \mathcal{U} and \mathcal{V} on the same underlying topological space M are called *equivalent*, if $\mathcal{U} \cup \mathcal{V}$ is a smooth atlas on M , that is if all charts from \mathcal{U} are smoothly compatible with all charts in \mathcal{V} . An equivalence class of atlases is called a *smooth structure* on M . A *smooth manifold* consists of a Hausdorff second countable topological space and a smooth structure.

To explain the point of the above definition, consider the 2-sphere. In the introduction we constructed a smooth atlas on S^2 , namely $\mathcal{U} := \{(S^2 \setminus \{N\}, \varphi_N), (S^2 \setminus \{S\}, \varphi_S)\}$. However, there are many ways to construct another smooth atlas, for example as follows:

$$\mathcal{U}' := \{S^2 \setminus \{N\}, \varphi_N\} \cup \{(S_+^2, \varphi_+)\}.$$

Here $S_+^2 := \{x \in S^2 \mid x_3 > 0\}$ and $\varphi_+(x) = (x_1, x_2)$.

Exercise 2.8. Check that \mathcal{U}' is a smooth atlas equivalent to \mathcal{U} .

It should be intuitively clear, that the description of S^2 via smooth atlases \mathcal{U} and \mathcal{U}' are ‘essentially equal’. Hence, it is natural to identify (S^2, \mathcal{U}) and (S^2, \mathcal{U}') .

An atlas \mathcal{U} is called *maximal*, if for any chart (V, ψ) smoothly compatible with all charts in \mathcal{U} is already contained in \mathcal{U} .

The importance of maximal atlases stems from the following result.

Lemma 2.9. *Each equivalence class of smooth atlases is represented by a unique maximal atlas.*

¹The origin of this terminology will be clear below.

Proof. For a smooth atlas \mathcal{U} on M define

$$\mathcal{U}_{max} := \{(V, \psi) \text{ is a chart on } M \text{ s.t. (2.6) are both smooth for all } (U, \varphi) \in \mathcal{U}\}.$$

Exercise 2.10. Check that \mathcal{U}_{max} is a smooth atlas on M .

By the construction of \mathcal{U}_{max} , we have $\mathcal{U} \subset \mathcal{U}_{max}$. Hence, any chart smoothly compatible with any chart in \mathcal{U}_{max} is also smoothly compatible with any chart in \mathcal{U} and therefore is contained in \mathcal{U}_{max} . Hence, \mathcal{U}_{max} is maximal. Clearly, \mathcal{U} and \mathcal{U}_{max} represent the same smooth structure. \square

By the above lemma, a smooth manifold may be considered as being equipped with a maximal atlas. In particular, if \mathcal{U} is any smooth atlas on M , we may freely add any chart smoothly compatible with all charts in \mathcal{U} without changing the smooth structure. For example, if (U, φ) is a chart near m_0 , then $(U, \hat{\varphi})$ with

$$\hat{\varphi}(m) = \varphi(m) - \varphi(m_0)$$

is also a chart near $m_0 \in M$ smoothly compatible with all charts in \mathcal{U} . The chart $(U, \hat{\varphi})$ satisfies

$$\hat{\varphi}(m_0) = 0,$$

which is commonly expressed by saying that $(U, \hat{\varphi})$ is *centered at* m_0 .

Remark 2.11. In what follows only smooth manifolds will be considered. Therefore, by saying that M is a manifold, we always mean a *smooth* manifold, unless explicitly stated otherwise.

Let us finish this section with some further examples of manifolds.

Example 2.12 (Products). Let M and N be smooth manifolds of dimensions k and ℓ respectively. Let $\mathcal{U} = \{(U_\alpha, \varphi_\alpha) \mid \alpha \in A\}$ and $\mathcal{V} = \{(V_\lambda, \psi_\lambda) \mid \lambda \in \Lambda\}$ be smooth atlases on M and N respectively. Then the product $M \times N$ is a Hausdorff second countable topological space. We define a C^0 -atlas on $M \times N$ by setting

$$\mathcal{W} := \{(U_\alpha \times V_\lambda, \varphi_\alpha \times \psi_\lambda) \mid \alpha \in A, \lambda \in \Lambda\}.$$

Given any two charts $(U_\alpha \times V_\lambda, \varphi_\alpha \times \psi_\lambda)$ and $(U_\beta \times V_\mu, \varphi_\beta \times \psi_\mu)$ the corresponding coordinate transformation is given by $\theta_{\alpha\beta} \times \eta_{\lambda\mu}$, where $\theta_{\alpha\beta} = \varphi_\alpha \circ \varphi_\beta^{-1}$ and $\eta_{\lambda\mu} = \psi_\lambda \circ \psi_\mu^{-1}$ are smooth maps. More precisely, this means the following:

$$\begin{aligned} \theta_{\alpha\beta} \times \eta_{\lambda\mu} &: \mathbb{R}^k \times \mathbb{R}^\ell \rightarrow \mathbb{R}^k \times \mathbb{R}^\ell, \\ \theta_{\alpha\beta} \times \eta_{\lambda\mu}(x, y) &= (\theta_{\alpha\beta}(x), \eta_{\lambda\mu}(y)), \quad x \in \mathbb{R}^k, y \in \mathbb{R}^\ell. \end{aligned}$$

In particular, $\theta_{\alpha\beta} \times \eta_{\lambda\mu}$ is a smooth map, which means that the atlas constructed above is smooth. Hence, $M \times N$ is a smooth manifold of dimension $k + \ell$. This yields in particular that the following

(i) the k -dimensional torus $\mathbb{T}^k := S^1 \times \cdots \times S^1$ and

(ii) the cylinder $\mathbb{R} \times S^1$

are smooth manifolds. In the latter case, the dimension of $\mathbb{R} \times S^1$ equals 2.

Example 2.13 (Real projective spaces). The real projective space \mathbb{RP}^k of dimension k is defined to be the set of all lines in \mathbb{R}^{k+1} through the origin. Since each line through the origin is uniquely determined by a point on this line distinct from the origin, we have

$$\mathbb{RP}^k = (\mathbb{R}^{k+1} \setminus \{0\}) / \sim,$$

where $x, y \in \mathbb{R}^{k+1} \setminus \{0\}$ are defined to be equivalent if and only if there exists $\lambda \in \mathbb{R} \setminus \{0\}$ such that $y = \lambda x$. In particular, we have the canonical surjective quotient map

$$\pi: \mathbb{R}^{k+1} \setminus \{0\} \rightarrow \mathbb{RP}^k, \quad \pi(x) = [x].$$

If $x = (x_0, x_1, \dots, x_k) \in \mathbb{R}^{k+1} \setminus \{0\}$, it is customary to write $[x_0 : x_1 : \dots : x_k]$ for $[x]$.

We endow \mathbb{RP}^k with the quotient topology, that is $U \subset \mathbb{RP}^k$ is open if and only if $\pi^{-1}(U)$ is open in $\mathbb{R}^{k+1} \setminus \{0\}$. It is straightforward to check that this yields a Hausdorff second countable topological space.

To construct a C^0 -atlas on \mathbb{RP}^k , observe that each

$$U_j := \{[x_0 : x_1 : \dots : x_k] \in \mathbb{RP}^k \mid x_j \neq 0\}, \quad j = 0, 1, \dots, k,$$

is an open subset of \mathbb{RP}^k . Indeed, this follows from the fact that

$$\pi^{-1}(U_j) = \{(x_0, \dots, x_k) \in \mathbb{R}^{k+1} \setminus \{0\} \mid x_j \neq 0\}$$

is an open subset of $\mathbb{R}^{k+1} \setminus \{0\}$.

The map

$$\begin{aligned} \varphi_j: U_j &\rightarrow \mathbb{R}^k, \\ \varphi_j[x_0 : x_1 : \dots : x_{j-1} : x_j : x_{j+1} : \dots : x_k] &= \left(\frac{x_0}{x_j}, \frac{x_1}{x_j}, \dots, \frac{x_{j-1}}{x_j}, \frac{x_{j+1}}{x_j}, \dots, \frac{x_k}{x_j} \right) \end{aligned}$$

is well-defined and continuous. Moreover, the map

$$\psi_j: \mathbb{R}^k \rightarrow U_j, \quad \psi_j(y_0, y_1, \dots, y_{k-1}) = [y_0 : y_1 : \dots : y_{j-1} : 1 : y_j : \dots : y_{k-1}] \quad (2.14)$$

is a continuous inverse of φ_j , that is φ_j is a homeomorphism. Since the collection U_0, \dots, U_k clearly covers all of \mathbb{RP}^k , $\mathcal{U} := \{(U_j, \varphi_j) \mid j = 0, 1, \dots, k\}$ is a C^0 -atlas on \mathbb{RP}^k .

Next, let us consider the coordinate transformations. To simplify the notations we consider only the map $\theta_{01} = \varphi_0 \circ \varphi_1^{-1} = \varphi_0 \circ \psi_1$. We have

$$\theta_{01}(y_0, \dots, y_{k-1}) = \varphi_0([y_0 : 1 : y_1, \dots, y_{k-1}]) = \left(\frac{1}{y_0}, \frac{y_1}{y_0}, \dots, \frac{y_{k-1}}{y_0} \right),$$

which is smooth on

$$\varphi_1(U_0 \cap U_1) = \{y \in \mathbb{R}^k \mid y_0 \neq 0\}.$$

A similar argument yields that all coordinate transformations $\theta_{ij} = \varphi_i \circ \psi_j$ are smooth on their domains of definition. Thus, \mathcal{U} is a smooth atlas and \mathbb{RP}^k is a smooth manifold of dimension k .

It may be useful to keep some non-examples of manifolds in mind.

- (a) The set $M = \{(x, y) \in \mathbb{R}^2 \mid x^2 = y^2\}$ consisting of two straight lines $y = \pm x$ intersecting at the origin, is not a manifold. Indeed, if M were a manifold, its dimension must be one. However, the origin does not have a neighbourhood in M homeomorphic to an open subset of \mathbb{R}^1 (Why?).

- (b) A disjoint union of manifolds is a manifold. However, a disjoint uncountable union of non-empty manifolds is *not* a manifold, since the second countability axiom is violated. For example, the disjoint union of real lines labeled by $\alpha \in (0, 1)$

$$N := \bigsqcup_{\alpha \in (0,1)} \mathbb{R}_\alpha$$

is not a manifold. Notice that the above example is *not* homeomorphic to $(0, 1) \times \mathbb{R}$, which is a manifold indeed, since, for example, each line $\mathbb{R}_\alpha \subset N$ is an open subset.

- (c) Consider the following line “with a double point”:

$$L := (-\infty, 0) \cup \{a, b\} \cup (0, +\infty).$$

Here $\{a, b\}$ is understood as a set consisting of two distinct elements. The following two subsets

$$U_a := (-\infty, 0) \cup \{a\} \cup (0, +\infty) \quad \text{and} \quad U_b := (-\infty, 0) \cup \{b\} \cup (0, +\infty)$$

cover all of L . Define $\varphi_a: U_a \rightarrow \mathbb{R}$ by $\varphi_a(x) = x$ if $x \neq a$ and $\varphi_a(a) = 0$. By the same token we can define $\varphi_b: U_b \rightarrow \mathbb{R}$.

A topology on L is defined simply by saying that V is open if and only if $\varphi_a(V \cap U_a)$ and $\varphi_b(V \cap U_b)$ are open in \mathbb{R} .

This yields a second countable topological space with a smooth atlas. However, L is non-Hausdorff.

L 1

2.2 Smooth maps

Given a smooth structure on M , we can make sense of smoothness of functions defined on M as follows.

Definition 2.15. Let M be a manifold with a smooth structure represented by a smooth atlas $\mathcal{U} = \{(U_\alpha, \varphi_\alpha)\}$. A function $f: M \rightarrow \mathbb{R}$ is said to be smooth, if for any chart $(U_\alpha, \varphi_\alpha)$ the function $f \circ \varphi_\alpha^{-1}: \mathbb{R}^k \rightarrow \mathbb{R}$ is smooth.

Notice that since an open subset V of M is again a smooth manifold, it makes sense to say that a function is smooth on V . The smoothness of functions is then a local property in the following sense: f is smooth if and only if the restriction of f to any open subset of M is smooth. In particular, if $\{V_\alpha \mid \alpha \in A\}$ is an open covering of M and f is smooth on each V_α , then f is smooth on M .

Strictly speaking, we still have to show that the notion of smoothness in Definition 2.15 is independent of the choice of an atlas. Indeed, assume that f is smooth with respect to \mathcal{U} and pick an atlas $\mathcal{V} = \{(V_\mu, \psi_\mu)\}$ equivalent to \mathcal{U} . Then on $\psi_\mu(U_\alpha \cap V_\mu) \subset \mathbb{R}^n$ we have

$$f \circ \psi_\mu^{-1}|_{\psi_\mu(U_\alpha \cap V_\mu)} = f \circ \varphi_\alpha^{-1} \circ \varphi_\alpha \circ \psi_\mu^{-1}|_{\psi_\mu(U_\alpha \cap V_\mu)} = f \circ \varphi_\alpha^{-1} \circ \theta_{\alpha\mu}|_{\psi_\mu(U_\alpha \cap V_\mu)},$$

where $\theta_{\alpha\mu} = \varphi_\alpha \circ \psi_\mu^{-1}$ is a smooth map. Hence, f is smooth with respect to \mathcal{V} on any subset $U_\alpha \cap V_\mu$. Since these subsets cover all of M , f is smooth on M with respect to \mathcal{V} .

Example 2.16. Let $F: \mathbb{R}^3 \rightarrow \mathbb{R}$ be any smooth function. Define $f: S^2 \rightarrow \mathbb{R}$ as the restriction of F to S^2 . I claim that f is a smooth function on S^2 . Indeed, let (U, φ) be any chart on S^2 constructed in the introduction. For concreteness, let us pick the chart $(S^2 \setminus \{N\}, \varphi_N)$. Then

$$f \circ \varphi_N^{-1}(y_1, y_2) = F\left(\frac{2y_1}{1 + y_1^2 + y_2^2}, \frac{2y_2}{1 + y_1^2 + y_2^2}, \frac{-1 + y_1^2 + y_2^2}{1 + y_1^2 + y_2^2}\right), \quad (y_1, y_2) \in \mathbb{R}^2.$$

Hence, $f \circ \varphi_N^{-1}$ is smooth and it is clear that this is also the case for $(S^2 \setminus \{S\}, \varphi_S)$. Thus, f is a smooth function on S^2 .

Example 2.17. Let $F: \mathbb{R}^{k+1} \setminus \{0\} \rightarrow \mathbb{R}$ be a smooth homogeneous function of degree 0, that is $F(\lambda x) = F(x)$ for all $\lambda \in \mathbb{R} \setminus \{0\}$ and $x \in \mathbb{R}^{k+1} \setminus \{0\}$. Define $f: \mathbb{RP}^k \rightarrow \mathbb{R}$ by setting $f([x]) = F(x)$. This yields a well-defined function, which I claim is smooth. Indeed, pick any chart (U_j, φ_j) constructed in Example 2.13. Using (2.14), we obtain

$$f \circ \varphi_j^{-1}(y_0, \dots, y_{k-1}) = F(y_0, \dots, y_{j-1}, 1, y_j, \dots, y_{k-1}),$$

which is smooth everywhere on \mathbb{R}^k . Hence, f is smooth.

Proposition 2.18. *The set $C^\infty(M)$ of all smooth functions on a manifold M is an algebra, that is*

- $f, g \in C^\infty(M), \lambda, \mu \in \mathbb{R} \implies \lambda f + \mu g \in C^\infty(M);$
- $f, g \in C^\infty(M) \implies f \cdot g \in C^\infty(M).$

Proof. Let f, g be any two smooth functions and λ, μ two real numbers. For any chart (U, φ) the functions

$$\begin{aligned} (\lambda f + \mu g) \circ \varphi^{-1} &= \lambda(f \circ \varphi^{-1}) + \mu(g \circ \varphi^{-1}), \\ (f \cdot g) \circ \varphi^{-1} &= f \circ \varphi^{-1} \cdot g \circ \varphi^{-1} \end{aligned}$$

are clearly smooth, hence $\lambda f + \mu g$ and $f \cdot g$ are smooth functions on M . \square

Let $f: M \rightarrow \mathbb{R}^\ell$ be a map, which can be written as an ℓ -tuple of functions: $f = (f_1, \dots, f_\ell)$. We say that f is smooth, if each component f_j is a smooth function on M .

It is also possible to define the notion of smoothness for maps between manifolds. To this end, let M and N be two manifolds of dimensions k and ℓ respectively. Pick an atlas \mathcal{U} on M and an atlas \mathcal{V} on N .

Definition 2.19. A continuous map $f: M \rightarrow N$ is said to be smooth, if for any $(U, \varphi) \in \mathcal{U}$ and any $(V, \psi) \in \mathcal{V}$ the map

$$\psi \circ f \circ \varphi^{-1}: \mathbb{R}^k \rightarrow \mathbb{R}^\ell$$

is smooth.

Remark 2.20. The requirement that f is continuous in the above definition is only needed to ensure that $\psi \circ f \circ \varphi^{-1}$ is defined on an open subset of \mathbb{R}^k . The map $\psi \circ f \circ \varphi^{-1}$ is called *the coordinate presentation of f* (with respect to charts (U, φ) and (V, ψ)).

The argument used to verify that the notion of smoothness of a function is well-defined is very common in the theory of manifolds and will be typically omitted below. However, the reader may wish to prove the following proposition as an exercise.

Proposition 2.21. *If $f: M \rightarrow N$ and $g: N \rightarrow L$ are smooth maps between smooth manifolds, then $g \circ f$ is also smooth.* \square

Definition 2.22. A smooth map $f: M \rightarrow N$ such that $f^{-1}: N \rightarrow M$ exists and is also smooth is called a *diffeomorphism*.

Observe that if (U, φ) is a chart, then $\varphi: U \rightarrow \varphi(U) \subset \mathbb{R}^k$ is a diffeomorphism.

To obtain a somewhat non-trivial example of a diffeomorphism, consider the tangent function:

$$\tan: (-\pi/2, \pi/2) \rightarrow \mathbb{R}, \quad \tan(x) = \frac{\sin x}{\cos x}.$$

This is smooth, bijective, the inverse function \arctan exists and is smooth. Hence the interval $(-\pi/2, \pi/2)$ is diffeomorphic to \mathbb{R} . In fact any open interval is diffeomorphic to \mathbb{R} (Why?).

A standard non-example is given by the map

$$f: \mathbb{R} \rightarrow \mathbb{R}, \quad f(x) = x^3,$$

which is clearly smooth and bijective. The inverse map however fails to be smooth at the origin so that f is *not* a diffeomorphism.

If there exists a diffeomorphism between M and N , we say that M and N are diffeomorphic. Notice that in this case we must have $k = \dim M = \dim N = \ell$. Indeed, if f is a diffeomorphism between M and N , then $F := \psi \circ f \circ \varphi^{-1}$ is a diffeomorphism between open subsets of \mathbb{R}^k and \mathbb{R}^ℓ . Let us denote $G := F^{-1} = \varphi \circ f^{-1} \circ \psi^{-1}$, which is smooth by the definition of smoothness for f^{-1} . Differentiating $G \circ F = \text{id}_{\mathbb{R}^k}$ at the point $G(x)$, we obtain

$$\text{id}_{\mathbb{R}^k} = D_{G(x)}(\text{id}_{\mathbb{R}^k}) = D_{G(x)}(G \circ F) = D_{F(G(x))}G \cdot D_{G(x)}F = D_x G \cdot D_{G(x)}F,$$

where $\text{id}_{\mathbb{R}^k}$ is the identity map. In particular, DG is surjective at each point. Furthermore, by a similar argument applied to the identity $F \circ G = \text{id}_{\mathbb{R}^\ell}$, we obtain that DG is injective at each point. In other words, $D_y G: \mathbb{R}^\ell \rightarrow \mathbb{R}^k$ is a linear isomorphism, which is only possible if $k = \ell$.

Definition 2.23. A map $f: M \rightarrow N$ is called a *local diffeomorphism*, if for any point $m \in M$ there exists an open neighbourhood $U \ni m$ in M and an open neighbourhood $V \ni f(m)$ in N such that

$$f|_U: U \rightarrow V$$

is a diffeomorphism.

A non-trivial example of a local diffeomorphism can be obtained as follows. The map

$$f: \mathbb{R} \rightarrow S^1, \quad f(x) = (\sin x, \cos x)$$

is a local diffeomorphism (why?), which is not a diffeomorphism, since $f(0) = f(\pm 2\pi) = f(\pm 4\pi) = \dots$

A non-trivial result from the course of analysis we need here is the following.

Theorem 2.24. Let U be open in \mathbb{R}^n and $f: U \rightarrow \mathbb{R}^k$ be smooth. Assume that at some $x \in U$ the differential $D_x f$ of f is invertible. Then $n = k$ and f is a local diffeomorphism at x , that is there exist open subsets $U' \ni x$ and $V \ni f(x)$ such that

$$f|_{U'}: U' \rightarrow V$$

is a diffeomorphism.

A proof of this theorem can be found for example in [BT03, Thm 9.4.1].

2.3 The fundamental theorem of algebra

As an application of the notions introduced in the preceding sections, we prove the fundamental theorem of algebra in this section. This requires some additional notions and constructions, which are of independent interest.

Let M and N be two manifolds of dimensions k and ℓ respectively. Pick a smooth map $f: M \rightarrow N$, a point $m \in M$ and charts (U_0, φ_0) and (V_0, ψ_0) such that $m \in U$ and $f(m) \in V$.

Definition 2.25. We say that m is a *critical point* of f if the differential of the coordinate representation

$$D_{\varphi_0(m)}(\psi_0 \circ f \circ \varphi_0^{-1}) \quad (2.26)$$

is non-surjective at $\varphi_0(m)$.

Lemma 2.27. *The notion of a critical point is well-defined, i.e., this is independent of the choice of charts.*

Proof. Pick any charts (U_1, φ_1) and (V_1, ψ_1) such that $m \in U_1$ and $f(m) \in V_1$. We have

$$\psi_0 \circ f \circ \varphi_0^{-1} = \psi_0 \circ \psi_1^{-1} \circ \psi_1 \circ f \circ \varphi_1^{-1} \circ \varphi_1 \circ \varphi_0^{-1} = \theta_{01}^\psi \circ (\psi_1 \circ f \circ \varphi_1^{-1}) \circ \theta_{10}^\varphi,$$

which is valid on an open subset containing m . Hence, by the chain rule, we obtain for the differentials

$$D(\psi_0 \circ f \circ \varphi_0^{-1}) = D\theta_{01}^\psi \circ D(\psi_1 \circ f \circ \varphi_1^{-1}) \circ D\theta_{10}^\varphi.$$

Since $D\theta_{01}^\psi$ and $D\theta_{10}^\varphi$ are invertible everywhere on the domain of their definition, we obtain that $D(\psi_0 \circ f \circ \varphi_0^{-1})$ is non-surjective at $\varphi_0(m)$ if and only if $D(\psi_1 \circ f \circ \varphi_1^{-1})$ is non-surjective at $\varphi_1(m)$. This finishes the proof of this lemma. \square

Definition 2.28. Any non-critical point is called *regular*. We say that $n \in N$ is a *regular value* of f , if $f^{-1}(n)$ consists of regular points only. If $f^{-1}(n)$ contains at least one singular point, then n is called a *singular value* of f .

Let me stress that any point, which does not lie in the image of f , is a regular value of f (this fact of course follows from the definition but may be easily missed at first).

Notice that in the particular case $k = \ell$, (2.26) is a linear map $\mathbb{R}^k \rightarrow \mathbb{R}^k$. Hence, (2.26) is non-surjective if and only if it has a non-trivial kernel, or, still if and only if $\det D_{\varphi_0(m)}(\psi_0 \circ f \circ \varphi_0^{-1}) = 0$.

With these preliminaries at hand, we can prove the following.

Theorem 2.29 (The fundamental theorem of algebra). *Let $p(z) := a_k z^k + a_{k-1} z^{k-1} + \dots + a_1 z + a_0$ be a polynomial with complex coefficients of degree $k \geq 1$. Then p has at least one (complex) root.*

Proof. Identify \mathbb{R}^2 with \mathbb{C} by writing

$$y = (y_1, y_2) \equiv y_1 + y_2 i = z.$$

For a fixed polynomial $p = a_k z^k + \dots + a_1 z + a_0$ such that $a_k \neq 0$, where $k \geq 1$, define a map $f: S^2 \rightarrow S^2$ by the rule

$$f(x) = \begin{cases} N, & \text{if } x = N, \\ \varphi_N^{-1} \circ p \circ \varphi_N(x), & \text{if } x \neq N. \end{cases}$$

The proof proceeds in a number of steps.

Step 1. f is smooth.

It is enough to check that f is smooth near N , i.e., that $\varphi_S \circ f \circ \varphi_S^{-1}$ is smooth. To see this, consider

$$\varphi_S \circ f \circ \varphi_S^{-1} = \varphi_S \circ \varphi_N^{-1} \circ \varphi_N \circ f \circ \varphi_N^{-1} \circ \varphi_N \circ \varphi_S^{-1} = \theta_{SN} \circ p \circ \theta_{SN}^{-1},$$

which is valid on $\mathbb{C} \setminus \{0\}$. By (1.2),

$$\theta_{SN}(z) = \frac{z}{z\bar{z}} = \frac{1}{\bar{z}}.$$

Consequently, for $z \neq 0$ we have

$$\varphi_S \circ f \circ \varphi_S^{-1}(z) = \frac{1}{p(1/\bar{z})} = \frac{1}{\frac{\bar{a}_k}{z^k} + \dots + \frac{\bar{a}_1}{z} + \bar{a}_0} = \frac{z^k}{\bar{a}_k + \dots + \bar{a}_1 z^{k-1} + \bar{a}_0 z^k}.$$

Since $\varphi_S \circ f \circ \varphi_S^{-1}(0) = 0$, $\varphi_S \circ f \circ \varphi_S^{-1}$ is clearly smooth on \mathbb{C} . Hence, f is smooth as claimed.

Step 2. The differential of the map $z \mapsto p(z)$ at the point z can be identified with $h \mapsto p'(z)h$, where

$$p'(z) = ka_k z^{k-1} + \dots + 2a_2 z + a_1.$$

Denote $p(z) = u(y_1, y_2) + v(y_1, y_2)i$. Since p is a holomorphic function of z , by the Cauchy-Riemann equations we have

$$\begin{aligned} D_z p(h) &= \begin{pmatrix} \frac{\partial u}{\partial y_1} & \frac{\partial u}{\partial y_2} \\ \frac{\partial v}{\partial y_1} & \frac{\partial v}{\partial y_2} \end{pmatrix} \begin{pmatrix} h_1 \\ h_2 \end{pmatrix} = \begin{pmatrix} \frac{\partial u}{\partial y_1} h_1 - \frac{\partial v}{\partial y_1} h_2 \\ \frac{\partial v}{\partial y_1} h_1 + \frac{\partial u}{\partial y_1} h_2 \end{pmatrix} \\ &\equiv \left(\frac{\partial u}{\partial y_1} h_1 - \frac{\partial v}{\partial y_1} h_2 \right) + \left(\frac{\partial v}{\partial y_1} h_1 + \frac{\partial u}{\partial y_1} h_2 \right) i \\ &= \frac{\partial p}{\partial y_1} h = p'(z) h. \end{aligned}$$

Step 3. The set of critical values of f is finite.

Indeed, any critical point of f is either N or of the form $\varphi_N^{-1}(z)$, where z is a zero of the polynomial

$$p'(z) = ka_k z^{k-1} + \dots + 2a_2 z + a_1.$$

Hence, the number of critical points of f is finite and therefore the number of critical values is also finite.

Remark 2.30. Notice that N is a critical point of f as long as $k \geq 2$, which we can assume without loss of generality, since any polynomial of degree 1 obviously has a root.

Step 4. For any regular value $x \in S^2$, the number of points in $f^{-1}(x)$ is finite and independent of x .

First notice that the set $\mathcal{R}(f)$ of regular values is open and connected as a complement of a finite number of points in S^2 .

Furthermore, for any $n \in \mathcal{R}(f)$ and any $m \in f^{-1}(n)$, the map $\varphi_N \circ f \circ \varphi_N^{-1} = p$ is a local diffeomorphism at $\varphi(m)$ by **Theorem 2.24**. Since φ_N is a homeomorphism, f is a local homeomorphism at m . In particular, there is a neighbourhood $U' \ni m$ such that $U' \cap f^{-1}(n) =$

$\{m\}$. Thus, $f^{-1}(n)$ is discrete. Since S^2 is compact, $f^{-1}(n)$ is also compact as a closed subset of a compact space. Hence, $f^{-1}(n)$ must in fact be a finite set.

The above argument actually shows that the map

$$\mathcal{R}(f) \ni x \mapsto \#f^{-1}(x) \in \mathbb{N} \quad (2.31)$$

is locally constant. Since $\mathcal{R}(f)$ is connected, this function must be constant.

Step 5. *We prove this theorem.*

Observe first that (2.31) cannot vanish everywhere on $\mathcal{R}(f)$. Indeed, the image of f is obviously infinite, whereas the set of critical values is finite by Step 3. Hence, there are regular values, which are in the image of f .

In fact, (2.31) vanishes nowhere as a locally constant function on an open connected space. Hence, $f^{-1}(S) \neq \emptyset$ as long as $S \in \mathcal{R}(f)$. Also, if S is a critical value, then $f^{-1}(S)$ contains at least one critical point. In either case, $f^{-1}(S)$ is non-empty, which means that p has at least one root. \square

The above proof turns out to contain a few ideas which can be used in other circumstances too. However, this requires some technical results, which are proved first.

L 2

2.4 Tangent spaces

We begin with the following consideration. Let γ be a smooth curve in \mathbb{R}^k through some $p \in \mathbb{R}^k$, that is a smooth map $\gamma: (a, b) \rightarrow \mathbb{R}^k$, such that $\gamma(t_0) = p$ for some $t_0 \in (a, b)$. Recall that the tangent vector of γ at p is

$$\dot{\gamma}(t_0) := \left. \frac{d}{dt} \right|_{t=t_0} \gamma(t) \in \mathbb{R}^k.$$

Let now γ be a smooth curve on the 2-sphere through some $p \in S^2$. Since S^2 is a subset of \mathbb{R}^3 , we may think of γ as a curve in \mathbb{R}^3 satisfying

$$\gamma_1^2(t) + \gamma_2^2(t) + \gamma_3^2(t) = 1 \quad \forall t \in (a, b). \quad (2.32)$$

It is reasonable to call the set

$$T_p S^2 := \{v \in \mathbb{R}^3 \mid v \text{ is the tangent vector of some smooth curve on } S^2 \text{ through } p\}$$

the tangent space of S^2 at the point p .

To determine $T_p S^2$ more explicitly, differentiate (2.32) with respect to t and set $t = t_0$:

$$\gamma_1(t_0)\dot{\gamma}_1(t_0) + \gamma_2(t_0)\dot{\gamma}_2(t_0) + \gamma_3(t_0)\dot{\gamma}_3(t_0) = 0 \quad \Longleftrightarrow \quad \langle p, \dot{\gamma}(t_0) \rangle = 0.$$

In other words, the tangent vector of any smooth curve on S^2 through p is necessarily orthogonal to p . Moreover, it is clear that any vector orthogonal to p arises in this way (consider all great circles through p). Hence,

$$T_p S^2 = p^\perp.$$

However, for an abstract manifold M a smooth curve γ on M does not lie in an Euclidean space in any obvious way so that the above definition of $T_p S^2$ does not immediately generalize. A nice workaround goes as follows.

Let γ be a smooth curve through $m \in M$. We can assume that γ is defined on $(-\varepsilon, \varepsilon)$ for some $\varepsilon > 0$ and $\gamma(0) = m$.

Definition 2.33. Two smooth curves γ_1 and γ_2 through m as above are said to be equivalent, if for some chart (U, φ) such that $m \in U$ we have

$$\left. \frac{d}{dt} \right|_{t=0} (\varphi \circ \gamma_1(t)) = \left. \frac{d}{dt} \right|_{t=0} (\varphi \circ \gamma_2(t)). \quad (2.34)$$

In other words, $\gamma_1 \sim \gamma_2$ if and only if the smooth curves $\varphi \circ \gamma_1$ and $\varphi \circ \gamma_2$ in \mathbb{R}^k (!) have the same tangent vector.

Furthermore, an equivalence class of smooth curves is called a *tangent vector* at m . The set $T_m M$ of all tangent vectors at m is called *the tangent space* at m .

Exercise 2.35. Show that the above equivalence relation is independent of the choice of a chart.

Exercise 2.36. Show that the tangent space to S^2 at some $p \in S^2$ in the sense of Definition 2.33 is p^\perp .

While Definition 2.33 is has a clear geometric meaning, the algebraic structure of $T_m M$ is opaque in this approach. For this reason we adopt an alternative definition, which is more algebraic.

First notice that given any tangent vector $[\gamma]$ through m we can define the map $\partial_{[\gamma]}: C^\infty(M) \rightarrow \mathbb{R}$ by setting

$$\partial_{[\gamma]}(f) = \left. \frac{d}{dt} \right|_{t=0} f(\gamma(t)) = \lim_{t \rightarrow 0} \frac{f(\gamma(t)) - f(m)}{t}.$$

Proposition 2.37. The map $\partial_{[\gamma]}$ is well defined and has the following properties:

(i) $\partial_{[\gamma]}$ is \mathbb{R} -linear, that is

$$\partial_{[\gamma]}(\lambda f + \mu g) = \lambda \partial_{[\gamma]}(f) + \mu \partial_{[\gamma]}(g)$$

holds for all $\lambda, \mu \in \mathbb{R}$ and $f, g \in C^\infty(M)$.

(ii) $\partial_{[\gamma]}$ satisfies

$$\partial_{[\gamma]}(fg) = \partial_{[\gamma]}(f)g(m) + f(m)\partial_{[\gamma]}(g)$$

for all $f, g \in C^\infty(M)$.

Proof. We only need to prove that $\partial_{[\gamma]}$ is well-defined, since (i) and (ii) are clear from the definition.

Thus, pick two smooth equivalent curves γ_1 and γ_2 through m . Pick also a chart (U, φ) near m and denote $F := f \circ \varphi^{-1}: \mathbb{R}^k \rightarrow \mathbb{R}$ and $\beta_j := \varphi \circ \gamma_j: (-\varepsilon, \varepsilon) \rightarrow \mathbb{R}^k$. Notice, that

$$\dot{\beta}_1(0) = \dot{\beta}_2(0) =: v,$$

since γ_1 and γ_2 are equivalent.

We have

$$\left. \frac{d}{dt} \right|_{t=0} (f \circ \gamma_1(t)) = \left. \frac{d}{dt} \right|_{t=0} (f \circ \varphi^{-1} \circ \varphi \circ \gamma_1(t)) = \left. \frac{d}{dt} \right|_{t=0} (F \circ \beta_1(t)) = D_v F(\varphi(m)),$$

where $D_v F$ is the derivative of F in the direction of v , that is

$$D_v F = \sum_{j=1}^k \frac{\partial F}{\partial x_j} v_j = \langle \nabla F, v \rangle. \quad (2.38)$$

A similar computation yields also

$$\left. \frac{d}{dt} \right|_{t=0} (f \circ \gamma_2(t)) = D_v F(\varphi(m)),$$

thus demonstrating that $\partial_{[\gamma]}$ depends on the equivalence class of γ only as the notation suggests. \square

Motivated by the above proposition we give the following

Definition 2.39. An \mathbb{R} -linear map $\partial: C^\infty(M) \rightarrow \mathbb{R}$ satisfying the Leibnitz rule

$$\partial(fg) = \partial(f)g(m) + f(m)\partial(g) \quad \forall f, g \in C^\infty(M)$$

is called a *derivation at m* .

Notice that a constant function, which takes value 1 everywhere on M , is annihilated by any derivation. Indeed, this follows from the following computation:

$$\partial(1) = \partial(1^2) = \partial(1) \cdot 1 + 1 \cdot \partial(1) = 2\partial(1).$$

By the linearity of derivations, any constant function is annihilated by each derivation.

For the proof of **Proposition 2.41** below, we need the following technical result, whose proof is deferred till the next section.

Proposition 2.40. *For any manifold M and any point $m_0 \in M$ the following holds.*

- (i) *Suppose f is a smooth function defined on a neighborhood U of m_0 . Then there is a smooth function \hat{f} defined everywhere on M and a neighborhood $\hat{U} \subset U$ of m_0 such that f and \hat{f} coincide everywhere on \hat{U} .*
- (ii) *Let ∂ be a derivation at m_0 . If f and \hat{f} are two smooth functions defined everywhere on M such that the restrictions of f and \hat{f} to some neighborhood \hat{U} of m_0 are equal, then $\partial(f) = \partial(\hat{f})$.*

Thus, for any tangent vector at m we constructed an explicit derivation at m . It turns out that this map is a bijection as we show next.

Denote temporarily by $Der_m M$ the set of all derivations at m .

Proposition 2.41. *The map*

$$T_m M \rightarrow Der_m M, \quad [\gamma] \mapsto \partial_{[\gamma]} \tag{2.42}$$

is a bijection.

Proof. We continue to use notations of the proof of **Proposition 2.37**. In addition, the chart (U, φ) is assumed to be centered at m_0 .

Step 1. (2.42) *is injective.*

Assume $\partial_{[\gamma_1]} = \partial_{[\gamma_2]}$, that is $\partial_{[\gamma_1]}(f) = \partial_{[\gamma_2]}(f)$ holds for any $f \in C^\infty(M)$. This implies in turn that

$$D_{v_1} F(\varphi(m)) = D_{v_2} F(\varphi(m))$$

holds for any $F \in C^\infty(\mathbb{R}^k)$, where $v_j = \dot{\beta}_j(0)$. Substituting $F = x_j$ in the above equality, we obtain that the j th components of v_1 and v_2 are equal for any j , i.e., $v_1 = v_2$, which yields in turn that γ_1 and γ_2 are equivalent. Hence, the injectivity of (2.42) follows.

Step 2. For any $F \in C^\infty(\mathbb{R}^k)$ there exist smooth functions G_1, \dots, G_k such that

$$F(x) = F(0) + \sum_{j=1}^k x_j G_j(x). \quad (2.43)$$

This follows by the following computation:

$$F(x) - F(0) = \int_0^1 \frac{d}{dt} F(tx) dt = \int_0^1 \langle \nabla F(tx), x \rangle dt,$$

which yields (2.43) with $G_j(x) = \int_0^1 \frac{\partial F}{\partial x_j}(tx) dt$.

Step 3. (2.42) is surjective.

Denote the j th component of φ by x_j so that $\varphi = (x_1, \dots, x_k)$. Notice that each x_j is a smooth function defined on U .

By Proposition 2.40, we can find $\hat{U} \subset U$ and a smooth function \hat{x}_j defined everywhere on U such that x_j and \hat{x}_j coincide on \hat{U} .

Pick any $\partial \in \text{Der}_m M$ and define

$$v_j := \partial(\hat{x}_j) \in \mathbb{R}, \quad v = (v_1, \dots, v_k),$$

Notice that by Proposition 2.40, (ii), v_j does not depend on the choice of \hat{x}_j .

Furthermore, define

$$\begin{aligned} \beta: (-\varepsilon, \varepsilon) &\rightarrow \mathbb{R}^k, & \beta(t) &= (v_1 t, \dots, v_k t), \\ \gamma: (-\varepsilon, \varepsilon) &\rightarrow M, & \gamma &:= \varphi^{-1} \circ \beta. \end{aligned}$$

By the previous step, there exist some functions $G_j: \mathbb{R}^k \rightarrow \mathbb{R}$ such that $f \circ \varphi^{-1}(x) = \sum x_j G_j(x)$. Hence,

$$f = f(m) + \sum_{j=1}^k x_j g_j, \quad (2.44)$$

where we think of x_j as a function on U and $g_j = G_j \circ \varphi$. In particular, $x_j(m) = 0$ for all $j = 1, \dots, k$.

Applying Proposition 2.40, (i) again, we can find some \hat{g}_j defined globally on M and a neighbourhood² $\hat{U} \subset U$ such that

$$f|_{\hat{U}} = f(m) + \sum_{j=1}^k \hat{x}_j \hat{g}_j|_{\hat{U}} = f(m) + \sum_{j=1}^k x_j g_j|_{\hat{U}}.$$

By Proposition 2.40, (ii) we obtain

$$\begin{aligned} \partial(f) &= \partial(f(m)) + \sum_{j=1}^k \partial(x_j) g_j(m) + x_j(m) \partial(g_j) \\ &= 0 + \sum_{j=1}^k v_j g_j(m) + 0 = \sum_{j=1}^k v_j g_j(m). \end{aligned} \quad (2.45)$$

²Shrinking the neighbourhoods if necessary, without loss of generality we can assume that \hat{U} is the same neighbourhood for all \hat{x}_j and g_j .

Furthermore, recalling (2.38), we obtain

$$\begin{aligned}\partial_{[\gamma]}(f) &= \sum_{j=1}^k v_j \frac{\partial F}{\partial x_j}(0) = \sum_{j=1}^k v_j \frac{\partial}{\partial x_j} \Big|_{x=0} \left(f(m) + \sum_{i=1}^k x_i G_i(x) \right) \\ &= \sum_{j=1}^k v_j \left(0 + \sum_{i=1}^k \delta_{ij} G_i(0) \right) = \sum_{j=1}^k v_j g_j(m).\end{aligned}$$

Comparing this with (2.45), we conclude that $\partial(f) = \partial_{[\gamma]}(f)$ holds for any $f \in C^\infty(M)$. Hence, $\partial = \partial_{[\gamma]}$, which finishes the proof of this step and the proof of this proposition too. \square

We use the bijective map of Proposition 2.41 to identify $T_m M$ with $Der_m M$. Since $Der_m M$ is clearly a vector space, we obtain the structure of a vector space on $T_m M$ in this way. Also, in view of this identification, we drop the notation $Der_m M$ in favour of $T_m M$ and we will switch freely between the two interpretations of tangent vectors as classes of curves and derivations.

Proposition 2.46. *For any $m \in M$ the tangent space $T_m M$ is a vector space of dimension $k = \dim M$.*

Proof. Pick a chart (U, φ) , $\varphi = (x_1, \dots, x_k)$ centered at m as in the proof of Proposition 2.37. For each $j = 1, \dots, k$ define a curve γ_j by

$$\varphi \circ \gamma_j(t) = (0, \dots, 0, t, 0, \dots, 0),$$

where the only non-trivial component is on the j th place. Correspondingly, we have k derivations:

$$\partial_j := \partial_{[\gamma_j]}.$$

Notice that if F is the coordinate representation of f , we have

$$\partial_j(f) = \frac{d}{dt} \Big|_{t=0} f \circ \gamma_j(t) = \frac{d}{dt} \Big|_{t=0} F(0, \dots, 0, t, 0, \dots, 0) = \frac{\partial F}{\partial x_j}(0).$$

We want to show that $\partial_1, \dots, \partial_k$ is a basis of $T_m M$.

To show that $\partial_1, \dots, \partial_k$ are linearly independent, assume there are some real numbers $\lambda_1, \dots, \lambda_k$ such that $\lambda_1 \partial_1 + \dots + \lambda_k \partial_k = 0$, that is

$$\lambda_1 \partial_1(f) + \dots + \lambda_k \partial_k(f) = 0$$

holds for any $f \in C^\infty(M)$. Substituting $f = x_j$ in the above equality³, we obtain $\lambda_j = 0$. Hence, $\partial_1, \dots, \partial_k$ are linearly independent indeed.

Let us show that any derivation ∂ at m can be represented as a linear combination of $\partial_1, \dots, \partial_k$. By Proposition 2.41, there exists a curve γ through m such that $\partial = \partial_{[\gamma]}$. If $\beta = \varphi \circ \gamma$ is a coordinate representation of γ and $v = \dot{\beta}(0)$, then

$$\partial(f) = D_v F = \sum v_j \frac{\partial F}{\partial x_j}(0) = \sum v_j \partial_j(f).$$

Hence, $\partial = v_1 \partial_1 + \dots + v_k \partial_k$. \square

Notice that the proof of the above proposition yields in fact a basis of $T_m M$ for any choice of a chart (U, φ) such that $m \in U$. In fact, we have shown that $\partial_1, \dots, \partial_k$ is a basis of $T_m M$, where

$$\partial_j f = \frac{\partial}{\partial x_j} \Big|_{x=\varphi(m)} (f \circ \varphi^{-1}(x)). \quad (2.47)$$

L 3

Also, in the particular case $M = \mathbb{R}^k$ the proof of [Proposition 2.41](#) shows that we have a canonical isomorphism

$$T_m \mathbb{R}^k \rightarrow \mathbb{R}^k, \quad \partial \mapsto (\partial(x_1), \dots, \partial(x_k)). \quad (2.48)$$

Indeed, denote $v_j = \partial(x_j)$ and $v = (v_1, \dots, v_k)$. Assuming m is the origin and writing $F(x) = F(0) + \sum x_j G_j(x)$ as in [\(2.43\)](#), we obtain

$$\partial(F) = \partial(F(0)) + \sum_{j=1}^k \left(\partial(x_j) G_j(0) + x_j(0) \partial(G_j) \right) = \sum_{j=1}^k v_j \frac{\partial F}{\partial x_j}(0) = D_v F(0)$$

Hence, if $v = 0$, then $\partial(F) = 0$ for all $F \in C^\infty(\mathbb{R}^k)$. In other words, [\(2.48\)](#) is injective. This map is also surjective, since the image of $\partial = D_v$ equals v .

This isomorphism is particularly clear, if we interpret tangent vectors as classes of curves. Indeed, if $\partial = \partial_{[\gamma]}$, then $\partial_{[\gamma]}(x_j) = \dot{\gamma}_j(0)$ so that [\(2.48\)](#) becomes

$$[\gamma] \mapsto \dot{\gamma}(0). \quad (2.49)$$

Example 2.50. Let $f: \mathbb{R}^{k+1} \rightarrow \mathbb{R}$ be a smooth function. Assume that 0 is a regular value of f . We shall show below that

$$M := f^{-1}(0)$$

is a smooth k -manifold. Taking this as granted for now, we can ask the following question: Given $m \in M$, can we describe the tangent space $T_m M$ more explicitly?

Notice that we have a natural linear map

$$\iota: T_m M \rightarrow \text{Der}_m \mathbb{R}^{k+1} \cong \mathbb{R}^{k+1}, \quad \iota(\partial)h = \partial(h|_M). \quad (2.51)$$

If a tangent vector is interpreted as a class of curves, then we have

$$\iota(\partial_{[\gamma]}) = (\partial_{[\gamma]}(x_1), \dots, \partial_{[\gamma]}(x_{k+1})) = \dot{\gamma}(0) \quad \text{or} \quad \iota([\gamma]) = \dot{\gamma}(0). \quad (2.52)$$

This map is injective but not surjective. Indeed, viewing γ as a curve in \mathbb{R}^{k+1} , this satisfies $f(\gamma(t)) = 0$ for all t . By differentiating this equation with respect to t , we obtain

$$\langle \nabla f(m), \dot{\gamma}(0) \rangle = 0.$$

Hence, $\dot{\gamma}(0)$ lies in the orthogonal complement to $\nabla f(m)$. Since $\dim T_m M = \dim M = k = \dim (\nabla f(m))^\perp$, we obtain

$$\iota(T_m M) = (\nabla f(m))^\perp.$$

Typically, this is expressed simply as $T_m M = (\nabla f(m))^\perp$.

An interested reader may find the following exercise to be instructive: Show that the image of [\(2.51\)](#) equals $(\nabla f(m))^\perp$ directly, that is without using [Proposition 2.41](#).

³Technically, we should first fix an “extension” \hat{x}_j of x_j as in the proof of [Proposition 2.41](#) above. However, this should be clear by now and we will omit this sort of argument below.

2.5 Cut off and bump functions

Recall that the function

$$\lambda: \mathbb{R} \rightarrow \mathbb{R}, \quad \lambda(t) := \begin{cases} 0, & \text{if } t \leq 0, \\ e^{-\frac{1}{t}}, & \text{if } t > 0 \end{cases}$$

is smooth everywhere on \mathbb{R} including $t = 0$.

Furthermore, notice that for any $r > 0$ we have $\lambda(t) + \lambda(r - t) > 0$ for any $t \in \mathbb{R}$. Indeed, for positive t the first term is positive and for negative t the second one is positive. Using this we define the function

$$\hat{\chi}_r(t) := \frac{\lambda(r - t)}{\lambda(t) + \lambda(r - t)},$$

where $r > 0$ is a parameter. Notice that $\hat{\chi}_r$ is smooth everywhere on \mathbb{R} , takes values in $[0, 1]$, $\hat{\chi}_r(t) = 0$ for $t \geq r$, and $\hat{\chi}_r(t) = 1$ for all $t \leq 0$. It is convenient to define

$$\chi_r(t) := \hat{\chi}_r(t - 1) = \frac{\lambda(r + 1 - t)}{\lambda(t - 1) + \lambda(r + 1 - t)}, \quad (2.53)$$

which is called a *cut off function*. The graph of χ_r is shown schematically on Figure 2.1 below.




Figure 2.1: Graph of χ_r .

Proposition 2.54. *For any point m_0 on M and any neighbourhood $U \ni m_0$ there exists a neighbourhood $V \subset U$ and a smooth function $\rho: M \rightarrow [0, 1]$ such that*

$$\rho|_V \equiv 1 \quad \text{and} \quad \rho|_{M \setminus U} \equiv 0.$$

Proof. Pick a chart centered at m_0 . Without loss of generality, we can assume that the local homeomorphism φ is defined everywhere on U . We can find $R > 0$ such that the ball $B_{2R}(0) := \{x \in \mathbb{R}^k \mid |x| < 2R\}$ is contained in $\varphi(U)$.

Furthermore, the function

$$\hat{\rho}(x) = \chi_1\left(\frac{|x|}{R}\right)$$

is smooth, equals 1 on $B_R(0)$ and vanishes outside of $B_{2R}(0)$. Hence,

$$\rho(m) = \begin{cases} \hat{\rho}(\varphi(m)), & \text{if } m \in U, \\ 0, & \text{otherwise,} \end{cases}$$

is a well-defined smooth function, which equals 1 on $\varphi^{-1}(B_R(0))$ and vanishes outside of $\varphi^{-1}(B_{2R}(0))$. \square

The function ρ provided by the above proposition is called *a bump function*.

With these preliminaries at hand, we are ready to prove **Proposition 2.40**.

Proof of Proposition 2.40. Let V and ρ be as in **Proposition 2.54**. The function

$$\hat{f}(m) = \begin{cases} f(m) \cdot \rho(m) & \text{if } m \in U \\ 0 & \text{otherwise,} \end{cases}$$

has the required properties. This proves (i).

Let us prove (ii). We can assume that $V \subset \hat{U}$ and, moreover, that ρ vanishes outside of \hat{U} . Then the function $(\hat{f} - f) \cdot \rho$ vanishes everywhere and therefore for any derivation ∂ at m_0 we have

$$0 = \partial((\hat{f} - f) \cdot \rho) = \partial(\hat{f} - f) \cdot \rho(m_0) + (\hat{f}(m_0) - f(m_0))\partial(\rho) = \partial(\hat{f}) - \partial(f).$$

This proves (ii). \square

Exercise 2.55. Show that the open ball

$$B := \{x \in \mathbb{R}^k \mid |x| < 1\}$$

in \mathbb{R}^k is diffeomorphic to \mathbb{R}^k . Deduce from this that any point on a manifold admits a chart (U, φ) such that $\varphi: U \rightarrow \mathbb{R}^k$ is a diffeomorphism. (*Hint:* Show first that there is a diffeomorphism $f: (0, 1) \rightarrow (0, \infty)$ such that $f(r) = r$ for all $r \leq r_0 < 1$.)

2.6 The differential of a smooth map

Let $f: M^k \rightarrow N^\ell$ be a smooth map between two smooth manifolds.

Definition 2.56. For any $m \in M$ the map

$$f_*(m): T_m M \rightarrow T_{f(m)} N \quad \text{defined by} \quad f_*(p)[\gamma] = [f \circ \gamma]$$

is called *the differential of f at the point m* . Here γ is a smooth curve through m .

Think of a tangent vector at $m \in M$ as a derivation ∂ at m . By **Proposition 2.41**, there exists a smooth curve γ through m such that $\partial = \partial_{[\gamma]}$. Then we have $\partial_{f_*(m)[\gamma]} = \partial_{[f \circ \gamma]}$. Hence, for any $h \in C^\infty(N)$ we obtain

$$\partial_{f_*(m)[\gamma]}(h) = \partial_{[f \circ \gamma]}(h) = \frac{d}{dt} \Big|_{t=0} (h \circ (f \circ \gamma))(t) = \frac{d}{dt} \Big|_{t=0} ((h \circ f) \circ \gamma)(t) = \partial_{[\gamma]}(h \circ f).$$

Hence, thinking of tangent vectors at m as derivations at m we can identify the differential of f at m with the map

$$\partial \mapsto f_* \partial, \quad \text{where} \quad (f_* \partial)h = \partial(h \circ f). \quad (2.57)$$

More precisely, this means that the following diagram commutes:

$$\begin{array}{ccc} T_m M & \xrightarrow{f_*(m)} & T_{f(m)} N \\ \downarrow & & \downarrow \\ \text{Der}_m M & \xrightarrow{f_*(m)} & \text{Der}_{f(m)} N. \end{array}$$

Here the vertical arrows are given by (2.42), the upper horizontal arrow represents the differential of f in the sense of Definition 2.56, whereas the lower arrow represents (2.57).

Since (2.57) is obviously a linear map, we obtain the following.

Proposition 2.58. *The differential is a linear map.* \square

Pick a chart (U, φ) on M centered at m and a chart (V, ψ) on N centered at $f(m) \in V$. Write $\varphi = (x_1, \dots, x_k)$ and $\psi = (y_1, \dots, y_\ell)$, where x_i are functions on U and y_j are functions on V just like in the proof of Equation 2.42. By the proof of Proposition 2.46, we obtain the following bases

$$(\partial_1^x, \dots, \partial_k^x) = \left(\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_k} \right) \quad \text{and} \quad (\partial_1^y, \dots, \partial_\ell^y) = \left(\frac{\partial}{\partial y_1}, \dots, \frac{\partial}{\partial y_\ell} \right). \quad (2.59)$$

It is worthwhile to recall that $\partial_i^x g = \frac{\partial G}{\partial x_i}(0)$, where $G = g \circ \varphi^{-1}$ is the coordinate representation of g .

Since the differential is a linear map, this can be represented by a matrix relative to the above bases. Thus, a natural question arises: Can we compute the matrix of the differential relative to Bases (2.59)?

To answer this question, recall that the coordinate representation of f is

$$F = \psi \circ f \circ \varphi^{-1}: \mathbb{R}^k \rightarrow \mathbb{R}^\ell. \quad (2.60)$$

Pick any smooth function h on N and consider its coordinate representation $H := h \circ \psi^{-1}$. Then the coordinate representation of the function $g := h \circ f$ is

$$G = g \circ \varphi^{-1} = h \circ f \circ \varphi^{-1} = h \circ \psi^{-1} \circ \psi \circ f \circ \varphi^{-1} = H \circ F.$$

In other words,

$$G(x) = H(F_1(x), \dots, F_\ell(x)),$$

where $F = (F_1, \dots, F_\ell)$. Hence, we compute:

$$\partial_i^x g = \frac{\partial G}{\partial x_i} = \sum_{j=1}^{\ell} \frac{\partial H}{\partial y_j} \frac{\partial F_j}{\partial x_i} = \sum_{j=1}^{\ell} \frac{\partial F_j}{\partial x_i} \partial_j^y h,$$

where all partial derivatives are computed at the origin, however we suppressed this in the notations to keep those simple. In other words

$$f_*(m) \partial_i^x = \sum_{j=1}^{\ell} \frac{\partial F_j}{\partial x_i} \partial_j^y \quad \Longleftrightarrow \quad f_*(m) \partial^x = \partial^y \cdot DF,$$

where in the last equation $\partial^x = (\partial_1^x, \dots, \partial_k^x)$ is interpreted as a k -tuple of vectors in $T_m M$ and, similarly, $\partial^y = (\partial_1^y, \dots, \partial_\ell^y)$; Moreover,

$$DF = \begin{pmatrix} \frac{\partial F_1}{\partial x_1} & \cdots & \frac{\partial F_1}{\partial x_k} \\ \vdots & \ddots & \vdots \\ \frac{\partial F_\ell}{\partial x_1} & \cdots & \frac{\partial F_\ell}{\partial x_k} \end{pmatrix}$$

is the Jacobi matrix of F (evaluated at the origin). Thus, our computation shows that the following result holds.

Proposition 2.61. *Let $f: M^k \rightarrow N^\ell$ be a smooth map between two smooth manifolds. Pick charts $(U, \varphi = (x_1, \dots, x_k))$ and $(V, \psi = (y_1, \dots, y_\ell))$ centered at m and $f(m)$ respectively. Denote by F the coordinate representation of f , i.e., $F = \psi \circ f \circ \varphi^{-1}$. Then the matrix of $f_*(m)$ with respect to Bases (2.59) is given by the Jacobi matrix of F . \square*

We finish this section by the following result.

Proposition 2.62. *For any two smooth maps $f: M \rightarrow N$ and $g: N \rightarrow K$ between smooth manifolds, we have*

$$(g \circ f)_*(m) = g_*(f(m)) \circ f_*(m).$$

Proof. The proof follows directly from the definition of the differential. Indeed, for any smooth curve γ through m , we have

$$(g \circ f)_*[\gamma] = [(g \circ f) \circ \gamma] = [g \circ (f \circ \gamma)] = g_*(f(m)) [f \circ \gamma] = g_*(f(m)) (f_*(m)[\gamma]).$$

This immediately implies the statement of this proposition. \square

Exercise 2.63. Let $A: \mathbb{R}^k \rightarrow \mathbb{R}^k$ be a linear map. Show that the differential of A at any point can be identified with A itself. More precisely, this means that the diagram

$$\begin{array}{ccc} \mathbb{R}^k & \xrightarrow{A_*(x)} & \mathbb{R}^k \\ \Psi \downarrow & & \downarrow \Psi \\ \mathbb{R}^k & \xrightarrow{A} & \mathbb{R}^k \end{array}$$

commutes for any $x \in \mathbb{R}^k$, where Ψ denotes the canonical isomorphism $T_x \mathbb{R}^k \rightarrow \mathbb{R}^k$ given by (2.48), or, equivalently, by (2.49). \square

L4

Chapter 3

Submanifolds and partitions of unity

3.1 Submanifolds

Represent $\mathbb{R}^{k+\ell}$ as a product:

$$\mathbb{R}^{k+\ell} = \mathbb{R}^k \times \mathbb{R}^\ell.$$

Corresponding to this representation, we have the following maps:

$$\begin{aligned} \iota_2: \mathbb{R}^\ell &\rightarrow \mathbb{R}^{k+\ell}, & \iota_2(y) &= (0, y), \\ \pi_2: \mathbb{R}^{k+\ell} &\rightarrow \mathbb{R}^\ell, & \pi_2(x, y) &= y, \end{aligned}$$

where $x \in \mathbb{R}^k$ and $y \in \mathbb{R}^\ell$.

Let $f: \mathbb{R}^{k+\ell} \rightarrow \mathbb{R}^\ell$ be a smooth map, which is defined on some neighbourhood U of the origin. For any point $p_0 = (x_0, y_0) \in U$ we have the linear map

$$D_y f(p_0): \mathbb{R}^\ell \xrightarrow{\iota_2} \mathbb{R}^{k+\ell} \xrightarrow{D_{p_0} f} \mathbb{R}^\ell. \quad (3.1)$$

For example, if $k = \ell = 1$, we have $D_y f(p_0) = \frac{\partial f}{\partial y}(p_0)$. For this reason, we call (3.1) *the partial derivative of f with respect to y (at the point p_0)*.

To simplify the notations it is convenient to assume that p_0 is the origin and $f(0) = 0$, although this is immaterial.

Theorem 3.2. *If $D_y f(0)$ is an isomorphism, then there exists a smooth map $\theta: \mathbb{R}^{k+\ell} \rightarrow \mathbb{R}^{k+\ell}$, which is a local diffeomorphism at 0, such that $\theta(0) = 0$ and*

$$f \circ \theta = \pi_2$$

holds in a neighbourhood of the origin.

Proof. Define

$$g: \mathbb{R}^{k+\ell} \rightarrow \mathbb{R}^{k+\ell} \quad \text{by} \quad g(x, y) := (x, f(x, y)). \quad (3.3)$$

Then for the differential of g we have

$$Dg(0) = \begin{pmatrix} id_{\mathbb{R}^k} & 0 \\ D_x f(0) & D_y f(0) \end{pmatrix} \iff Dg(0) \begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} u \\ D_x f(0)u + D_y f(0)v \end{pmatrix},$$

where $u \in \mathbb{R}^k$ and $v \in \mathbb{R}^\ell$.

If $(u, v) \in \ker Dg(0)$, then $u = 0$ and $D_y f(0)v = 0$. However, $D_y f$ is an isomorphism by assumption of this theorem, so that $v = 0$. Therefore, $Dg(0)$ is injective and, hence, an isomorphism.

By [Theorem 2.24](#), there is a local inverse $\theta: \mathbb{R}^{k+\ell} \rightarrow \mathbb{R}^{k+\ell}$ to g , that is in a neighbourhood of the origin we have

$$g \circ \theta = \text{id}_{\mathbb{R}^{k+\ell}} \implies \pi_2 = \pi_2 \circ \text{id}_{\mathbb{R}^{k+\ell}} = \pi_2 \circ g \circ \theta = f \circ \theta.$$

Thus, the theorem is proved. \square

Corollary 3.4 (The implicit function theorem). *Suppose that the assumptions of [Theorem 3.2](#) hold. Then there exists a neighbourhood V_1 of $0 \in \mathbb{R}^k$, a neighbourhood V_2 of $0 \in \mathbb{R}^\ell$, and a unique smooth map $h: V_1 \rightarrow V_2$ such that*

$$f(x, y) = 0 \iff y = h(x) \quad (3.5)$$

whenever $(x, y) \in V_1 \times V_2$.

Furthermore, denoting $W := f^{-1}(0) \cap V_1 \times V_2$, the map

$$\psi := \pi_1|_W: W \rightarrow V_1, \quad (x, y) \mapsto x$$

is a homeomorphism, that is (W, ψ) is a chart on $f^{-1}(0) \cap U$.

Proof. Let $\theta: U \rightarrow \theta(U)$ be the local diffeomorphism provided by [Theorem 3.2](#). Pick any open subsets V_1 and V_2 as in the formulation of the theorem such that $V_1 \times V_2 \subset U$. For $x \in V_1$ define $h(x) := \pi_2 \circ \theta(x, 0)$. Furthermore, for $(x, y) \in V_1 \times V_2$, denote

$$(z, w) := \theta^{-1}(x, y) = g(x, y) = (x, f(x, y)).$$

Here we used the fact, that g , which is given by [\(3.3\)](#), is the inverse of θ . Then

$$\begin{aligned} f(x, y) = 0 &\implies 0 = f \circ \theta \circ \theta^{-1}(x, y) = f \circ \theta(z, w) = w \\ &\implies (z, 0) = (x, f(x, y)). \end{aligned}$$

Hence, $z = x$ and $(x, y) = \theta(x, 0)$, which yields in turn $y = h(x)$.

Furthermore, for any $x \in V_1$ we have

$$(x, 0) = g \circ \theta(x, 0) = g(\pi_1 \circ \theta(x, 0), \pi_2 \circ \theta(x, 0))$$

From the definition of g we obtain $x = \pi_1 \circ \theta(x, 0)$ and, hence, $0 = f(x, h(x))$.

To show the uniqueness, notice that

$$\begin{aligned} f(x, h(x)) = 0 &\implies g(x, h(x)) = (x, f(x, h(x))) = (x, 0) \\ f(x, \hat{h}(x)) = 0 &\implies g(x, \hat{h}(x)) = (x, 0). \end{aligned}$$

Since g is a local diffeomorphism, we obtain $h(x) = \hat{h}(x)$ provided x is sufficiently close to the origin.

Furthermore, notice that the map

$$V_1 \rightarrow W, \quad x \mapsto (x, h(x))$$

is a continuous inverse of ψ . Hence, ψ is a homeomorphism. \square

Remark 3.6. The hypothesis of **Corollary 3.4** implies that the differential of f at the origin is *surjective*. In fact, the surjectivity of the differential is decisive in **Theorem 3.2** and **Corollary 3.4**, whereas the hypothesis that $D_y f(0)$ is an isomorphism can be achieved by a linear change of coordinates, see the proof of **Theorem 3.10** below for some details.

The proofs of **Theorem 3.2** and **Corollary 3.4** show that there is a chart (U, φ) on $\mathbb{R}^{k+\ell}$ such that $\varphi(U \cap N) \subset \mathbb{R}^k \times \{0\}$. This motivates the following.

Definition 3.7. Let N be a manifold of dimension $k + \ell$. A subset $M \subset N$ is said to be a *submanifold* of dimension k (or *k-submanifold*), if for each point $m \in M$ there exists a chart (U, φ) on N centered at m such that

$$\varphi(U \cap N) = \varphi(U) \cap (\mathbb{R}^k \times \{0\}) \quad (3.8)$$

holds. Under these circumstances, the chart (U, φ) is said to be *adapted to M* .

Notice that if (U, φ) is an adapted chart, then $(M \cap U, \psi)$ is a chart on M , where

$$\psi := \pi_1 \circ \varphi|_{U \cap M} : U \cap M \rightarrow \mathbb{R}^k.$$

Proposition 3.9. *A k -submanifold is a smooth k -manifold.*

Proof. By its very definition, a k -submanifold is equipped with a C^0 -atlas \mathcal{U} , consisting of restrictions of all adapted charts.

I claim that this atlas is in fact smooth. Indeed, let (U_1, φ_1) and (U_2, φ_2) be two charts adapted to M . Denoting by $\iota_1 : \mathbb{R}^k \rightarrow \mathbb{R}^{k+\ell}$ the inclusion $\iota_1(x) = (x, 0)$, we have

$$\psi_1 \circ \psi_2^{-1}(x) = \psi_1(\varphi_2^{-1}(x, 0)) = \pi_1 \circ \varphi_1 \circ \varphi_2^{-1} \circ \iota_1(x) = \pi_1 \circ \theta_{12} \circ \iota_1(x).$$

Thus, \mathcal{U} is smooth. □

We are now in the position to state one of the central theorems of this chapter.

Theorem 3.10. *Let M and N be smooth manifolds. If n is a regular value of a smooth map $f : M \rightarrow N$ and $\dim M \geq \dim N$, then $f^{-1}(n)$ is a submanifold of M of dimension $k := \dim M - \dim N$.*

Proof. Denote

$$\ell = \dim N \quad \implies \quad \dim M = k + \ell.$$

Pick any $m \in f^{-1}(n)$ and any charts (U, φ) and (V, ψ) centered at m and n respectively. Let $F = \psi \circ f \circ \varphi^{-1}$ be the coordinate representation of f with respect to the charts (U, φ) and (V, ψ) . Since φ and ψ are diffeomorphisms, we obtain that the differential F_* of F is surjective at the origin (in fact, at any point from $F^{-1}(0)$). In particular, $\dim \ker F_*(0) = k$.

Choose a basis $(v_1, \dots, v_{k+\ell})$ of $\mathbb{R}^{k+\ell}$ such that (v_1, \dots, v_k) is a basis of $\ker F_*(0)$. Set

$$A : \mathbb{R}^{k+\ell} \rightarrow \mathbb{R}^{k+\ell}, \quad z \mapsto \sum_{j=1}^{k+\ell} z_j v_j.$$

Notice that by the definition of A and elementary facts from linear algebra, the following holds:

- A is an isomorphism;
- $A \circ \iota_1 : \mathbb{R}^k \rightarrow \ker F_*(0)$ is an isomorphism;

- $F_*(0) \circ A \circ \iota_2: \mathbb{R}^\ell \rightarrow \mathbb{R}^\ell$ is an isomorphism.

Furthermore, consider the map $G := F \circ A: \mathbb{R}^{k+\ell} \rightarrow \mathbb{R}^\ell$. By Exercise 2.63, we have

$$G_*(0) = F_*(0) \circ A \quad \implies \quad D_y G = F_*(0) \circ A \circ \iota_2.$$

Since the latter map is an isomorphism, by the proofs of Theorem 3.2 and Corollary 3.4 we obtain a chart (W, ξ) on $\mathbb{R}^{k+\ell}$ adapted to $G^{-1}(0)$, that is

$$\xi(W \cap G^{-1}(0)) = \xi(W) \cap (\mathbb{R}^k \times \{0\}).$$

Without loss of generality we can assume that W is contained in $A^{-1}(\varphi(U))$.

Various charts involved in the proof are shown schematically on Figure 3.1.




Figure 3.1: Scheme of the proof of Theorem 3.10.

Define a chart $(\hat{W}, \hat{\xi})$ on $\mathbb{R}^{k+\ell}$ by

$$(\hat{W}, \hat{\xi}) = (A^{-1}(W), \xi \circ A^{-1}).$$

Since $z \in G^{-1}(0) \Leftrightarrow Az \in F^{-1}(0)$, we obtain

$$\hat{\xi}(\hat{W} \cap F^{-1}(0)) = \xi(W \cap G^{-1}(0)) = \xi(W) \cap (\mathbb{R}^k \times \{0\}).$$

Finally, setting

$$\varphi_1 := \hat{\xi} \circ \varphi \quad \text{and} \quad U_1 = \varphi_1^{-1}(\hat{\xi}(\hat{W})) = \varphi^{-1}(\hat{W})$$

we obtain

$$\varphi_1(U_1 \cap f^{-1}(n)) = \hat{\xi}(\hat{W} \cap F^{-1}(0)) = \xi(W) \cap (\mathbb{R}^k \times \{0\}) = \varphi_1(U_1) \cap (\mathbb{R}^k \times \{0\}).$$

Thus, (U_1, φ_1) is a chart adapted to $f^{-1}(n)$ at m . \square

Notice the following: If $\dim M < \dim N$, then n is a regular value of smooth map $f: M \rightarrow N$ if and only if $n \notin \text{Im } f$, see the paragraph following Definition 2.28. In this case $f^{-1}(n) = \emptyset$ is also (by definition) a smooth manifold. Thus, the condition $\dim M \geq \dim N$ can be dropped in the formulation of Theorem 3.10.

Proposition 3.11. *In the setting of Theorem 3.10, for any $m \in f^{-1}(n)$ we have*

$$T_m f^{-1}(n) = \ker f_*(m).$$

Proof. Pick any curve γ in $f^{-1}(n)$ through m . Since γ lies in the level set of f , we have

$$f \circ \gamma(t) = n \quad \text{for all } t \in (-\varepsilon, \varepsilon). \quad (3.12)$$

Since the constant curve $t \mapsto n$ represents the zero vector in $T_n N$, by the definition of the differential of f and (3.12) we obtain $f_*(m)([\gamma]) = 0$. In other words any vector $[\gamma]$ tangent to $f^{-1}(n)$ lies in the kernel of $f_*(m)$. \square

Example 3.13.

- (i) Consider the map $f: \mathbb{R}^{n+1} \rightarrow \mathbb{R}$, $f(x) = |x|^2$. Then 1 is a regular value of f . In particular, $S^n = f^{-1}(1)$ is a manifold of dimension n . Of course, the reader knows this fact by now very well.
- (ii) Let $M_n(\mathbb{R})$ be the space of all $n \times n$ matrices with real entries. One can show that 1 is a regular value of the function $\det: M_n(\mathbb{R}) \rightarrow \mathbb{R}$, $A \mapsto \det A$. Consequently,

$$\text{SL}_n(\mathbb{R}) := \{A \in M_n(\mathbb{R}) \mid \det A = 1\}$$

is a manifold of dimension $\dim M_n(\mathbb{R}) - 1 = n^2 - 1$.

Let us compute the tangent space to $\text{SL}_n(\mathbb{R})$ at the point $\mathbb{1}$. To this end, it is convenient to identify $M_n(\mathbb{R})$ with \mathbb{R}^{n^2} . Recalling that

$$\det A = \sum_{\sigma} \text{sign } \sigma a_{1\sigma(1)} \cdots a_{n\sigma(n)},$$

where σ runs through all permutations of the set $\{1, \dots, n\}$, for any $B \in M_n(\mathbb{R})$ we obtain

$$\begin{aligned} \det(\mathbb{1} + tB) &= (1 + tb_{11})(1 + tb_{22}) \cdots (1 + tb_{nn}) \\ &\quad + \sum_{\sigma \neq \text{id}} \text{sign } \sigma (\delta_{1\sigma(1)} + tb_{1\sigma(1)}) (\delta_{2\sigma(2)} + tb_{2\sigma(2)}) \cdots (\delta_{n\sigma(n)} + tb_{n\sigma(n)}). \end{aligned}$$

Notice that for any $\sigma \neq id$, $\sigma(i) \neq i$ at least for two values of i . Hence, the last term in the above expression is $o(t)$. This yields

$$\det(\mathbb{1} + tB) = (1 + t \operatorname{tr} B + o(t)) + o(t).$$

Consequently, $\det_*(\mathbb{1}) B = \operatorname{tr} B$ and therefore

$$T_{\mathbb{1}} \operatorname{SL}_n(\mathbb{R}) = \{B \in M_n(\mathbb{R}) \mid \operatorname{tr} B = 0\}.$$

(iii) Let $\operatorname{Sym}^n(\mathbb{R}) \subset M_n(\mathbb{R})$ denote the subspace of all symmetric matrices. One can show that the identity matrix $\mathbb{1} \in \operatorname{Sym}^n(\mathbb{R})$ is a regular value of the map

$$f: M_n(\mathbb{R}) \rightarrow \operatorname{Sym}^n(\mathbb{R}), \quad f(A) = A \cdot A^t. \quad (3.14)$$

Consequently,

$$O(n) := \{A \in M_n(\mathbb{R}) \mid A \cdot A^t = \mathbb{1}\}$$

is a manifold and

$$\begin{aligned} \dim O(n) &= \dim M_n(\mathbb{R}) - \dim \operatorname{Sym}^n(\mathbb{R}) = n^2 - \frac{n(n+1)}{2} \\ &= \frac{n(n-1)}{2}. \end{aligned}$$

Notice that if we would consider (3.14) as a map $M_n(\mathbb{R}) \rightarrow M_n(\mathbb{R})$, then $\mathbb{1}$ would not be a regular value.

Just like in the case of $\operatorname{SL}_n(\mathbb{R})$, let us compute the tangent space to $O(n)$ at the point $\mathbb{1}$. We have

$$f(\mathbb{1} + sB) = (\mathbb{1} + sB) \cdot (\mathbb{1} + sB)^t = \mathbb{1} + s(B + B^t) + o(s).$$

Hence, $f_*(\mathbb{1})B = B + B^t$ and

$$T_{\mathbb{1}} O(n) = \{B \in M_n(\mathbb{R}) \mid B^t = -B\}.$$

We finish this section by Sard's theorem, which, loosely speaking, says that for any smooth map almost any point is a regular value. More precisely, we say that a subset A of a smooth k -manifold M is of measure zero, if for any chart (U, φ) on M the set $\varphi(A \cap U) \subset \mathbb{R}^k$ is of measure zero.

Theorem 3.15 (Sard). *Let $f: M \rightarrow N$ be a smooth map between smooth manifolds. Then almost any point $n \in N$ is a regular value of f , that is the set of critical values for f is of measure zero.* \square

A proof of Sard's theorem can be found for example in [BT03, 9.4] or [Mil65, §3].

L5

3.2 Immersions and embeddings

Just like maps with surjective differentials can be conveniently described as projections after applying a diffeomorphism, the maps with injective differentials admit an analogous description.

Theorem 3.16. *Let U be an open subset of \mathbb{R}^k containing the origin and $f: U \rightarrow \mathbb{R}^k \times \mathbb{R}^\ell$ be a smooth map such that $f(0) = 0$ and*

$$f_{1*}(0): \mathbb{R}^k \rightarrow \mathbb{R}^k, \quad \text{where } f_1 := \pi_1 \circ f,$$

is an isomorphism. Then there exists a neighbourhood $V \subset \mathbb{R}^{k+\ell}$ of the origin and a diffeomorphism $\theta: V \rightarrow \theta(V) \subset \mathbb{R}^{k+\ell}$ such that $\theta \circ f = \iota_1$ and $\theta(V \cap f(U)) = \theta(V) \cap (\mathbb{R}^k \times \{0\})$.

Proof. The proof of this theorem is similar to the proof of [Theorem 3.2](#).

Thus, consider the map

$$F: U \times \mathbb{R}^\ell \rightarrow \mathbb{R}^{k+\ell} = \mathbb{R}^k \times \mathbb{R}^\ell, \quad F(x, y) := f(x) + (0, y) = (f_1(x), f_2(x) + y).$$

The differential of this map

$$F_*(0) = \begin{pmatrix} f_{1*}(0) & 0 \\ f_{2*}(0) & \text{id}_{\mathbb{R}^\ell} \end{pmatrix}$$

is an isomorphism. Hence, there exists a neighbourhood V of the origin and a diffeomorphism $\theta: V \rightarrow \theta(V)$ such that

$$\theta \circ F = \text{id}_{\theta(V)}.$$

In particular, for any $(x, 0) \in \theta(V)$ the above equality yields:

$$\theta \circ F(x, 0) = \theta \circ f(x) = \iota_1(x) \quad \implies \quad \theta \circ f = \iota_1.$$

Hence, $\theta(V) \cap (\mathbb{R}^k \times \{0\}) \subset \theta(V \cap f(U))$.

To show the converse inclusion, let $(x, y) \in \theta(V \cap f(U))$. Hence, there exists some $(z, w) \in V \cap f(U)$ such that $(x, y) = \theta(z, w)$. In this case we must have $(z, w) = f(x)$ for some $x \in U$ and therefore

$$(x, y) = \theta(z, w) = \theta \circ f(x) = (x, 0).$$

Thus, $y = 0$ and $(x, 0) \in V$, which yields $\theta(V \cap f(U)) \subset \theta(V) \cap (\mathbb{R}^k \times \{0\})$. □

Definition 3.17. A smooth map $f: M^k \rightarrow N^\ell$ such that $f_*(m)$ is injective at each point $m \in M$ is called an *immersion*. An immersion, which is a diffeomorphism onto a k -submanifold of N , is called an *embedding*.

Clearly, an immersion of $f: M \rightarrow N$ can exist only if $\dim M \leq \dim N$. Notice also, that by [Theorem 3.16](#) each immersion is locally injective, however an immersion does not need to be globally injective. Even if an immersion is injective, this may fail to be an embedding. This is shown schematically on Figures [3.2](#) and [3.3](#) below. In particular, the image of an immersion does not need to be a submanifold.




Figure 3.2: The image of a non-injective immersion $\mathbb{R} \rightarrow \mathbb{R}^2$.




Figure 3.3: The image of an injective immersion $\mathbb{R} \rightarrow \mathbb{R}^2$, which is not an embedding.

Proposition 3.18. *An immersion which is a homeomorphism onto its image is an embedding.*

Proof. Denote $k := \dim M$ and $\ell := \dim N$. The proof consists of the following 3 steps.

Step 1. *For any $m \in M$ there exists a chart (V, ψ) on N centered at $n := f(m)$ with the following properties:*

- $\psi_{1*}(n)|_{\text{Im } f_*(m)}: \text{Im } f_*(m) \rightarrow \mathbb{R}^k$ is an isomorphism, where $\psi_1 = \pi_1 \circ \psi$ and $\pi_1: \mathbb{R}^\ell = \mathbb{R}^k \oplus \mathbb{R}^{\ell-k} \rightarrow \mathbb{R}^k$ is the projection.
- There exists a neighbourhood U of m such that $f(U) = V \cap f(M)$.

Since f is a homeomorphism onto its image, $f: M \rightarrow f(M)$ is an open map. In particular, for any open $\hat{U} \subset M$ there exists an open subset $\hat{V} \subset N$ such that $f(\hat{U}) = \hat{V} \cap f(M)$. If \hat{U} is a neighbourhood of m , we can choose a chart (V, ξ) centered at n such that $V \subset \hat{V}$.

Furthermore, since $\xi_*: T_n N \rightarrow \mathbb{R}^\ell$ is an isomorphism and $\text{Im } f_*(m)$ is a k -dimensional subspace of $T_n N$, we can find a linear isomorphism $A: \mathbb{R}^\ell \rightarrow \mathbb{R}^\ell$ such that

$$A(\xi_*(\text{Im } f_*(n))) = \mathbb{R}^k \times \{0\}.$$

Then $(V, \psi) = (V, A \circ \xi)$ is the required chart. Also, setting $U := f^{-1}(V)$ we obtain $f(U) = V \cap f(M)$.

Step 2. *$f(M)$ is a submanifold of N .*

Pick any $m \in M$ and a chart (U, φ) centered at m . Pick also a chart (V, ψ) as in the previous step. Denote also $W := \psi(V) \subset \mathbb{R}^\ell$.

Let $F = \psi \circ f \circ \varphi^{-1}$ be the coordinate representation of f . Denoting $F_1 := \pi_1 \circ F: \mathbb{R}^k \rightarrow \mathbb{R}^k$, we have

$$F_{1*}(0) = \pi_{1*}(0) \circ F_*(0) = \pi_{1*}(0) \circ \psi_*(n) \circ f_*(m) \circ \varphi_*^{-1}(0) = \psi_{1*}(0) \circ f_*(m) \circ \varphi_*^{-1}(0)$$

Since $\varphi_*^{-1}(0)$ is an isomorphism, by Step 1 we obtain that $F_{1*}(0)$ is injective. Hence, $F_{1*}(0)$ is an isomorphism. Hence, by Theorem 3.16 we can find a diffeomorphism¹ $\theta: W \rightarrow \theta(W) \subset \mathbb{R}^\ell$ such that

$$\theta \circ F = \iota_1 \iff (\theta \circ \psi) \circ f \circ \varphi^{-1} = \iota_1$$

¹Without loss of generality we may assume that V was chosen so that θ is defined everywhere on W .

Denote $\hat{\psi} := \theta \circ \psi$. Then $(W, \hat{\psi})$ is a chart on N adapted to $f(M)$.

Step 3. f is a diffeomorphism between M and $f(M)$.

Let (U, φ) and $(W, \hat{\psi})$ be as in the preceding step. By the construction of $\hat{\psi}$, the coordinate representation of f is $\hat{\psi} \circ f \circ \varphi^{-1} = \iota_1: \mathbb{R}^k \rightarrow \mathbb{R}^\ell$. Since the restriction of $\pi_1 \circ \psi$ to $f(M) \cap W$ is a chart on $f(M)$, the coordinate representation of f viewed as a map $f: M \rightarrow f(M)$ is given by

$$\pi_1 \circ \hat{\psi} \circ f \circ \varphi^{-1} = \pi_1 \circ \iota_1 = \text{id}.$$

Hence, f is a local diffeomorphism. Since $f: M \rightarrow f(M)$ is bijective, this is a diffeomorphism. \square

Corollary 3.19. *If M is compact, then any injective immersion $f: M \rightarrow N$ is an embedding.*

Proof. Pick a closed subset $A \subset M$. Since A is closed in M , A is compact and therefore $f(A)$ is compact in N . Since N is Hausdorff, $f(A)$ is closed. Hence, f is a closed map, i.e., the image of any closed subset is closed. This means that $f^{-1}: f(M) \rightarrow M$ is continuous, that is, $f: M \rightarrow f(M)$ is a homeomorphism. The statement of this corollary now follows immediately from **Proposition 3.18**. \square

Theorem 3.10 combined with Sard's theorem allows us to construct many smooth manifolds, which are in fact submanifolds of Euclidean spaces. It turns out that any smooth manifold can be realized as a submanifold of an Euclidean space.

Theorem 3.20 (Whitney's embedding theorem). *For any smooth manifold M there is an embedding of M into \mathbb{R}^n for some $n \in \mathbb{N}$.*

Proof. We prove Whitney's embedding theorem only in the case when M is compact.

For any $m \in M$ choose a chart (U_m, φ_m) . Pick also open neighbourhoods $V_m \subset W_m$ and a bump function ρ_m such that the following holds:

- $\overline{V_m} \subset U_m$;
- $\rho_m|_{\overline{W_m}} \equiv 1$ and $\rho_m < 1$ outside of $\overline{W_m}$;
- ρ_m vanishes outside of V_m .

Since M is compact, there is a finite subset $\{m_1, \dots, m_p\}$ of M such that $\{W_i\}$ cover all of M , where $W_i := W_{m_i}$. Consider each $\psi_i := \rho_i \cdot \psi_i := \rho_{m_i} \cdot \varphi_{m_i}$ as a smooth map $M \rightarrow \mathbb{R}^k$ (extended by zero outside of V_{m_i}), where $k = \dim M$. Finally, define

$$f: M \rightarrow \mathbb{R}^{pk+p}, \quad f(m) = (\psi_1(m), \dots, \psi_p(m), \rho_1(m), \dots, \rho_p(m)).$$

Clearly, f is smooth. I claim that this map is also injective. Indeed, pick any two distinct points m and \hat{m} . Without loss of generality, we can assume $m \in W_1$. If $\hat{m} \in \overline{W_1}$, then $\psi_1(m) = \varphi_1(m) \neq \varphi_1(\hat{m}) = \psi_1(\hat{m})$. If $\hat{m} \notin \overline{W_1}$, then $1 = \rho_1(m) \neq \rho_1(\hat{m})$, so that f is injective indeed.

Furthermore, assuming $m \in W_1$ again, $\psi_{1*}(m): T_m M \rightarrow \mathbb{R}^k$ is an isomorphism. Hence, $f_*(m): T_m M \rightarrow \mathbb{R}^{kp+p}$ is injective at any $m \in M$. By **Corollary 3.19**, f is an embedding. \square

Whitney's embedding theorem shows that any (abstract) manifold M can be thought of as a submanifold of an Euclidean space. In other words, we could have defined manifolds as the subspaces of Euclidean spaces admitting charts² at each point. Some authors do take this point of view arguing that this yields the same pool of examples. While it is true of course that this would yield the same pool of examples, manifolds often do not arise as subsets of Euclidean spaces. For example, the real projective space is not obviously contained in any Euclidean space (and it is even not so obvious how to construct an embedding). Even if one decides to work with the submanifolds only, one finds out pretty soon that certain useful constructions, for example taking quotients by group actions, are incompatible with this setting. More importantly, it is useful to distinguish "inner" properties of manifolds from those of an embedding. All these reasons led to the necessity to separate abstract manifolds from their particular realizations as submanifolds.

3.3 Partitions of unity

Let $f: M \rightarrow \mathbb{R}$ be a (continuous) map.

Definition 3.21. The set

$$\text{supp } f := \overline{\{m \in M \mid f(m) \neq 0\}}$$

is called *the support* of f . In other words, $\text{supp } f$ is the closure of the set, where f does not vanish.

Example 3.22.

- (i) For $f: \mathbb{R} \rightarrow \mathbb{R}$, $f(x) = x$ we have $\text{supp } f = \mathbb{R}$.
- (ii) For the cut off function (2.53) we have $\text{supp } \chi_r = (-\infty, 1 + r]$.
- (iii) For the bump function $\hat{\rho}: \mathbb{R}^k \rightarrow \mathbb{R}$, $\hat{\rho}(x) = \chi_1(|x|)$ we have

$$\text{supp } \hat{\rho} = \{x \in \mathbb{R}^k \mid |x| \leq 2\}.$$

Definition 3.23. A family of functions $\{\rho_\alpha: M \rightarrow \mathbb{R}_{\geq 0} \mid \alpha \in A\}$, where A is an index set, is called a *partition of unity*, if the following holds:

- (i) The family $\{\text{supp } \rho_\alpha \mid \alpha \in A\}$ is locally finite, that is for each $m \in M$ there exists a neighbourhood $W \ni m$ such that the set $\{\alpha \in A \mid W \cap \text{supp } \rho_\alpha \neq \emptyset\}$ is finite;
- (ii) For each $m \in M$ we have

$$\sum_{\alpha \in A} \rho_\alpha(m) = 1. \quad (3.24)$$

Furthermore, a partition of unity $\{\rho_\alpha\}$ is said to be *subordinate to a covering* $\{U_\beta \mid \beta \in B\}$, if for each $\alpha \in A$ there exists some $\beta = \beta(\alpha)$ such that $\text{supp } \rho_\alpha \subset U_{\beta(\alpha)}$.

Notice that (i) implies that for each $m \in M$ the set $\{\alpha \mid \rho_\alpha(m) \neq 0\}$ is finite so that (3.24) is a finite sum.

Any manifold trivially admits a partition of unity consisting of a single constant function. To obtain a non-trivial example, consider $M = \mathbb{R}$ and the family $\{\hat{\rho}_j \mid j \in \mathbb{Z}\}$, where

$$\hat{\rho}_j: \mathbb{R} \rightarrow \mathbb{R}, \quad \hat{\rho}_j(x) = \chi_1(|x - j|). \quad (3.25)$$

²More precisely, admitting adapted charts.

In particular, $\hat{\rho}_j$ equals 1 everywhere on the interval $[j - 1, j + 1]$ and $\text{supp } \hat{\rho}_j = [j - 2, j + 2]$. Hence, the function

$$\hat{\rho}(x) := \sum_{j \in \mathbb{Z}} \hat{\rho}_j(x)$$

is well-defined and positive everywhere on \mathbb{R} . Therefore, $\{\rho_j := \hat{\rho}_j / \hat{\rho} \mid j \in \mathbb{Z}\}$ is a partition of unity on \mathbb{R} . This partition of unity is subordinate for example to the following coverings:

$$\{(j - 3, j + 3) \mid j \in \mathbb{Z}\} \quad \text{and} \quad \{(-j, j) \mid j \in \mathbb{N}\}.$$

L 6

Theorem 3.26 (Existence of Partition of Unity). *Given an open covering \mathcal{U} on a manifold, there is a partition of unity subordinate to \mathcal{U} .*

Proof. We prove this theorem only under a simplifying hypothesis that the manifold M under consideration is compact.

Thus, pick any point $m \in M$ and a set $U_{\beta(m)} \in \mathcal{U}$ containing m . By [Proposition 2.54](#), there exist a neighbourhood $V_m \subset U_{\beta(m)}$ and a bump function $\hat{\rho}_m$ such that $\hat{\rho}_m \equiv 1$ on V_m and $\text{supp } \hat{\rho}_m \subset U_{\beta(m)}$.

Since M is compact, we can choose a finite subset $\{m_1, \dots, m_p\}$ such that $\{V_1, \dots, V_p\}$ is a covering of M , where $V_j := V_{m_j}$. Redenoting $\hat{\rho}_j := \hat{\rho}_{m_j}$, we obtain that

$$\hat{\rho}(m) := \sum_{j=1}^p \hat{\rho}_j(m)$$

is positive everywhere on M . Hence, $\{\rho_j := \hat{\rho}_j / \hat{\rho} \mid j = \overline{1, p}\}$ is a partition of unity on M . Moreover, this is subordinate to \mathcal{U} , since $\text{supp } \rho_j = \text{supp } \hat{\rho}_j \subset U_{\beta(m_j)}$. \square

A couple of remarks are in place here. First, a proof of the above theorem in full generality can be found for example in [\[War83, 1.11\]](#) and uses the axiom of second countability, which we have not really used so far. This is one of the main reasons that the manifolds are required to be second countable.

Second, it is straightforward to generalize the definition of a smooth manifold to the complex holomorphic setting. Namely, we could have defined a complex manifold as a (Hausdorff second countable) topological space equipped with an atlas $\mathcal{U} := \{(U_\alpha, \varphi_\alpha) \mid \alpha \in A\}$, where $\varphi_\alpha: U_\alpha \rightarrow \varphi_\alpha(U) \subset \mathbb{C}^k$ is a homeomorphism, such that all transition maps

$$\theta_{\alpha\beta} := \varphi_\alpha \circ \varphi_\beta^{-1}: \mathbb{C}^k \rightarrow \mathbb{C}^k$$

are holomorphic. Most of the results we have seen so far are still valid in this setting (with an obvious replacement of the adjective “smooth” by “holomorphic”), except, most notably, [Theorem 3.26](#) (and related [Theorem 3.20](#) and [Proposition 2.54](#)). The existence of the partition of unity on smooth manifolds and its non-existence on complex manifolds gives these two theories somewhat different flavours.

A typical application of the partition of unity is to existence questions, which we illustrate on the following result.

Theorem 3.27. *Let $M^k \subset N^\ell$ be an embedded submanifold. Then for any $h \in C^\infty(N)$ the restriction of h to M is a smooth function on M . Conversely, any smooth function f on M admits a smooth extension to N , that is there exists some $h \in C^\infty(N)$ such that $h|_M = f$.*

Proof. First notice that $h|_M = h \circ \iota_M$, where $\iota_M: M \rightarrow N$ is the embedding. The smoothness of the restriction then follows from the smoothness of ι_M .

Thus, given $f \in C^\infty(M)$ we wish to find some $h \in C^\infty(N)$ such that $h|_M = f$. Let \mathcal{U} be a covering of M by adapted charts. For the covering $\mathcal{U}_N := \mathcal{U} \cup \{N \setminus M\}$ of N pick a partition of unity $\{\rho_\alpha\}$ on N adapted to \mathcal{U}_N .

For each α define $f_\alpha \in C^\infty(M)$ by

$$f_\alpha(m) := \rho_\alpha(m) \cdot f(m) \quad \implies \quad f = \sum_{\alpha} f_\alpha,$$

where at each point $m \in M$ only finitely many f_α are not vanishing.

I claim that each f_α admits an extension h_α . First recall that if (U, φ) is an adapted chart, then $(U \cap M, \psi)$ is a chart on M , where $\psi = \pi_1 \circ \varphi|_M$. Denoting by $F = f \circ \psi^{-1}$ the coordinate representation of f , define

$$H_\alpha(x, y) := F(x) \cdot (\rho_\alpha \circ \varphi^{-1})(x, y) \quad \text{and} \quad h_\alpha(n) := \begin{cases} H_\alpha \circ \varphi(n) & \text{if } n \in U, \\ 0 & \text{otherwise.} \end{cases}$$

Notice that H_α is a smooth function on \mathbb{R}^ℓ , which vanishes outside of $\varphi(U)$ so that h_α is well-defined and smooth. Moreover, the family $\{\text{supp } h_\alpha \mid \alpha \in A\}$ is locally finite, since

$$(h_\alpha(n) \neq 0 \implies \rho_\alpha(n) \neq 0) \implies \text{supp } h_\alpha \subset \text{supp } \rho_\alpha.$$

Therefore, we can define $h(n) := \sum_{\alpha} h_\alpha(n)$, which is smooth, because in a neighbourhood of each point h is a finite sum of smooth functions. For $m \in M$, we have

$$h(m) = \sum_{\alpha} h_\alpha(m) = \sum_{\alpha} f_\alpha(m) = f(m).$$

Hence, h is a smooth extension of f . □

Chapter 4

The tangent bundle and the group of diffeomorphisms

4.1 Some elements of linear algebra

In what follows it may be useful to recall the following elements of linear algebra. Let V be a linear vector space of dimension k . Any basis $v = (v_1, \dots, v_k)$ of V gives rise to the linear isomorphism

$$\mathbb{R}^k \rightarrow V, \quad y \mapsto \sum_{j=1}^k y_j v_j = v \cdot y. \quad (4.1)$$

The very last term is a shortcut for the sum of the products on the left hand side and should be understood as a matrix multiplication, where the first “matrix” v consists of 1 row and k columns (and its entries are vectors from V), whereas y is a matrix with k rows and 1 column, i.e., y is just a column-vector.

Conversely, given a linear isomorphism $\mathbb{R}^k \rightarrow V$ we can construct a basis of V just by taking the image of the standard basis of \mathbb{R}^k . This yields a bijective correspondence between the set of all bases of V and the set of all isomorphisms $\mathbb{R}^k \rightarrow V$.

Furthermore, let $w = (w_1, \dots, w_k)$ be another basis of V . Then w and v are related by the so called change-of-basis matrix B , which is obtained as follows. Decompose w_i in terms of the basis v , that is write

$$w_i = \sum_{j=1}^k b_{ij} v_j.$$

Then $B = (b_{ij})$ is the change-of-basis matrix between w and v . In terms of the matrix multiplication used above, the relation between v , w , and B can be elegantly expressed as follows:

$$w = v \cdot B.$$

Just to familiarize ourselves better with these notations, let $\varphi: V \rightarrow V$ be a linear map. The reader knows that given a basis of V , say v , φ can be represented by a $k \times k$ -matrix A . This means, that if (y_1, \dots, y_k) are coordinates of a vector $v \in V$, then the coordinates of $\varphi(v)$ are given by $A \cdot y$ (in this formula y is interpreted as a column-vector). An elementary computation yields that the j th column of A consists of coordinates of $\varphi(v_j)$ with respect to v . In other words, A can be characterized by the equality $\varphi(v) = v \cdot A$. If $w = v \cdot B$ is another basis of V as above, then we have

$$\varphi(w) = \varphi(v \cdot B) = \varphi(v) \cdot B = v \cdot A B = w \cdot B^{-1} A B.$$

This yields that the matrix of φ with respect to the basis w is $B^{-1}AB$. The reader surely knows this fact from the linear algebra, however the typical proof of this boils down to a tedious computation.

4.2 The tangent and cotangent bundles

Consider the set

$$TM := \bigsqcup_{m \in M} T_m M,$$

which comes equipped with a map

$$\pi: TM \rightarrow M, \quad \pi(v) = m \Leftrightarrow v \in T_m M.$$

For example, in the case $M = \mathbb{R}^k$ we can identify $T_m \mathbb{R}^k$ with \mathbb{R}^k just as in (2.48) so that

$$T\mathbb{R}^k = \bigsqcup_{m \in \mathbb{R}^k} T_m \mathbb{R}^k = \mathbb{R}^k \times \mathbb{R}^k \quad \text{and} \quad \pi = \pi_1.$$

Furthermore, let (U, φ) be a chart on M . Write $\varphi = (x_1, \dots, x_k)$ and recall that at each point $m \in M$ we have constructed the following basis of $T_m M$:

$$\partial_x|_m := (\partial_1, \dots, \partial_k)|_m, \quad \text{where} \quad \partial_j|_m f = \frac{\partial}{\partial x_j} \Big|_{x=\varphi(m)} (f \circ \varphi^{-1}(x)).$$

See (2.47) and the proof of Proposition 2.46 for further details. Therefore, we obtain the bijection

$$U \times \mathbb{R}^k \rightarrow \pi^{-1}(U) = \bigsqcup_{m \in U} T_m M, \quad (m, y) \mapsto \sum_{j=1}^k y_j \partial_j|_m.$$

Combining this with $\varphi: U \rightarrow \varphi(U)$, which is also a bijection, we obtain finally a bijective map

$$\tau = \tau_\varphi: \varphi(U) \times \mathbb{R}^k \rightarrow \pi^{-1}(U), \quad \tau(x, y) := \sum_{j=1}^k y_j \partial_j|_{\varphi^{-1}(x)}.$$

Proposition 4.2. *Let $\mathcal{U} := \{(U_\alpha, \varphi_\alpha) \mid \alpha \in A\}$ be a smooth atlas on M . There is a unique second countable and Hausdorff topology on M such that*

$$\mathcal{V} := \left\{ \left(\pi^{-1}(U_\alpha), \tau_\alpha^{-1} \right) \mid \alpha \in A \right\}$$

is a C^0 -atlas on TM , where $\tau_\alpha := \tau_{\varphi_\alpha}$. This atlas is in fact smooth so that TM is a smooth manifold of dimension $2k$. Moreover, π is a smooth map with surjective differential at each point.

Proof. The proof consists of the following steps.

Step 1. Write $\varphi_\alpha = (s_1, \dots, s_k)$ and $\varphi_\beta = (x_1, \dots, x_k)$. At each point $m \in U_\alpha \cap U_\beta$ we obtain two bases of $T_m M$, namely $\partial_s|_m$ and $\partial_x|_m$. Then the change-of-basis matrix between these two bases is the Jacobi-matrix of $\theta_{\alpha\beta}$, i.e.,

$$\partial_x|_m = \partial_s|_m \cdot B \quad \implies \quad B = D\theta_{\alpha\beta}. \quad (4.3)$$

Pick any function f and let $F_\alpha := f \circ \varphi_\alpha^{-1}$ be the coordinate representation of f with respect to the chart φ_α . Then we have

$$F_\beta = f \circ \varphi_\beta^{-1} = f \circ \varphi_\alpha^{-1} \circ \varphi_\alpha \circ \varphi_\beta^{-1} = F_\alpha \circ \theta_{\alpha\beta}.$$

Therefore, using the above equality we obtain

$$\partial_j^x f = \frac{\partial F_\beta}{\partial x_j} = \sum_{i=1}^k \frac{\partial F_\alpha}{\partial x_i} \frac{\partial \theta_{\alpha\beta,i}}{\partial x_j} = \sum_{i=1}^k \frac{\partial \theta_{\alpha\beta,i}}{\partial x_j} \partial_i^s f, \quad (4.4)$$

where we dropped the point where the derivatives are computed from the notations. Since f is an arbitrary function, we obtain (4.3).

Step 2. For the coordinate transformation $\Theta_{\alpha\beta} := \tau_\alpha^{-1} \circ \tau_\beta$ on TM we have

$$\Theta_{\alpha\beta}(x, y) = (\theta_{\alpha\beta}(x), \theta_{\alpha\beta*}(x) y).$$

In particular, $\Theta_{\alpha\beta}$ is smooth.

Denote $\tau_\beta(x, y) = v$. This means

$$\varphi_\beta(\pi(v)) = x \quad \text{and} \quad v = \sum_{j=1}^k y_j \partial_j^x = \partial_x \cdot y,$$

where the right hand side of this equality is interpreted in the sense of (4.1). Therefore, by (4.3) we have

$$v = \partial_x \cdot y = \partial_s \cdot D\theta_{\alpha\beta} \cdot y.$$

Denoting $\tau_\alpha^{-1}(v) = (s, t)$, we have

$$s = \varphi_\alpha(\pi(v)) = \varphi_\alpha(\varphi_\beta^{-1}(x)) = \theta_{\alpha\beta}(x) \quad \text{and} \quad t = D\theta_{\alpha\beta} \cdot y.$$

Step 3. We construct a topology on TM .

Declare a set $V \subset TM$ to be open in TM if and only if $\tau_\alpha^{-1}(V) = \tau_\alpha(V \cap U_\alpha)$ is open in \mathbb{R}^{2k} for any $\alpha \in A$. We have

- (i) \emptyset is open and $\tau_\alpha^{-1}(M) = \varphi_\alpha(U_\alpha) \times \mathbb{R}^k \implies M$ is open.
- (ii) V_1 and V_2 are open $\implies \tau_\alpha^{-1}(V_1 \cap V_2) = \tau_\alpha^{-1}(V_1) \cap \tau_\alpha^{-1}(V_2)$ is open.
- (iii) If V_β are open for each $\beta \in B$, then $\tau_\alpha^{-1}(\cup_\beta V_\beta) = \cup_\beta \tau_\alpha^{-1}(V_\beta)$ is open.

Hence, this yields a topology on TM such that π is continuous. Moreover, each $(\pi^{-1}(U_\alpha), \tau_\alpha^{-1})$ is a chart on TM .

Step 4. The topology of TM is Hausdorff and second countable.

Indeed, pick any two distinct points v_1, v_2 and consider the following cases:

- (a) $\pi(v_1) \neq \pi(v_2) \implies \pi^{-1}(U_1)$ and $\pi^{-1}(U_2)$ separate v_1 and v_2 provided $U_1 \cap U_2 = \emptyset$.
- (b) $\pi(v_1) = \pi(v_2) =: m \implies$ for any chart $(U_\alpha, \varphi_\alpha)$ containing m the sets $\tau_\alpha(U_\alpha \times V_1)$ and $\tau_\alpha(U_\alpha \times V_2)$ are open and separate v_1 and v_2 provided $V_1, V_2 \subset \mathbb{R}^k$ separate $\pi_2(\tau_\alpha^{-1}(v_1))$ and $\pi_2(\tau_\alpha^{-1}(v_2))$.

Furthermore, the constructed topology is second countable for the following reason: Let U_i be a countable basis of the topology of M and V_j be a countable basis for \mathbb{R}^k . Without loss of generality we can assume that each U_i is contained in some chart U_{α_i} . Then the collection of all sets of the form

$$\tau_{\alpha_i}(\varphi_{\alpha_i}(U_i) \times V_j)$$

is a countable basis for the topology of TM .

To finish the proof, pick a chart $(\pi^{-1}(U_\alpha), \tau_\alpha^{-1})$ and consider the coordinate representation of π with respect to this chart on TM and the chart $(U_\alpha, \varphi_\alpha)$ on M :

$$\varphi_\alpha \circ \pi \circ \tau_\alpha(x, y) = \varphi_\alpha(\varphi_\alpha^{-1}(x)) = x \quad \implies \quad \varphi_\alpha \circ \pi \circ \tau_\alpha = \pi_1.$$

Thus π is smooth and its differential is surjective at each point. \square

To get a better understanding of the tangent bundle let us consider the case of an embedded submanifold $M \subset \mathbb{R}^\ell$. If $\iota: M \rightarrow \mathbb{R}^\ell$ denotes the embedding, for each $m \in M$ the map $\iota_*(m): T_m M \rightarrow \mathbb{R}^\ell$ is injective. Consider the map

$$j: TM \rightarrow T\mathbb{R}^\ell = \mathbb{R}^\ell \times \mathbb{R}^\ell, \quad j(v) = (\iota(m), \iota_*(m)v), \quad \text{where } m := \pi(v).$$

Clearly, j is injective.

Let (U, φ) be a chart on \mathbb{R}^ℓ adapted to M . Thinking of $\mathbb{R}^\ell \times \mathbb{R}^\ell$ as the tangent bundle of \mathbb{R}^ℓ , we obtain a chart $(\pi_1^{-1}(U), \tau_\varphi) = (U \times \mathbb{R}^\ell, \tau_\varphi)$ on $\mathbb{R}^\ell \times \mathbb{R}^\ell$.

Furthermore, recall that $m \in U$ belongs to M if and only if $m = \varphi^{-1}(x, 0)$ for some $x \in \mathbb{R}^k$. Also, $v \in T_m \mathbb{R}^\ell$ belongs to $T_m M$ if and only if

$$v = \sum_{j=1}^k y_j \partial_j|_{\varphi^{-1}(x)} \iff \tau_\varphi^{-1}(v) = ((x, 0), (y, 0))$$

(notice that the summation runs to k !). Hence, $(U \times \mathbb{R}^\ell, \tau_\varphi)$ is a chart adapted to $j(TM)$ (up to a permutation of coordinates). In particular, $j(TM)$ is a $2k$ -submanifold of $\mathbb{R}^{2\ell}$.

Finally, notice that j is just the differential of ι . Here we interpret the differential of ι as a map $TM \rightarrow T\mathbb{R}^\ell$.

Exercise 4.5. Recall that the restriction of $\psi := \pi_1 \circ \varphi$ is a chart on M , where $\pi_1: \mathbb{R}^\ell = \mathbb{R}^k \times \mathbb{R}^{\ell-k} \rightarrow \mathbb{R}^k$ is the projection. Show that the coordinate representation of j with respect to the charts $(\pi^{-1}(U \cap M), \tau_\psi^{-1})$ and $(U \times \mathbb{R}^\ell, \tau_\varphi^{-1})$ is the map

$$\mathbb{R}^k \times \mathbb{R}^k \ni (x, y) \mapsto (\iota_1(x), \iota_1(y)).$$

Deduce that j is an immersion and, hence, an embedding.

Thus, we can simply identify TM with $j(TM)$ so that

$$TM = \{(u, v) \in \mathbb{R}^{2\ell} \mid u \in M \text{ and } v \in T_u M\}.$$

Example 4.6. For $M = S^k \subset \mathbb{R}^{k+1}$, we have

$$TS^k = \{(x, y) \in S^k \times \mathbb{R}^{k+1} \mid \langle x, y \rangle = 0\} \subset \mathbb{R}^{2k+2}.$$

In particular, for $k = 1$ we obtain that TS^1 is a 2-submanifold of \mathbb{R}^4 . In fact, we can realize TS^1 as a submanifold of \mathbb{R}^3 in the following sense. Consider the map

$$f: S^1 \times \mathbb{R} \rightarrow \mathbb{R}^4, \quad f(x_0, x_1; t) = (x_0, x_1, tx_1, -tx_0).$$

One can check that f is a diffeomorphism between $S^1 \times \mathbb{R} \subset \mathbb{R}^3$ and TS^1 so that we can in fact identify TS^1 with an infinite cylinder.

L7

Bibliography

- [BT03] D. Barden and C. Thomas, *An introduction to differential manifolds*, Imperial College Press, London, 2003. MR1992457 ↑10, 28
- [Mil65] J. Milnor, *Topology from the differentiable viewpoint*, 1965 (English). ↑28
- [War83] F. Warner, *Foundations of differentiable manifolds and Lie groups*, Graduate Texts in Mathematics, vol. 94, Springer-Verlag, New York-Berlin, 1983. Corrected reprint of the 1971 edition. ↑33