

Stochastic Processes

MATH 447

Nicholas Hayek

Taught by Prof. Louigi Addario-Berry

CONTENTS

I	Markov Chains	3
	<i>Time-Homogeneous Markov Chains</i>	
	<i>Multi-Step Transition Probabilities</i>	
	<i>Long Term Behavior</i>	
	Index of Definitions	

We assume working knowledge of probability and no knowledge of measure theory (though a grasp of analysis is essential). See these [MATH 356 notes](#), also taught by Louigi!

I Markov Chains

Before providing definitions, we give some examples of stochastic processes:

Eg. 1.1 A simple random walk: $S_{i+1} = S_i + X_i$, where $X_i \sim \text{Ber}(p)$ and $S_0 = 0$. We might ask: does S_i ever return to 0, i.e.

$$\mathbb{P}(\exists i > 0 : S_i = 0)$$

Eg. 1.2 A branching process: as in asexual reproduction, we have an initial node. Each node n has a number of children X_n , where $\frac{X_n}{2} \sim \text{Ber}(p)$. We denote Z_i to be the number of individuals in the i -th generation. We might ask: does Z_i ever have no children, i.e.

$$\mathbb{P}(\exists i > 0 : Z_i = 0)$$

Eg. 1.3 Choose k independent random points in the square $[0, \sqrt{k}]^2$. On average, then, there is 1 point within any unit square $U \subseteq [0, \sqrt{k}]^2$.

DEF 1.1

DEF 1.2


Given a finite or countable set V , a *Markov chain* with *state space* V is a sequence $X_n : n \geq 0$ of random variables, with $X_n \in V$, such that:

$$\underbrace{\mathbb{P}(X_{n+1} = v_{n+1})}_{\text{future}} \mid \underbrace{X_0 = v_0, \dots, X_{n-1} = v_{n-1}}_{\text{past}}, \underbrace{X_n = v_n}_{\text{present}} = \mathbb{P}(X_{n+1} = v_{n+1} \mid X_n = v_n)$$

In other words, the future only depends on the past via the present. This is called the *Markov property*.

DEF 1.3

Sometimes we allow Markov chains to be only finitely large (i.e. $0 \leq n \leq m$). For instance, we limit ourselves to one weekend of gambling in Las Vegas. A graphical example would look something like:



By repeated Bayes' Law, we observe

PROP 1.1

$$\begin{aligned} & \mathbb{P}(X_1 = v_1, \dots, X_n = v_n | X_0 = v_0) \\ &= \mathbb{P}(X_1 = v_1 | X_0 = v_0) \cdot \mathbb{P}(X_2 = v_2 | X_0 = v_0, X_1 = v_1) \cdots \mathbb{P}(X_n = v_n | X_0 = v_0, \dots, X_{n-1} = v_{n-1}) \\ &= \mathbb{P}(X_1 = v_1 | X_0 = v_0) \cdot \mathbb{P}(X_2 = v_2 | X_1 = v_1) \cdots \mathbb{P}(X_n = v_n | X_{n-1} = v_{n-1}) \quad \text{by Markov property} \end{aligned}$$

Time-Homogeneous Markov Chains

We often write
THMC

We say that a Markov chain is *time-homogeneous* if, for all $u, v \in V$ and $n \geq 0$

DEF 1.4

$$\mathbb{P}(X_{n+1} = v | X_n = u) = \mathbb{P}(X_1 = v | X_0 = u)$$

In other words, the chain's behavior is described entirely by $\mathbb{P}(X_1 = v | X_0 = u)$ for each $(v, u) \in V \times V$. In this case, we can describe such probabilities in a *transition matrix* P :

DEF 1.5

$$P = (p_{u,v})_{(u,v) \in V^2} = (\mathbb{P}(X_1 = v | X_0 = u))_{(u,v) \in V^2}$$

Eg. 1.4 Recall the game Snakes and Ladders. A 6×6 grid is indexed $1, \dots, 36$. Players start at the 1 cell. The game ends when a player reaches the 36 cell. A die roll dictates how many spots one advances. There are some directed edges between cells (increasing: "ladders", decreasing: "snakes"). One must follow these edges when one lands at its tail. Suppose a ladder exists from 11 to 27. Then

$$\mathbb{P}(X_{11} = 27 | X_{10} = 6, X_9 = 3) = \frac{1}{6} = \mathbb{P}(X_{11} = 27 | X_{10} = 6) = \mathbb{P}(X_2 = 27 | X_1 = 6)$$

We see that Snakes and Ladders is naturally modeled as a time-homogeneous Markov chain.

Eg. 1.5 Sampling without replacement is *not* a Markov chain. If we sample from $|X| = 10$, we have

$$\mathbb{P}(X_3 = a | X_2 = b) = 1/9$$

$$\mathbb{P}(X_3 = a | X_2 = b, X_1 = c) = 1/8$$

$$\mathbb{P}(X_3 = a | X_2 = b, X_1 = a) = 0$$

so we do not satisfy the Markov property.

Eg. 1.6 Returning to the Snakes and Ladders example, consider $S \subseteq V$. Let $T_S = \inf\{n \geq 0 : X_n \in S\}$. We may ask...

- What is the average number of rounds to finish? We can write this as $\mathbb{E}[T_{\{36\}} | X_0 = 1]$.
- What is the probability of landing on 18 or 19 before the game ends? We can write this as $\mathbb{P}(T_{\{18,19\}} < T_{\{36\}} | X_0 = 1)$.
- What is the average number of visits to 6 before the game ends? We can write this as

$$\mathbb{E}[\#\{n \in [T_{\{36\}}] : X_n = 6\} | X_0 = 1]$$

- What is the expected proportion of time spent on state 5 before the game ends?
- If we allow two players, what is the probability that player 1 wins? Is this still a Markov chain?

DEF 1.6 A matrix $P = (p_{u,v})_{(u,v) \in V^2}$ is called a *stochastic matrix* if every row sums to 1, i.e.

$$\forall u \in V, \sum_{v \in V} p_{u,v} = 1$$

Note that any stochastic matrix is the transition matrix of some time-homogeneous Markov chain with state space V and transition probabilities

$$\mathbb{P}(X_{n+1} = v | X_n = u) = \mathbb{P}(X_1 = v | X_0 = u) = p_{u,v}$$

A directed graph, together with its stochastic matrix, can visualize any THMC:




Fig. 1.7 Random walks on an undirected weighted graph, where edge weights dictate the proportional probability of transitioning between two states, are a special class of THMCs. In particular, given a graph $G = (V, E)$ with weights $w(e) > 0 : e \in E$, we set

$$p_{u,v} = \frac{w(\{u,v\})}{\sum_{z \in N(u)} w(\{u,z\})}$$

If there are no edges $u \leftrightarrow v$, we write $p_{u,v} = 0$.

Not every THMC can be represented by a random walk on an undirected weighted graph. In particular, see the directed graph listed above, or any transition matrix which is not symmetric.

As a concrete example, we can consider a random walk on the number line \mathbb{Z} , where, if $w(k, k+1) = \alpha$, $w(k-1, k) = \frac{\alpha}{2}$.

$$\dots \frac{1}{16} \quad -3 \quad \frac{1}{8} \quad -2 \quad \frac{1}{4} \quad -1 \quad \frac{1}{2} \quad 0 \quad 1 \quad 2 \quad 4 \quad 3 \quad 8 \quad \dots$$

Multi-Step Transition Probabilities

Given a THMC $X = X_n : n \geq 0$ with a transition matrix P , we write

$$\begin{aligned} \mathbb{P}(X_2 = v | X_0 = u) &= \sum_{w \in V} \mathbb{P}(X_2 = v, X_1 = w | X_0 = u) \\ &= \sum_{w \in V} \mathbb{P}(X_1 = w | X_0 = u) \mathbb{P}(X_2 = v | X_1 = w, X_0 = u) \quad \text{by Markov property} \\ &= \sum_{w \in V} p_{u,w} p_{w,v} = (P^2)_{u,v} \quad \text{or write } P_{u,v}^2 \end{aligned}$$

Hence, to determine a two-step transition probability, and by extension an n -step transition probability from u to w , we consider $P_{u,v}^n$. PROP 1.2

See [Prop 1.1](#) to expand probabilities, using Bayes', as needed. We get that PROOF.

$$\begin{aligned} \mathbb{P}(X_n = v | X_0 = u) &= \sum_{v_1, \dots, v_{n-1} \in V} \mathbb{P}(X_1 = v_1, \dots, X_{n-1} = v_{n-1}, X_n = v | X_0 = u) \\ &= \sum_{v_1, \dots, v_{n-1} \in V} p_{u,v_1} p(v_1, v_2) \cdots p(v_{n-1}, v) = (P^n)_{u,v} \quad \square \end{aligned}$$

Thus, if P is a stochastic matrix, then so is P^n . PROP 1.3

$$\sum_{v \in V} P_{u,v}^n = \sum_{v \in V} \mathbb{P}(X_n = v | X_0 = u) = 1. \quad \square$$
PROOF.

Theorem 1.1 Markov Property

If $X_n : n \geq 0$ is a THMC with state space V , then for all $u_0, \dots, u_{n-1}, u, v \in V$,

$$\mathbb{P}(X_{n+m} = v | X_0 = u_0, \dots, X_{n-1} = u_{n-1}, X_n = u) = \mathbb{P}(X_{n+m} = v | X_n = u) = P_{u,v}^m$$

One shows this by combining the Markov property with [Prop 1.2](#) via induction. PROOF.

Somewhat nonsensically, we *also* call this the Markov property. When talking about THMCs, this will be the default notion.

We say that a Markov chain has an *initial distribution* $\alpha = (\alpha_v : v \in V)$ if $\mathbb{P}(X_0 = v) = \alpha_v$ for each $v \in V$. If this is the case, we often write α as a subscript of our state probabilities. For instance, DEF 1.7

$$\mathbb{P}_\alpha(X_n = v) = \sum_{u \in V} \mathbb{P}_\alpha(X_0 = u, X_n = v) = \sum_{u \in V} \mathbb{P}_\alpha(X_0 = u) \mathbb{P}_\alpha(X_n = v | X_0 = u) = \sum_{u \in V} \alpha_u P_{u,v}^n$$

For any event E depending only on X_0, \dots, X_n , with $\mathbb{P}(X_n = u, E) > 0$, we have PROP 1.4

$$\mathbb{P}(X_{n+m} = v | X_n = u, E) = P_{u,v}^m$$

PROOF.

For any such event E , we can determine whether E occurs exactly when we know the realized values u_i of X_i for $i = 1, \dots, n-1$. Hence, we may write S to be the set of tuples (u_0, \dots, u_{n-1}) that guarantee E . It follows that

$$\mathbb{P}(X_n = u, E) = \sum_{\mathbf{s} \in S} \mathbb{P}(\mathbf{x} = \mathbf{s}, X_n = u)$$

Similarly, we have

$$\begin{aligned} \mathbb{P}(X_{n+m} = v, X_n = u, E) &= \sum_{\mathbf{s} \in S} \mathbb{P}(X_{n+m} = v, X_n = u, \mathbf{x} = \mathbf{s}) \\ &= \sum_{\mathbf{s} \in S} \mathbb{P}(X_{n+m} = v | X_n = u, \mathbf{x} = \mathbf{s}) \mathbb{P}(X_n = u, \mathbf{x} = \mathbf{s}) \\ &= P_{u,v}^m \sum_{\mathbf{s} \in S} \mathbb{P}(X_n = u, \mathbf{x} = \mathbf{s}) = P_{u,v}^m \mathbb{P}(X_n = u, E) \end{aligned}$$

Divide and use Bayes, and the result follows. \square

PROP 1.5 If X is a THMC with transition matrix P , then, for all $k \geq 1$, $X_{kn} : n \geq 0$ is a THMC with transition matrix P^k .

PROOF.

For any $n \neq 0$, any sequence $u_0, \dots, u_{n+1} \in V$ satisfies

$$\mathbb{P}(X_{(n+1)k} = u_{n+1} | X_0 = u_0, X_k = u_1, \dots, X_{nk} = u_n) = P_{u_n, u_{n+1}}^k \quad \square$$

Theorem 1.2 Chapman-Kolmogorov

For any Markov chain X with state space V , any $m, n \geq 0$, and $u, v \in V$,

$$\mathbb{P}(X_{m+n} = v | X_0 = u) = \sum_{w \in V} \mathbb{P}(X_n = w | X_0 = u) \mathbb{P}(X_{m+n} = v | X_n = w)$$

If the X is time homogeneous, then this is $P_{u,v}^{n+m}$, which agrees with [Prop 1.2](#).

Long Term Behavior

DEF 1.8 Recall from probability the *law of large numbers*: if $Y_n : n \geq 1$ are IID with common mean μ , then $\frac{S_n}{n} \rightarrow \mu$ in probability, where $S_n = \sum_{i=1}^n Y_i$, i.e. $\forall \varepsilon > 0$,

$$\limsup_{n \rightarrow \infty} \mathbb{P}\left(\left|\frac{S_n}{n} - \mu\right| > \varepsilon\right) = 0$$

If $Y_i \in \mathbb{Z}$ then, for $k, \ell, u_i \in \mathbb{Z}$ and $i = 1, \dots, n-1$,

$$\begin{aligned} \mathbb{P}(S_{n+1} = \ell | S_n = k, S_i = u_i \forall i) &= \mathbb{P}(Y_{n+1} = \ell - k | S_n = k, S_i = u_i \forall i) \\ &= \mathbb{P}(Y_{n+1} = \ell - k | Y_1 = u_1 - u_0, Y_2 = u_2 - u_1, \dots, Y_n = k - u_{n-1}) \\ &= \mathbb{P}(Y_{n+1} | \ell - k) = \mathbb{P}(Y_1 = \ell - k) = P_{k, \ell} \end{aligned}$$

where $S_n : n \geq 0$ has transition matrix P , noting that it may be viewed as a THMC.

From now on, we denote by $\mathbb{P}_v(E)$ the probability $\mathbb{P}(E|v)$.

Fig. 1.8 A general two-state chain, with states A and B , can be described by

$$P = \begin{pmatrix} 1 - \alpha & \alpha \\ \beta & 1 - \beta \end{pmatrix}$$

Let $q_n = \mathbb{P}_A(X_n = A) = \mathbb{P}(X_n = A|X_0 = A)$. Then

$$\begin{aligned} q_{n+1} &= \mathbb{P}_A(X_{n+1} = A, X_n = A) + \mathbb{P}_A(X_{n+1} = A, X_n = B) \\ &= \mathbb{P}_A(X_{n+1} = A|X_n = A)\mathbb{P}_A(X_n = A) + \mathbb{P}_A(X_{n+1} = A|X_n = B)\mathbb{P}_A(X_n = B) \\ &= (1 - \alpha)q_n + \beta(1 - q_n) = \beta + (1 - \alpha - \beta)q_n \end{aligned}$$

This recurrence has a unique solution. In particular, one can find

$$q_n = \frac{\beta}{\alpha + \beta} + (1 - \alpha - \beta)^n \frac{\alpha}{\alpha + \beta}$$

It follows that $q_n \rightarrow \frac{\beta}{\alpha + \beta}$, and hence $\mathbb{P}_A(X_n = B) = 1 - q_n \rightarrow \frac{\alpha}{\alpha + \beta}$. Likewise:

$$\mathbb{P}_B(X_n = B) = \frac{\alpha}{\alpha + \beta} + (1 - \alpha - \beta)^n \frac{\beta}{\alpha + \beta}$$

So $\mathbb{P}_B(X_n = B) \rightarrow \frac{\alpha}{\alpha + \beta}$.

Let $\pi := (\pi_A, \pi_B) := \left(\frac{\beta}{\alpha + \beta}, \frac{\alpha}{\alpha + \beta}\right)$ be the distribution of our initial state X_0 , Then

$$\mathbb{P}_\pi(X_1 = A) = \pi_A \mathbb{P}_A(X_1 = A) + \pi_B \mathbb{P}_B(X_1 = A) = \pi_A$$

and, similarly, $\mathbb{P}_\pi(X_1 = B) = \pi_B$. Hence, if X_0 has initial distribution π , then X_1 also has distribution π . By induction, X_n has distribution $\pi \forall n \geq 0$.

INDEX OF DEFINITIONS

initial distribution [1.7](#)

law of large numbers [1.8](#)

Markov chain [1.1](#)

Markov property [1.3](#)

state space [1.2](#)

stochastic matrix [1.6](#)

time-homogeneous [1.4](#)

transition matrix [1.5](#)