

# Stochastic Processes

MATH 447

Nicholas Hayek

*Taught by Prof. Louigi Addario-Berry*

## CONTENTS

<b>I</b>	<b>Markov Chains</b>	<b>3</b>
	Introduction . . . . .	3
	<i>Time-Homogeneous Markov Chains</i>	
	<i>Multi-Step Transition Probabilities</i>	
	Stationary and Limiting Distributions . . . . .	9
	<i>Periodicity of States</i>	
	<i>Finding Stationary Distributions</i>	
	<i>Transience and Recurrence</i>	
	<i>Canonical Decompositions</i>	
	<i>Proof of Fundamental Theorem of Markov Chains</i>	
	Hitting Times and Absorbing States . . . . .	20
	<i>Reversibility</i>	

## Index of Definitions

We assume working knowledge of probability and no knowledge of measure theory (though a grasp of analysis is essential). See these [MATH 356 notes](#), also taught by Louigi!

# I Markov Chains

Conditional expectations will be important in this course. Recall  $\mathbb{E}[X|Y = y_0]$ , where  $X, Y$  are random variables. If  $Y$  is continuous, writing  $\mathbb{E}[X|Y = y_0] = \frac{\mathbb{P}(X, Y=y_0)}{\mathbb{P}(Y=y_0)}$ , will not work. Instead, we consider the slice of the joint density function  $f(x, y)$  at  $y = y_0$ . The result is a one dimensional function  $g(x)$  which may not have probability 1. Hence, we divide by  $\int g(x)$  to make it into a density function:

$$\mathbb{E}[X|Y = y_0] = \int_{\mathbb{R}} \frac{f(x, y_0)}{\int_{\mathbb{R}} f(x, y_0) dx} x dx$$

DEF 1.1 We frequently write  $f_{X|Y}(x) = f(x, y) / \int_{\mathbb{R}} f(x, y) dx$ , and call this the *conditional density* of  $X$  given  $Y$ . For fixed  $y$ , then,  $\mathbb{E}[X|Y = y] = \mathbb{E}[Z]$ , where  $Z \sim f_{X|Y}$ .

## INTRODUCTION

Before providing definitions, we give some examples of stochastic processes:

**Eg. 1.1** A simple random walk:  $S_{i+1} = S_i + X_i$ , where  $X_i \sim \text{Ber}(p)$  and  $S_0 = 0$ . We might ask: does  $S_i$  ever return to 0, i.e.

$$\mathbb{P}(\exists i > 0 : S_i = 0)$$

**Eg. 1.2** A branching process: as in asexual reproduction, we have an initial node. Each node  $n$  has a number of children  $X_n$ , where  $\frac{X_n}{2} \sim \text{Ber}(p)$ . We denote  $Z_i$  to be the number of individuals in the  $i$ -th generation. We might ask: does  $Z_i$  ever have no children, i.e.

$$\mathbb{P}(\exists i > 0 : Z_i = 0)$$

**Eg. 1.3** Choose  $k$  independent random points in the square  $[0, \sqrt{k}]^2$ . On average, then, there is 1 point within any unit square  $U \subseteq [0, \sqrt{k}]^2$ .

DEF 1.2 Given a finite or countable set  $V$ , a *Markov chain* with *state space*  $V$  is a sequence  $X_n : n \geq 0$  of random variables, with  $X_n \in V$ , such that:

DEF 1.3

$$\underbrace{\mathbb{P}(X_{n+1} = v_{n+1})}_{\text{future}} \mid \underbrace{X_0 = v_0, \dots, X_{n-1} = v_{n-1}}_{\text{past}}, \underbrace{X_n = v_n}_{\text{present}} = \mathbb{P}(X_{n+1} = v_{n+1} \mid X_n = v_n)$$

In other words, the future only depends on the past via the present. This is called the

DEF 1.4 *Markov property*.

Sometimes we allow Markov chains to be only finitely large (i.e.  $0 \leq n \leq m$ ). For instance, we limit ourselves to one weekend of gambling in Las Vegas. A graphical example would look something like:



We can decompose walks on Markov chains by discrete "steps":

PROP 1.1

$$\mathbb{P}(X_1 = v_1, \dots, X_n = v_n | X_0 = v_0) = \prod_{i=1}^n \mathbb{P}(X_i = v_i | X_{i-1} = v_{i-1})$$

PROOF.

$$\begin{aligned} & \mathbb{P}(X_1 = v_1, \dots, X_n = v_n | X_0 = v_0) \\ &= \mathbb{P}(X_2 = v_2, \dots, X_n = v_n | X_0 = v_0, X_1 = v_1) \mathbb{P}(X_1 = v_1 | X_0 = v_0) \quad \text{by cond. prob.} \\ & \vdots \\ &= \mathbb{P}(X_1 = v_1 | X_0 = v_0) \cdot \mathbb{P}(X_2 = v_2 | X_0 = v_0, X_1 = v_1) \cdots \mathbb{P}(X_n = v_n | X_0 = v_0, \dots, X_{n-1} = v_{n-1}) \\ &= \mathbb{P}(X_1 = v_1 | X_0 = v_0) \cdot \mathbb{P}(X_2 = v_2 | X_1 = v_1) \cdots \mathbb{P}(X_n = v_n | X_{n-1} = v_{n-1}) \quad \text{by Markov prop} \end{aligned}$$

□

### Time-Homogeneous Markov Chains

We often write  
THMC

We say that a Markov chain is *time-homogeneous* if, for all  $u, v \in V$  and  $n \geq 0$

DEF 1.5

$$\mathbb{P}(X_{n+1} = v | X_n = u) = \mathbb{P}(X_1 = v | X_0 = u)$$

In other words, the chain's behavior is described entirely by  $\mathbb{P}(X_1 = v | X_0 = u)$  for each  $(v, u) \in V \times V$ . In this case, we can describe such probabilities in a *transition matrix*  $P$ :

DEF 1.6

$$P = (p_{u,v})_{(u,v) \in V^2} = (\mathbb{P}(X_1 = v | X_0 = u))_{(u,v) \in V^2}$$

**Eg. 1.4** Recall the game Snakes and Ladders. A  $6 \times 6$  grid is indexed  $1, \dots, 36$ . Players start at the 1 cell. The game ends when a player reaches the 36 cell. A die roll dictates how many spots one advances. There are some directed edges between cells (increasing: "ladders", decreasing: "snakes"). One must follow these

edges when one lands at its tail. Suppose a ladder exists from 11 to 27. Then

$$\mathbb{P}(X_{11} = 27 | X_{10} = 6, X_9 = 3) = \frac{1}{6} = \mathbb{P}(X_{11} = 27 | X_{10} = 6) = \mathbb{P}(X_2 = 27 | X_1 = 6)$$

We see that Snakes and Ladders is naturally modeled as a time-homogeneous Markov chain.

**Eg. 1.5** Sampling without replacement is *not* a Markov chain. If we sample from  $|X| = 10$ , we have

$$\mathbb{P}(X_3 = a | X_2 = b) = 1/9$$

$$\mathbb{P}(X_3 = a | X_2 = b, X_1 = c) = 1/8$$

$$\mathbb{P}(X_3 = a | X_2 = b, X_1 = a) = 0$$

so we do not satisfy the Markov property.

**Eg. 1.6** Returning to the Snakes and Ladders example, consider  $S \subseteq V$ . Let  $T_S = \inf\{n \geq 0 : X_n \in S\}$ , which we call the "*hitting time*" of  $S$ . We may ask...

- What is the average number of rounds to finite? We can write this as  $\mathbb{E}[T_{\{36\}} | X_0 = 1]$ .
- What is the probability of landing on 18 or 19 before the game ends? We can write this as  $\mathbb{P}(T_{\{18,19\}} < T_{\{36\}} | X_0 = 1)$ .
- What is the average number of visits to 6 before the game ends? We can write this as

$$\mathbb{E}[\#\{n \in [T_{\{36\}}] : X_n = 6\} | X_0 = 1]$$

- What is the expected proportion of time spent on state 5 before the game ends?
- If we allow two players, what is the probability that player 1 wins? Is this still a Markov chain?

DEF 1.7

DEF 1.8 A matrix  $P = (p_{u,v})_{(u,v) \in V^2}$  is called a *stochastic matrix* if every row sums to 1, i.e.

$$\forall u \in V, \sum_{v \in V} p_{u,v} = 1$$

Note that any stochastic matrix is the transition matrix of some time-homogeneous Markov chain with state space  $V$  and transition probabilities

$$\mathbb{P}(X_{n+1} = v | X_n = u) = \mathbb{P}(X_1 = v | X_0 = u) = p_{u,v}$$

A directed graph, together with its stochastic matrix, can visualize any THMC:



**Eg. 1.7** Random walks on an undirected weighted graph, where edge weights dictate the proportional probability of transitioning between two states, are a special class of THMCs. In particular, given a graph  $G = (V, E)$  with weights  $w(e) > 0 : e \in E$ , we set

$$p_{u,v} = \frac{w(\{u, v\})}{\sum_{z \in N(u)} w(\{u, z\})}$$

If there are no edges  $u \leftrightarrow v$ , we write  $p_{u,v} = 0$ .

Not every THMC can be represented by a random walk on an undirected weighted graph. In particular, see the directed graph listed above, or any transition matrix which is not symmetric.

As a concrete example, we can consider a random walk on the number line  $\mathbb{Z}$ , where, if  $w(k, k+1) = \alpha$ ,  $w(k-1, k) = \frac{\alpha}{2}$ .

$$\dots \frac{1}{16} \quad -3 \quad \frac{1}{8} \quad -2 \quad \frac{1}{4} \quad -1 \quad \frac{1}{2} \quad 0 \quad \frac{1}{2} \quad 1 \quad 2 \quad \frac{4}{2} \quad 3 \quad \frac{8}{2} \quad \dots$$

### Multi-Step Transition Probabilities

Given a THMC  $X = X_n : n \geq 0$  with a transition matrix  $P$ , we write

$$\begin{aligned} \mathbb{P}(X_2 = v | X_0 = u) &= \sum_{w \in V} \mathbb{P}(X_2 = v, X_1 = w | X_0 = u) \\ &= \sum_{w \in V} \mathbb{P}(X_1 = w | X_0 = u) \mathbb{P}(X_2 = v | X_1 = w, \cancel{X_0 = u}) \quad \text{by Markov property} \\ &= \sum_{w \in V} p_{u,w} p_{w,v} = (P^2)_{u,v} \quad \text{or write } P_{u,v}^2 \end{aligned}$$

Hence, to determine a two-step transition probability, and by extension an  $n$ -step transition probability from  $u$  to  $w$ , we consider  $P_{u,v}^n$ . PROP 1.2

PROOF.

See [Prop 1.1](#) to expand probabilities, using Bayes', as needed. We get that

$$\begin{aligned}\mathbb{P}(X_n = v | X_0 = u) &= \sum_{v_1, \dots, v_{n-1} \in V} \mathbb{P}(X_1 = v_1, \dots, X_{n-1} = v_{n-1}, X_n = v | X_0 = u) \\ &= \sum_{v_1, \dots, v_{n-1} \in V} p_{u, v_1} p(v_1, v_2) \cdots p(v_{n-1}, v) = (P^n)_{u, v} \quad \square\end{aligned}$$

**PROP 1.3** Thus, if  $P$  is a stochastic matrix, then so is  $P^n$ .

PROOF.

$$\sum_{v \in V} P_{u, v}^n = \sum_{v \in V} \mathbb{P}(X_n = v | X_0 = u) = 1. \quad \square$$

### Theorem 1.1 Markov Property

If  $X_n : n \geq 0$  is a THMC with state space  $V$ , then for all  $u_0, \dots, u_{n-1}, u, v \in V$ ,

$$\mathbb{P}(X_{n+m} = v | X_0 = u_0, \dots, X_{n-1} = u_{n-1}, X_n = u) = \mathbb{P}(X_{n+m} = v | X_n = u) = P_{u, v}^m$$

PROOF.

One shows this by combining the Markov property with [Prop 1.2](#). □

We *also* call this the Markov property.

DEF 1.9

We say that a Markov chain has an *initial distribution*  $\alpha = (\alpha_v : v \in V)$  if  $\mathbb{P}(X_0 = v) = \alpha_v$  for each  $v \in V$ . If this is the case, we often write  $\alpha$  as a subscript of our state probabilities. For instance,

$$\mathbb{P}_\alpha(X_n = v) = \sum_{u \in V} \mathbb{P}_\alpha(X_0 = u, X_n = v) = \sum_{u \in V} \mathbb{P}_\alpha(X_0 = u) \mathbb{P}_\alpha(X_n = v | X_0 = u) = \sum_{u \in V} \alpha_u P_{u, v}^n$$

In particular, [Thm 1.1](#) is the Markov property combined with a claim about  $P_{u, v}^m$

**PROP 1.4** For an event  $E$ , depending only on  $X_0, \dots, X_n$ , with  $\mathbb{P}(X_n = u, E) > 0$ , we have

$$\mathbb{P}(X_{n+m} = v | X_n = u, E) = P_{u, v}^m$$

PROOF.

For any such event  $E$ , we can determine whether  $E$  occurs exactly when we know the realized values  $u_i$  of  $X_i$  for  $i = 1, \dots, n-1$ . Hence, we may write  $\mathcal{S}$  to be the set of tuples  $(u_0, \dots, u_{n-1})$  that guarantee  $E$ . It follows that

$$\mathbb{P}(X_n = u, E) = \sum_{\mathbf{s} \in \mathcal{S}} \mathbb{P}(\mathbf{X} = \mathbf{s}, X_n = u)$$

Similarly, we have

$$\begin{aligned}
 \mathbb{P}(X_{n+m} = v, X_n = u, E) &= \sum_{\mathbf{s} \in \mathcal{S}} \mathbb{P}(X_{n+m} = v, X_n = u, \mathbf{X} = \mathbf{s}) \\
 &= \sum_{\mathbf{s} \in \mathcal{S}} \mathbb{P}(X_{n+m} = v | X_n = u, \mathbf{X} = \mathbf{s}) \mathbb{P}(X_n = u, \mathbf{X} = \mathbf{s}) \\
 &= P_{u,v}^m \sum_{\mathbf{s} \in \mathcal{S}} \mathbb{P}(X_n = u, \mathbf{X} = \mathbf{s}) = P_{u,v}^m \mathbb{P}(X_n = u, E)
 \end{aligned}$$

Divide and use Bayes, and the result follows.  $\square$

If  $X$  is a THMC with transition matrix  $P$ , then, for all  $k \geq 1$ ,  $X_{kn} : n \geq 0$  is a THMC with transition matrix  $P^k$ . PROP 1.5

For any  $n \neq 0$ , any sequence  $u_0, \dots, u_{n+1} \in V$  satisfies PROOF.

$$\mathbb{P}(X_{(n+1)k} = u_{n+1} | X_0 = u_0, X_k = u_1, \dots, X_{nk} = u_n) = P_{u_n, u_{n+1}}^k \quad \square$$

### Theorem 1.2 Chapman-Kolmogorov

For any Markov chain  $X$  with state space  $V$ , any  $m, n \geq 0$ , and  $u, v \in V$ ,

$$\mathbb{P}(X_{m+n} = v | X_0 = u) = \sum_{w \in V} \mathbb{P}(X_n = w | X_0 = u) \mathbb{P}(X_{m+n} = v | X_n = w)$$

If the  $X$  is time homogeneous, then this is  $P_{u,v}^{n+m}$ , which agrees with [Prop 1.2](#).

$$\begin{aligned}
 \mathbb{P}(X_{m+n} = v | X_0 = u) &= \sum_{w \in V} \mathbb{P}(X_{m+n} = v, X_n = w | X_0 = u) \\
 &= \sum_{w \in V} \mathbb{P}(X_{m+n} = v | X_n = w, X_0 = u) \mathbb{P}(X_n = w | X_0 = u) \\
 &= \sum_{w \in V} \mathbb{P}(X_{m+n} = v | X_n = w) \mathbb{P}(X_n = w | X_0 = u)
 \end{aligned}$$

Below is a visualization of the Chapman-Kolmogorov equation: to get from  $u$  to  $v$  over 6 steps, we pick some partition (e.g.  $2 + 4 = 6$ ), and sum the probabilities of reaching  $v$  via



length 2 (red) and length 4 (blue) paths.

You should count 6 such combinations in the sample below.



## STATIONARY AND LIMITING DISTRIBUTIONS

DEF 1.10 Recall from probability the *law of large numbers*: if  $Y_n : n \geq 1$  are IID with common mean  $\mu$ , then  $\frac{S_n}{n} \rightarrow \mu$  in probability, where  $S_n = \sum_{i=1}^n Y_i$ , i.e.  $\forall \varepsilon > 0$ ,

$$\limsup_{n \rightarrow \infty} \mathbb{P} \left( \left| \frac{S_n}{n} - \mu \right| > \varepsilon \right) = 0$$

If  $Y_i \in \mathbb{Z}$  is a THMC, for  $k, \ell \in \mathbb{Z}$  and  $i \in [n-1]$ ,

$$\begin{aligned} P_{k,\ell} &:= \mathbb{P}(S_{n+1} = \ell | S_n = k, S_i = u_i \forall i) = \mathbb{P}(Y_{n+1} = \ell - k | S_n = k, S_i = u_i \forall i) \\ &= \mathbb{P}(Y_{n+1} = \ell - k | Y_1 = u_1 - u_0, Y_2 = u_2 - u_1, \dots, Y_n = k - u_{n-1}) \\ &= \mathbb{P}(Y_{n+1} = \ell - k) = \mathbb{P}(Y_1 = \ell - k) \end{aligned}$$

We conclude that  $S_n : n \geq 0$  may be viewed as a THMC, with transition matrix  $P$ .

From now on, we denote by  $\mathbb{P}_v(E)$  the probability  $\mathbb{P}(E|v)$ .

**Eg. 1.8** A general two-state chain, with states  $A$  and  $B$ , can be described by

$$P = \begin{pmatrix} 1 - \alpha & \alpha \\ \beta & 1 - \beta \end{pmatrix}$$

Let  $q_n = \mathbb{P}_A(X_n = A) = \mathbb{P}(X_n = A | X_0 = A)$ . Then

$$\begin{aligned} q_{n+1} &= \mathbb{P}_A(X_{n+1} = A, X_n = A) + \mathbb{P}_A(X_{n+1} = A, X_n = B) \\ &= \mathbb{P}_A(X_{n+1} = A | X_n = A) \mathbb{P}_A(X_n = A) + \mathbb{P}_A(X_{n+1} = A | X_n = B) \mathbb{P}_A(X_n = B) \\ &= (1 - \alpha)q_n + \beta(1 - q_n) = \beta + (1 - \alpha - \beta)q_n \end{aligned}$$

This recurrence has a unique solution. In particular, one can find

$$q_n = \frac{\beta}{\alpha + \beta} + (1 - \alpha - \beta)^n \frac{\alpha}{\alpha + \beta}$$

It follows that  $q_n \rightarrow \frac{\beta}{\alpha+\beta}$ , and hence  $\mathbb{P}_A(X_n = B) = 1 - q_n \rightarrow \frac{\alpha}{\alpha+\beta}$ . Likewise:

$$\mathbb{P}_B(X_n = B) = \frac{\alpha}{\alpha+\beta} + (1 - \beta)^n \frac{\beta}{\alpha+\beta}$$

So  $\mathbb{P}_B(X_n = B) \rightarrow \frac{\alpha}{\alpha+\beta}$ .

Let  $\pi := (\pi_A, \pi_B) := \left(\frac{\beta}{\alpha+\beta}, \frac{\alpha}{\alpha+\beta}\right)$  be the distribution of our initial state  $X_0$ . Then

$$\mathbb{P}_\pi(X_1 = A) = \pi_A \mathbb{P}_A(X_1 = A) + \pi_B \mathbb{P}_B(X_1 = A) = \pi_A$$

and, similarly,  $\mathbb{P}_\pi(X_1 = B) = \pi_B$ . Hence, if  $X_0$  has initial distribution  $\pi$ , then  $X_1$  also has distribution  $\pi$ . By induction,  $X_n$  has distribution  $\pi \forall n \geq 0$ .

When we say  $X = \text{Markov}(P)$ , we mean that  $X$  is a THMC with transition matrix  $P$ .

A probability distribution  $\pi$  is called **stationary** if  $\pi P = \pi$ . Similarly, a probability distribution  $\lambda$  is called a **limiting distribution** if, for each  $u, v \in V$

DEF 1.11

DEF 1.12

$$(P^n)_{u,v} \rightarrow \lambda_v \text{ as } n \rightarrow \infty$$

In other words,  $\mathbb{P}_u(X_n = v) \rightarrow \lambda_v$ . Note that, for any initial distribution  $\alpha$ , we have  $\alpha P^n \rightarrow \lambda$ , i.e.  $(\alpha P^n)_v \rightarrow \lambda_v$ , where  $\lambda$  is limiting. As proof:

PROOF.

$$(\alpha P^n)_v = \sum_{u \in V} \alpha_u P^n_{u,v} \rightarrow \sum_{u \in V} \alpha_u \lambda_v = \lambda_v$$

□

If  $\lambda$  is a limiting distribution for  $P$ , then  $\lambda$  is stationary for  $P$ .

PROP 1.6

Fix any initial distribution  $\alpha$ , we have

PROOF.

$$\lambda = \lim_{n \rightarrow \infty} (\alpha P^n) = \lim_{n \rightarrow \infty} (\alpha P^{n-1} P) = \left( \lim_{n \rightarrow \infty} \alpha P^{n-1} \right) P = \lambda P \quad \square$$

Stationary distributions need not be unique, but limiting distributions are (as the limit  $\lim_{n \rightarrow \infty} \alpha P^n$  is well-defined). In general, then, stationary distributions need not be limiting distributions.

A stochastic matrix  $P$  is called **regular** if  $\exists n \geq 1$  such that  $P^n > 0$  on all entries.

DEF 1.13

### Theorem 1.3 Fundamental Theorem of Markov Chains

Every finite, regular stochastic matrix  $P$  has a limiting distribution  $\pi$ .

Incorporating some of the formulations above, this is equivalent to saying: *For a regular stochastic matrix, there exists a unique distribution  $\pi = (\pi_v : v \in V)$  such that  $\pi P = \pi$  and  $\mathbb{P}_u(X_n = v) \rightarrow \pi_v \forall u, v \in V$ .*

When  $n = 0$ ,  $P^n = I$ , which encapsulates the idea that, at timestep 0, we will be at our initial positions.

In this case, there is a unique stationary distribution, and it is the unique limiting distribution.

### A stationary distribution always exists!

Let  $\rho = \langle 1, \dots, 1 \rangle$ . Then note that  $P\rho = \rho$ , since the sum of any row in  $P$  must be 1. Hence,  $P$  has eigenvalue 1. It follows that it has a left eigenvector, i.e.  $\pi : \pi P = \pi$ . This is exactly a stationary distribution, as long as we scale suitably such that  $\pi$  is a distribution.

However, then process of scaling into a distribution is non-trivial. Since  $\pi$  may have negative coordinates, and hence  $\sum \pi_i = 0$ , we must consider instead  $|\pi|$ , i.e. prove it is also an eigenvalue.

This is true, but requires the fact that  $P$  is stochastic

### Periodicity of States

DEF 1.14 For  $u, v \in V$ , we say that  $v$  is *accessible* from  $u$  if  $\exists n \geq 0$  such that  $(P^n)_{u,v} > 0$ . Equivalently, in the directed graph generated by  $P$ , there is a directed path from  $u$  to  $v$ . When  $v$  is accessible from  $u$ , we write  $u \rightarrow v$ .

DEF 1.15 States  $u$  and  $v$  *communicate* if  $u \rightarrow v$  and  $v \rightarrow u$ . When  $u$  and  $v$  communicate, we write  $u \leftrightarrow v$ . Observe that communication is an equivalence relation. Hence, the state space  $V$  can be written as a disjoint union of mutually-communicating states, called a

DEF 1.16 *communication class*. Note that, in the directed graph generated by  $P$ , these correspond to the strongly connected components.

DEF 1.17 We say that  $P$  is *irreducible* if there is only one communication class.

Clearly, if  $P$  is regular, then it is irreducible

PROP 1.7  $u \rightarrow v \iff \mathbb{P}_u(T_{\{v\}} < \infty) > 0$ .

DEF 1.18 The *period* of a state  $u \in V$  is

$$d(u) := \gcd(n > 0 : P_{u,u}^n > 0)$$

DEF 1.19 If  $d(u) = 1$ , we call  $u$  *aperiodic*. By extension,  $P$  is aperiodic if  $d(u) = 1 \forall u \in V$ , and  $X$  is aperiodic if  $X = \text{Markov}(P)$  for  $P$  aperiodic.

PROP 1.8 If  $u \leftrightarrow v$ , then  $d(u) = d(v)$ .

PROOF.

Let  $I = \{n > 0 : P_{u,u}^n > 0\}$ , and similarly  $J$  for  $v$ . Hence,  $d(u) = \gcd(I)$  and  $d(v) = \gcd(J)$ . Let  $a, b > 0$  such that  $P_{u,v}^a > 0$  and  $P_{v,u}^b > 0$ . Then

$$P_{u,u}^{a+b} \geq P_{u,v}^a P_{v,u}^b > 0$$

$\implies a + b \in I$ , so  $d(u) | a + b$ . Now, if  $n \in J$ , then

$$P_{u,u}^{a+b+n} \geq P_{u,v}^a P_{v,v}^n P_{v,u}^b > 0$$

$\implies a + b + n \in I$ , so  $d(u) | n + a + b$ . But, by the previous line,  $d(u) | n$ . Since  $n \in J$  is arbitrary, we can write  $d(u) | \gcd(J) = d(v)$ .

Symmetrically, we could conclude that  $d(v) | d(u)$ , so indeed  $d(v) = d(u)$ .  $\square$

PROP 1.9 Let  $I = \{n > 0 : P_{u,u}^n > 0\}$ . If  $\gcd(I) = 1$ , then  $\exists a, b \in I$  such that  $\gcd(a, b) = 1$ .

This is not true for any  $I$  (and thus relies not only on number theory). Let  $\ell, m \in I$ , with  $\ell < m$ . Let  $k = m - \ell$ . If  $k = 1$ , then  $\gcd(\ell, m) = 1$ . Otherwise, since  $\gcd(I) = 1$ , there is an  $n \in I$  with  $k \nmid n$ . We then write  $n = qk + r$ , with  $r \in [k - 1]$ . Then  $m' \in (q + 1)m \in I$ , since  $P_{u,u}^{(q+1)m} \geq (P_{u,u}^m)^{q+1}$ . Symmetrically, we can argue  $\ell' = (q + 1)\ell \in I$ .

Similarly,  $\ell^* := \ell' + n \in I$ , since  $P_{u,u}^{\ell'+n} \geq P_{u,u}^{\ell'} P_{u,u}^n$ . We have

$$\begin{aligned} m' - \ell^* &= (q + 1)m - (q + 1)\ell - n = (q + 1)(m - \ell) - n \\ &= (q + 1)k - n = k - r \in [k - 1] \end{aligned}$$

TODO...

PROOF.

#### Theorem 1.4 Postage Stamp Lemma

If  $P$  is irreducible and aperiodic, then  $\forall u, v \in V, \exists N$  such that  $P_{u,v}^n > 0 \forall n \geq N$ .

Before proving this, we note that, for  $a, b \geq 1$  with  $\gcd(a, b) = 1$ , then for any  $q \geq ab$ , we can write  $q = ja + kb$  for integers  $j, k \geq 0$ .

Fix  $u, v \in V$ . Since  $P$  is aperiodic, there are  $a, b \geq 1$  with  $P_{u,u}^a, P_{u,u}^b > 0$  and  $\gcd(a, b) = 1$ , by [Prop 1.9](#). Since  $P$  is irreducible, there is some  $m > 0$  with  $P_{u,v}^m > 0$ . Thus, let  $N = m + ab$ . For any  $n \geq N$ , let  $q = n - m$ . We have that  $q \geq ab$ , so we can find  $j, k \geq 0$  with  $q = ja + kb$ . Then

$$P_{u,v}^n = P_{u,v}^{q+m} = P_{u,v}^{ja+kb+m} \geq P_{u,u}^{ja} P_{u,u}^{kb} P_{u,v}^m \geq (P_{u,u}^a)^j (P_{u,u}^b)^k P_{u,v}^m$$

All are positive, so  $P_{u,v}^n > 0$ , as desired.

PROOF.

#### Theorem 1.5 Characterization of Regular Markov Chains

Let  $P = (p_{u,v})_{u,v \in V}$  be a stochastic matrix, where  $|V| < \infty$ . Then

$$P \text{ is regular} \iff P \text{ is irreducible and aperiodic}$$

We first note why finiteness is necessary. Consider:



with all edges having weight 1. This graph is clearly aperiodic and irreducible, but not regular.

( $\implies$ ) If  $P$  is regular, then  $\exists n > 0$  s.t.  $P_{u,v}^n > 0$  for all  $u, v \in V$ . Then, for all  $u, v \in V$ , we have  $u \rightarrow v$  and  $v \rightarrow u$ . Hence,  $P$  is irreducible. Now, if  $P$  is irreducible, then for all  $u \in V$ , there is some  $v \in V$  such that  $P_{v,u} > 0$  (think about this in graph theoretic

PROOF.

terms). Then, let  $n > 0$  be such that  $P_{u,u}^n$  is positive. We have

$$P_{u,u}^{n+1} \geq P_{u,v}^n P_{u,v} > 0$$

So, with  $I = \{m > 0 : P_{u,u}^m > 0\}$ ,  $d(u) = \gcd(I) \leq \gcd(n, n+1) = 1$ . It follows that  $d(u) = 1$ , so  $u$  is aperiodic (and hence  $P$  is aperiodic).

( $\Leftarrow$ ) By [Thm 1.4](#), for each  $u, v \in V$ , there exists  $N : P_{u,v}^n > 0 \forall n \geq N$ . Let  $N^*$  be the maximum value of  $N$  determined over all pairs  $(u, v) \in V^2$ . Then, for  $n \geq N^*$  and all  $u, v \in V$ ,  $P_{u,v}^n > 0$ . It follows that all entries of  $P^n$  are positive, and we are done.  $\square$

### Finding Stationary Distributions

Recall that  $x = (x_v : v \in V)$  is a stationary distribution if  $xP = x$ . Let  $V$  be finite. Then, for a stationary distribution  $x$ , we have

$$\begin{aligned} x_1 p_{1,1} + \cdots + x_n p_{n,1} &= x_1 \\ x_1 p_{1,2} + \cdots + x_n p_{n,2} &= x_2 \\ &\vdots \\ x_1 p_{1,n} + \cdots + x_n p_{n,n} &= x_n \end{aligned}$$

We have  $n$  equations,  $n$  unknowns, and a homogeneous system, so there is not a unique solution. If

$$P = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0.5 & 0.5 \\ 0.5 & 0 & 0.5 \end{pmatrix}$$

We can compute  $x = \langle t, 2t, 2t \rangle$ . But, noting that  $x$  is a probability distribution, and hence  $5t = 1$ , this yields  $x = \langle 1/5, 2/5, 2/5 \rangle$ . We'll consider some special cases.

### UNDIRECTED GRAPHS

Let  $G = (V, E)$  be undirected. Then we define a THMC by

$$P_{u,v} = \begin{cases} \frac{1}{\deg(u)} & \{u, v\} \in E \\ 0 & \text{o.w.} \end{cases}$$

This is distinct from [Example 1.7](#)

Let  $x = (\deg(v) : v \in V)$ . We have

$$\begin{aligned} (xP)_v &= \sum_{u \in V} \deg(u) P_{u,v} = \sum_{u \in N(v)} \deg(u) \cdot \frac{1}{\deg(u)} \\ &= \deg(v) \end{aligned}$$

Hence,  $xP = x$ . Recalling that  $\sum_{v \in V} \deg(v) = 2|E|$ , we conclude that

$$\left( \frac{\deg(v)}{2|E|} : v \in V \right)$$

is a stationary distribution.

## UNDIRECTED WEIGHTED GRAPHS

This is not distinct from [Example 1.7](#)

Let  $G = (V, E)$  be undirected. Then, we define a THMC by

$$P_{u,v} = \begin{cases} \frac{w(\{u,v\})}{\sum_{z \in N(u)} w(\{u,z\})} & v \in N(u) \\ 0 & \text{o.w.} \end{cases}$$

Let  $x = (\sum_{e: e \ni v} w(e) : v \in V)$ . Then we can compute  $xP = x$ , and similar to above,

$$x = \left( \frac{\sum_{e: e \ni v} w(e)}{2 \sum_{e \in E} w(e)} : v \in V \right)$$

is a stationary distribution.

*Transience and Recurrence*

Recall  $T_S = \inf\{n \geq 0 : X_n \in S\}$ , the "hitting time" of  $S$ . We let  $R_S = \inf\{n > 0 : X_n \in S\}$ . Note that if  $T_S > 0$ ,  $T_S = R_S$ . Otherwise,  $R_S$  gives the first "*return time*" to the set  $S$ .

DEF 1.20

A state  $v \in V$  is called *recurrent* if  $\mathbb{P}_v(R_{\{v\}} < \infty) = 1$ . If all states of  $v$  are recurrent, we may call  $P$  and  $X = \text{Markov}(P)$  recurrent. Otherwise, we call  $v$  *transient*, and similarly extend the notion to the transition matrix and chain when all state are transient.

DEF 1.21

DEF 1.22

For a given state  $v \in V$ , we call  $L_v = |\{n \geq 0 : X_n = v\}|$  the *local time* of  $v$ . Note that, if  $v = X_j$  and  $v$  is recurrent, then  $L_v = \infty$ .

DEF 1.23

Intuitively, if  $L_v > k$  when  $X_0 = v$ , then  $L_v > k - 1$  when  $X_{i_1} = v$ , where  $i_1$  is the first time we return to  $v$ .

Let  $X = \text{Markov}(P)$ . For any state  $v \in V$  and  $k > 1$ ,

PROP 1.10

$$\mathbb{P}_v(L_v > k) = \mathbb{P}_v(L_v > 1)^k$$

Using the law of total probability:

PROOF.

$$\begin{aligned} \mathbb{P}_v(L_v > k) &= \mathbb{E}[\mathbb{P}_v(L_v > k | R_v)] = \sum_{t=1}^{\infty} \mathbb{P}_v(R_v = t) \mathbb{P}_v(L_v > k | R_v = t) \\ &= \sum_{t=1}^{\infty} \mathbb{P}_v(R_v = t) \mathbb{P}_v(L_v > k | R_v = t, X_t = v) \\ &= \sum_{t=1}^{\infty} \mathbb{P}_v(R_v = t) \mathbb{P}_v(L_v > k - 1) \\ &= \mathbb{P}_v(L_v > k - 1) \sum_{t=1}^{\infty} \mathbb{P}_v(R_v = t) \\ &= \mathbb{P}_v(L_v > k - 1) \mathbb{P}_v(R_v < \infty) = \mathbb{P}_v(L_v > k - 1) \mathbb{P}_v(L_v > 1) \end{aligned}$$

The result follows by induction. □

PROP 1.11

$$\mathbb{P}_v(L_v = \infty) = \begin{cases} 1 & v \text{ recurrent} \\ 0 & v \text{ transient} \end{cases}$$

PROOF.

This follows directly from [Prop 1.10](#) + monotonicity of probability.  $\square$

PROP 1.12

$$\sum_{n=0}^{\infty} P_{v,v}^n = \begin{cases} \infty & v \text{ recurrent} \\ \frac{1}{1 - \mathbb{P}_v(R_{\{v\}} < \infty)} & v \text{ transient} \end{cases}$$

PROOF.

This follows from linearity of expectation, and the fact that, for a non-negative integer variable  $X$ ,  $\mathbb{E}[X] = \sum_{i=0}^{\infty} \mathbb{P}(X > i)$  (We use, and prove, this fact later).

$$\begin{aligned} \sum_{n=0}^{\infty} P_{v,v}^n &= \sum_{n=0}^{\infty} \mathbb{P}_v(X_n = v) = \sum_{i=0}^{\infty} \mathbb{E}[\mathbb{1}_{X_i=v}] \\ &= \mathbb{E}\left[\sum_{i=0}^{\infty} \mathbb{1}_{X_i=v}\right] = \mathbb{E}_v[L_v] = \sum_{i=0}^{\infty} \mathbb{P}(L_v > i) \\ &= \sum_{i=0}^{\infty} \mathbb{P}_v(L_v > 1)^i = \begin{cases} \infty & X \text{ recurrent} \\ \frac{1}{1 - \mathbb{P}_v(L_v > 1)} & X \text{ transient} \end{cases} \end{aligned}$$

Note that  $\mathbb{P}_v(L_v > 1) \equiv \mathbb{P}_v(R_v < \infty)$   $\square$

PROP 1.13 If  $u \leftrightarrow v$ , then  $u$  is transient  $\iff v$  is transient.

PROOF.

Fix  $a, b \geq 0$  with  $P_{u,v}^a, P_{v,u}^b > 0$ . Then

$$\begin{aligned} \sum_{n=0}^{\infty} P_{v,v}^n &\geq \sum_{n=0}^{\infty} P_{v,v}^{a+b+n} = \sum_{n=0}^{\infty} P_{v,u}^b P_{u,u}^n P_{u,v}^a \\ &= P_{v,u}^b P_{u,v}^a \sum_{n=0}^{\infty} P_{u,u}^n \end{aligned}$$

Thus, if  $v$  is transient, then  $\sum_{n=0}^{\infty} P_{v,v}^n < \infty$ , so it must be that  $\sum_{n=0}^{\infty} P_{u,u}^n < \infty$ , i.e.  $u$  is transient. The argument is identical in reverse.  $\square$

**Eg. 1.9** If  $u \leftrightarrow v$  and  $u$  is recurrent, then  $\mathbb{P}_u(T_{\{v\}} < \infty) = 1$ .

**Eg. 1.10** The following chain is completely transient:



In fact, we could replace  $2/3$  by  $p$ , and  $1/3$  by  $1 - p$ . In this case, the chain is irreducible. To see that it is transient, we have

$$P_{0,0}^{2n} = \binom{2n}{n} p^n (1-p)^n$$

Then

$$\sum_{n=0}^{\infty} P_{0,0}^{2n} < \sum_{n=0}^{\infty} 2^{2n} p^n (1-p)^n = \sum_{n=0}^{\infty} (4p(1-p))^n < \infty \quad \text{if } p \neq \frac{1}{2}$$

By [Prop 1.12](#). Notice that  $\sum_{n=0}^{\infty} P_{0,0}^n = \sum_{n=0}^{\infty} P_{0,0}^{2n}$ , since it is only possible to return on even-length cycles.

We conclude that the chain is transient when  $p \neq \frac{1}{2}$ .

Stirling's Formula provides

$$m! \sim \left(\frac{m}{e}\right)^m \sqrt{2\pi m}$$

FACT

in that

$$\lim_{m \rightarrow \infty} \frac{m!}{\left(\frac{m}{e}\right)^m \sqrt{2\pi m}} = 1$$

These facts, though out of the scope of this course, can be derived from a careful treatment of Reimann sums

This fact implies

$$e \left(\frac{n}{e}\right)^n \leq n! \leq \frac{e(n+1)}{4} \left(\frac{n+1}{e}\right)^n$$

**Fig. 1.11** We return to the previous example, letting  $p = \frac{1}{2}$ . Then

$$\begin{aligned} \sum_{n=0}^{\infty} P_{0,0}^{2n} &= \sum_{n=0}^{\infty} \frac{(2n)!}{(n!)^2} [p(1-p)]^n \sim \sum_{n=0}^{\infty} \frac{\left(\frac{2n}{e}\right)^{2n} \sqrt{2\pi(2n)}}{\left[\left(\frac{n}{e}\right)^n \sqrt{2\pi n}\right]^2} [p(1-p)]^n \\ &= \sum_{n=0}^{\infty} \frac{[4p(1-p)]^n}{\sqrt{\pi n}} = \sum_{n=0}^{\infty} \frac{1}{\sqrt{\pi n}} = \infty \end{aligned}$$

We conclude that the chain is recurrent when  $p = \frac{1}{2}$ .

If  $V$  is finite, then there is at least one recurrent state.

PROP 1.14

Fix an initial distribution  $\alpha = (\alpha_v : v \in V)$ . Then  $\mathbb{P}_\alpha(\sum_{v \in V} L_v = \infty) = 1$ . We conclude

PROOF.



that there is at least one state  $v \in V$  with  $\mathbb{P}_\alpha(L_v = \infty) > 0$ . But also:

$$\begin{aligned} \mathbb{P}_\alpha(L_v = \infty) &= \sum_{n=0}^{\infty} \mathbb{P}_\alpha(L_v = \infty, T_v = n) = \sum_{n=0}^{\infty} \mathbb{P}_\alpha(L_v = \infty | T_v = n) \mathbb{P}_\alpha(T_v = n) \\ &= \sum_{n=0}^{\infty} \mathbb{P}_\alpha(L_v = \infty | T_v = n, X_n = v) \mathbb{P}_\alpha(T_v = n) \\ &= \sum_{n=0}^{\infty} \mathbb{P}_\alpha(L_v = \infty | X_n = v) \mathbb{P}_\alpha(T_v = n) \\ &= \sum_{n=0}^{\infty} \mathbb{P}_v(L_v = \infty) \mathbb{P}_\alpha(T_v = n) \end{aligned}$$

So  $\mathbb{P}_v(L_v = \infty) > 0 \implies \mathbb{P}_v(L_v = \infty) = 1$ , by [Prop 1.11](#).  $\square$

**PROP 1.15** Finite, irreducible chains are recurrent.

PROOF.

Since the chain is finite it has at least one recurrent state, by [Prop 1.14](#). Then all states must be recurrent, since the chain is irreducible, by [Prop 1.13](#).  $\square$

### Canonical Decompositions

Fix a transition matrix  $P$  and list the communication classes of  $V$  as

$$D_1, D_2, \dots \quad (\text{transient}) \quad C_1, C_2, \dots \quad (\text{recurrent})$$

**DEF 1.24** Note that we can split the chain up in this way by [Prop 1.13](#). Set  $D = \cup_{i \geq 0} D_i$ . Then the *canonical decomposition* of the chain is defined to be

$$D \sqcup C_1 \sqcup C_2 \sqcup \dots$$

**DEF 1.25** We say that a communication class  $C$  is *closed* if, for any  $u \in C, v \notin C, p_{u,v} = 0$ . Intuitively, if  $X_0 \in C$ , or we enter  $C$  at some later time, we will never leave  $C$ .

**PROP 1.16** If  $C$  is a recurrent communication class, then  $C$  is closed.

PROOF.

Fix  $u \in C, v \notin C$ . Suppose  $v \mapsto u$ . If  $p_{u,v} > 0$ , then  $u \mapsto v$ , so  $v \in C$   $\nmid$ . Suppose  $v \not\mapsto u$ . Then  $\mathbb{P}_u(R_u = \infty) \geq \mathbb{P}_u(X_1 = v) = p_{u,v}$ . But  $\mathbb{P}_u(R_u = \infty) = 0$ , since  $u$  is recurrent. It follows that  $p_{u,v} = 0$ .  $\square$

The converse of [Prop 1.16](#) is not true in generality, but it is in the finite case:

**PROP 1.17** Finite, closed communication classes are recurrent.

PROOF.

From any starting state in  $C$ , we must visit some state  $u \in C$  infinitely often, as  $|C| < \infty$  and  $X_t \in C \forall t$ . But recurrence is a class property by [Prop 1.13](#). Hence, all of  $C$  is recurrent.  $\square$

When our communication classes are closed, we have



### *Proof of Fundamental Theorem of Markov Chains*

Recall [Thm 1.3](#):

**Every finite, regular stochastic matrix  $P$  has a limiting distribution  $\pi$ .**

We will prove this in two steps. First, we will find some stationary distribution. Then, we will prove that this is a limiting distribution.

#### **Theorem 1.6 Existence Theorem**

Let  $P$  be irreducible and recurrent. Let  $(X_n : n \geq 0) = \text{Markov}(P)$ . Fix  $u \in V$ , which we call a reference vertex, and, for any  $v \in V$ , define

$$\gamma_v = \mathbb{E}_u[|\{0 \leq n < R_u : X_n = v\}|]$$

Let  $\gamma = (\gamma_v : v \in V)$ . Then  $\gamma P = \gamma$ , and  $0 < \gamma_v < \infty \forall v \in V$ .

By  $\mathbb{E}_u$ , we mean the expectation, under the assumption that  $X_0 = u$

Observe that  $\gamma_u = 1$ . Write

$$\gamma_v = \mathbb{E}_u \left[ \sum_{n=0}^{R_u-1} \mathbb{1}_{X_n=v} \right] = \mathbb{E}_u \left[ \sum_{n=1}^{R_u} \mathbb{1}_{X_n=v} \right] = \mathbb{E}_u \left[ \sum_{n=1}^{\infty} \mathbb{1}_{X_n=v} \mathbb{1}_{n \leq R_u} \right]$$

We utilize a second indicator variable in order to use linearity of expectation (otherwise, our sum would index over a random variable, which is not valid). Then

$$\gamma_v = \sum_{n=1}^{\infty} \mathbb{E}_u[\mathbb{1}_{X_n=v} \mathbb{1}_{n \leq R_u}] = \sum_{n=1}^{\infty} \mathbb{P}_u(X_n = v, n \leq R_u)$$

Now, using the law of total probability,

$$\begin{aligned} \mathbb{P}_u(X_n = v, n \leq R_u) &= \sum_{w \in V} \mathbb{P}_u(X_{n-1} = w, X_n = v, n \leq R_u) \\ &= \sum_{w \in W} \mathbb{P}_u(X_n = v | X_{n-1} = w, n \leq R_u) \mathbb{P}_u(X_{n-1} = w, n \leq R_u) \\ &= \sum_{w \in V} \mathbb{P}_u(X_{n-1} = w, n \leq R_u) P_{w,v} \end{aligned}$$

PROOF.

So

$$\begin{aligned}
 \gamma_v &= \sum_{n=1}^{\infty} \sum_{w \in W} P_{w,v} \mathbb{P}_u(X_{n-1} = w, n \leq R_u) = \sum_{w \in V} P_{w,v} \left( \sum_{n=1}^{\infty} \mathbb{P}_u(X_{n-1} = w, n-1 < R_u) \right) \\
 &= \sum_{w \in V} P_{w,v} \sum_{n=0}^{\infty} \mathbb{P}_u(X_n = w, n < R_u) = \sum_{w \in V} P_{w,v} \sum_{n=0}^{\infty} \mathbb{E}_u[\mathbb{1}_{X_n=w} \mathbb{1}_{n < R_u}] \\
 &= \sum_{w \in V} P_{w,v} \mathbb{E}_u \left[ \sum_{n=0}^{R_u-1} \mathbb{1}_{X_n=w} \right] = \sum_{w \in V} P_{w,v} \gamma_w = (\gamma P)_v
 \end{aligned}$$

$\Rightarrow \gamma P = \gamma$ . Furthermore,  $\gamma_v > 0$ , since  $u \mapsto v$ . Letting  $n : P_{v,u}^n > 0$ , we have  $\gamma_u = (\gamma P^n)_u \geq \gamma_v P_{v,u}^n$ . Noting that  $\gamma_u = 1$ , this shows  $\gamma_v < \infty$ .  $\square$

**PROP 1.18** As a corollary, under the same conditions, if  $\mathbb{E}_u[R_u]$  is finite,  $\pi = (\pi_v : v \in V)$ , where

$$\pi_v = \frac{\gamma_v}{\sum_{w \in V} \gamma_w}$$

is a stationary distribution.

**PROOF.**

Observe

$$\sum_{w \in V} \gamma_w = \sum_{w \in V} \mathbb{E}_i \left[ \sum_{n=0}^{R_i-1} \mathbb{1}_{X_n=w} \right] = \mathbb{E}_i \left[ \sum_{n=0}^{R_i-1} \sum_{w \in V} \mathbb{1}_{X_n=w} \right] = \mathbb{E}_i[R_i]$$

So we require that  $\mathbb{E}_u[R_u] < \infty$ , then a stationary distribution exists.  $\square$

An exercise, if  $u$  is recurrent and  $u \mapsto v$ , then  $\mathbb{P}_u(T_v < \infty) = 1 \forall v \in V$ .

**PROP 1.19**  $\pi$ , as defined above, is a limiting distribution.

**PROOF.**

Consider two independent copies  $X_n, Y_n = \text{Markov}(P)$ . Then  $(X_n, Y_n)$  is a Markov chain with transition matrix  $Q = (q_{(v,w),(x,y)})_{(v,w),(x,y) \in V \times V}$ . In particular,

$$q_{(v,w),(x,y)} = p_{v,x} p_{w,y}$$

Fix some state  $u \in V$ . Let  $\alpha$  be the initial distribution that has  $X_0 = u$  and  $Y_0 \sim \pi$ .

$$\alpha_{(x,y)} = \begin{cases} \pi(y) & x = u \\ 0 & \text{o.w.} \end{cases}$$

Remark that, as  $\pi$  is stationary,  $Y_n \sim \pi \forall n \geq 0$ . For any  $u \in V$ , we'd like that  $P_{u,v}^n \rightarrow \pi(v)$ .

Let  $M = \inf\{n \geq 0 : X_n = Y_n\}$  be the first meeting time of the  $X_n$  and  $Y_n$  chains. If  $P$  is

finite and regular, then  $Q$  is finite and regular. Then

$$\mathbb{P}_\alpha(M < \infty) \geq \mathbb{P}_\alpha(T_{(v,v)} < \infty) = \sum_{(x,y) \in V \times V} \alpha(x,y) \mathbb{P}_{(x,y)}(T_{v,v} < \infty) = 1$$

It follows that  $\lim_{n \rightarrow \infty} \mathbb{P}_\alpha(M > n) = 0$ . We claim that  $\mathbb{P}_\alpha(X_n = v, M \leq n) = \mathbb{P}_\alpha(Y_n = v, M \leq n) \forall n \geq 0$ . Assuming this, then

$$\begin{aligned} \mathbb{P}_u(X_n = v) &= \mathbb{P}_\alpha(X_n = v) = \mathbb{P}_\alpha(X_n = v, M \leq n) + \mathbb{P}_\alpha(X_n = v, M > n) \\ \mathbb{P}_\pi(Y_n = v) &= \mathbb{P}_\alpha(Y_n = v) = \mathbb{P}_\alpha(Y_n = v, M \leq n) + \mathbb{P}_\alpha(Y_n = v, M > n) \\ \implies \mathbb{P}_u(X_n = v) - \pi(v) &= \mathbb{P}_\alpha(X_n = v, M > n) - \mathbb{P}_\alpha(Y_n = v, M > n) \\ \implies |\mathbb{P}_u(X_n = v) - \pi(v)| &= |\mathbb{P}_\alpha(X_n = v, M > n) - \mathbb{P}_\alpha(Y_n = v, M > n)| \leq \mathbb{P}_\alpha(M > n) \end{aligned}$$

But  $\mathbb{P}_\alpha(M > n) \rightarrow 0$ , so it must be that  $\mathbb{P}_u(X_n = v) \rightarrow \pi(v)$ .

We still must show the claim, however.

$$\begin{aligned} \mathbb{P}_\alpha(X_n = v, M \leq n) &= \sum_{k=0}^n \sum_{w \in V} \mathbb{P}_\alpha(X_n = v, M = k, X_k = w) \\ &= \sum_{k=0}^n \sum_{w \in V} \mathbb{P}_\alpha(M = k, X_k = w) \mathbb{P}_\alpha(X_n = v | X_k = w, M = k) \\ &\stackrel{\text{MP}}{=} \sum_{k=0}^n \sum_{w \in V} \mathbb{P}_\alpha(M = k, X_k = w) P_{w,v}^{n-k} \\ &\text{and now we reverse!} \\ &= \sum_{k=0}^n \sum_{w \in V} \mathbb{P}_\alpha(M = k, Y_k = w) P_{w,v}^{n-k} \\ &\stackrel{\text{MP}}{=} \sum_{k=0}^n \sum_{w \in V} \mathbb{P}_\alpha(M = k, Y_k = w) \mathbb{P}_\alpha(X_n = v | Y_k = w, M = k) \\ &= \mathbb{P}_\alpha(Y_n = v, M \leq n) \quad \square \end{aligned}$$

## HITTING TIMES AND ABSORBING STATES

We call a state  $v \in V$  *absorbing* if  $\mathbb{P}_v(X_1 = v) = 1$ .

DEF 1.26

**Eg. 1.12** Consider a game where we continually bet \$1 with winning probability  $p$ , until we earn \$ $k$  or run out of money. Modeling the state space as our balance, we have absorbing states at \$0 and \$ $k$ .

Observe that each absorbing state forms a recurrent communication class of size 1. As a consequence, we have a variant of the canonical form, where  $A \subseteq V$  are absorbing states,

and we write  $V = A \sqcup (V \setminus A)$ . For a stochastic matrix  $P$  and block matrices  $Q, R$ , we have:

$$P = \begin{array}{c|c} V \setminus A & A \\ \hline Q & R \\ 0 & I_{|A|} \end{array} \Rightarrow P^2 = \begin{array}{c|c} V \setminus A & A \\ \hline Q^2 & QR + RI \\ 0 & I_{|A|} \end{array}$$

$$\Rightarrow P^n = \begin{array}{c|c} V \setminus A & A \\ \hline Q^n & (I_{|V \setminus A|} + Q + \cdots + Q^{n-1})R \\ 0 & I_{|A|} \end{array}$$

Schematically, then

$$\begin{array}{ccc} \begin{pmatrix} Q \\ \downarrow \end{pmatrix} & & \begin{pmatrix} I_{|A|} \\ \downarrow \end{pmatrix} \\ V \setminus A & \xrightarrow{R} & A \end{array}$$

**PROP 1.20** Let  $V$  be finite. Then all states are either absorbing or transient.

**PROOF.**

Fix an initial distribution  $\alpha = (\alpha_v : v \in V)$  and an absorbing state  $c \in V$ . Then

$$\mathbb{E}_\alpha(T_c) = \sum_{v \in V} \mathbb{P}_\alpha(X_0 = v) \mathbb{E}_\alpha(T_c | X_0 = v) = \sum_{v \in V} \alpha_v \mathbb{E}_v(T_c) =$$

Recall from probability that  $\mathbb{E}X = \sum_{n \geq 1} \mathbb{P}(X \geq n)$  whenever  $X$  is supported on the non-negative integers. A pictorial proof is as follows

$$\begin{aligned} \mathbb{E}[X] &= \mathbb{P}(X = 1) \\ &+ \mathbb{P}(X = 2) + \mathbb{P}(X = 2) \\ &+ \mathbb{P}(X = 3) + \mathbb{P}(X = 3) + \mathbb{P}(X = 3) \\ &\vdots \\ &+ \mathbb{P}(X = n) + \mathbb{P}(X = n) + \mathbb{P}(X = n) + \cdots + \mathbb{P}(X = n) \\ &= \mathbb{P}(X \geq 1) + \mathbb{P}(X \geq 2) + \cdots + \mathbb{P}(X \geq n) \end{aligned}$$

Assume  $v$  is not absorbing. Then

$$\begin{aligned} \mathbb{E}_v(T_c) &= \sum_{n \geq 1} \mathbb{P}_v(T_c \geq n) = \sum_{n \geq 1} \sum_{w \neq c \in V} \mathbb{P}_v(X_{n-1} = w) = \sum_{n \geq 1} \sum_{w \neq c \in V} P_{v,w}^{n-1} \\ &= \sum_{n \geq 1} \sum_{w \in V \setminus A} Q_{w,v}^{n-1} + \sum_{n \geq 1} \sum_{w \neq c \in A} P_{v,w}^{n-1} \end{aligned}$$

where  $Q$  is as in the absorbing canonical form above. If there is an absorbing state  $b \neq c$  such that  $v \mapsto b$ , then  $\mathbb{E}_v(T_c) = \infty$ . Otherwise, the second sum above is 0, and we get

$$\mathbb{E}_v(T_c) = \sum_{w \in V \setminus A} \sum_{n \geq 1} Q_{v,w}^{n-1} = \sum_{w \in V \setminus A} \sum_{n \geq 0} Q_{v,w}^n$$

Here,  $I + Q + Q^2 + \cdots = (I - Q)^{-1}$ . □

Why is this invertible?  
Generally, if  $A^n \rightarrow 0$  as  $n \rightarrow \infty$ , then  $I - A$  is invertible.

**Theorem 1.7 Hitting State Properties**

Let  $P$  be a finite transition matrix, such that all states are either transient or absorbing. Let  $A$  be the absorbing states, with  $|A| = m$ . Then

$$\lim_{n \rightarrow \infty} P^n = \begin{bmatrix} 0 & (I - Q)^{-1}R \\ 0 & I_m \end{bmatrix}$$

Furthermore, for any transient states  $u, v$ ,

$$\mathbb{E}_u(L_v) = (I - Q)_{u,v}^{-1} \quad \text{and} \quad \mathbb{E}_u[T_A] = \sum_{v \in V} (I - Q)_{u,v}^{-1}$$

Finally, for any transient state  $u$  and absorbing state  $c$ ,

$$\mathbb{P}_u(X_{T_A} = c) = ((I - Q)^{-1}R)_{u,c}$$

**Eg. 1.13** Consider Gambler's Ruin, with reward limit  $k$ :



Then  $A = \{0, k\}$ . Let  $p_i = \mathbb{P}_i(X_{T_A} = k)$ . Computing manually, this is  $p_i = pp_{i+1} + (1-p)p_{i-1}$ , with boundary conditions  $p_0 = 0$  and  $p_k = 1$ . We conclude that

$$p_i = \begin{cases} \frac{1 - \left(\frac{1-p}{p}\right)^i}{1 - \left(\frac{1-p}{p}\right)^k} & p \neq \frac{1}{2} \\ \frac{i}{k} & p = \frac{1}{2} \end{cases}$$

Similarly, we manually compute  $\mathbb{E}_i(T_A) = 1 + p\mathbb{E}_{i+1}(T_A) + (1-p)\mathbb{E}_{i-1}(T_A)$ , with boundary conditions  $\mathbb{E}_0(T_A) = \mathbb{E}_k(T_A) = 0$ . This gives

$$\mathbb{E}_i(T_A) = \begin{cases} \frac{k(1-r^i) - i(1-r^k)}{p(1-r)(1-r^k)} & p \neq \frac{1}{2} \\ i(k-i) & p = \frac{1}{2} \end{cases} \quad \text{where } r = \frac{1-p}{p}$$

*Reversibility*

Some motivation: on an atomic scale, we cannot pin-point an "arrow of time," meaning, it is impossible to distinguish between a video on the atomic scale and that video in reverse. Entropy is hard to gauge! For instance, consider the following chains:



By symmetry, it must be that  $(\pi_1, \pi_2, \pi_3) = (1/3, 1/3, 1/3)$  is the stationary and limiting distribution for each chain. Can we tell if a "video" of the left chain is played in reverse? By symmetry, no. What about for the right video? We should observe "net counter-clockwise" behavior, as the clockwise transition from a given state only has probability  $1/3$ . Thus, we could distinguish between a video and its reverse.

Sometimes, it may only be possible to observe the arrow of time in a chain while close to initial conditions. The Ehrenfest Urn is a box with a partition down the middle, with particles initially on the left side of the box.

At each time step, we choose a random point in the box, and move it to its opposing side. When the urn reaches equilibrium, it will be impossible to distinguish an arrow of time. However, at the start, we expect left particles to move to the right.

These notions motivate the following:

DEF 1.27 Let  $P$  be a stochastic matrix with stationary distribution  $\pi$ . Then the *time reversal* of  $P$  is the matrix  $\hat{P}$  with

$$\pi_u \hat{P}_{u,v} = \pi_v P_{v,u}$$

In other words, after one time step, the probability of reaching state  $v$  from  $u$  in the time reversal is equal to the probability of reaching state  $u$  from  $v$  in the original chain.

### Theorem 1.8 Norris' Theorem

Let  $P$  be irreducible with stationary distribution  $\pi$ . Let  $X_0 \sim \pi$  and consider the chain  $X_n : n \in [N] = \text{Markov}(P)$ . Then  $(Y_0, \dots, Y_N) := (X_N, \dots, X_0)$  is  $\text{Markov}(\hat{P})$ .

PROOF.

Not provided in class. See textbook. □

**Fig. 1.14** Consider the Ehrenfest Urn example. Let  $X_n$  denote the number of particles on the left container at time  $n$ . Suppose that there are  $k$  total particles, such that  $V = [k]$ . Then

$$\mathbb{P}(X_{n+1} = i + 1 | X_n = i) = \frac{k - i}{k} \quad \mathbb{P}(X_{n+1} = i - 1 | X_n = i) = \frac{i - 1}{k}$$

What might a stationary distribution be? We could guess  $\text{Bin}(k, \frac{1}{2})$ , so that  $\pi(i) = \frac{1}{2^k} \binom{k}{i}$  (why does this make sense?). Checking,

$$\begin{aligned} \mathbb{P}_\pi(X_1 = i) &= \pi(i - 1) \mathbb{P}_{i-1}(X_1 = i) + \pi(i + 1) \mathbb{P}_{i+1}(X_1 = i) \\ &= \frac{1}{2^k} \left[ \binom{k}{i-1} \frac{k - (i - 1)}{k} + \binom{k}{i+1} \frac{i + 1}{k} \right] \\ &= \frac{1}{2^k} \left[ \frac{(k - 1)!}{i(k - i)!} \cdot i + \frac{(k - 1)!}{i!(k - 1)!} \cdot (k - 1) \right] = \frac{1}{2^k} \binom{k}{i} = \pi_i \end{aligned}$$

We conclude that  $X_0 \sim \pi \implies X_1 \sim \pi$ , so  $\pi P = \pi$ .

We can now compute the time reversal matrix, based on [Def 1.27](#):

$$\hat{P}_{i,i+1} = \frac{\pi_{i+1}}{\pi_i} P_{i+1,i} = \frac{\binom{k}{i+1}}{\binom{k}{i}} \frac{i+1}{k} = \frac{k-i}{k} = P_{i,i+1}$$

One shows a similar result for  $P_{i,i-1}$ .  $P = \hat{P}$ . In conclusion,  $P = \hat{P}$ .

A chain  $X = \text{Markov}(P)$  and its stochastic matrix  $P$  are called **reversible** if  $\hat{P} = P$ .

DEF 1.28

### Theorem 1.9 Reversible Chains are Random Weighted Walks

Let  $P$  be a stochastic matrix, with a stationary distribution  $\pi$ . Then  $P$  is reversible if and only if  $P$  can be represented as a random walk on an undirected weighted graph, i.e. there exists a graph  $G = (V, E)$  with weights  $w : E \rightarrow [0, \infty)$  such that

$$P_{u,v} = \frac{w(\{u, v\})}{\sum_{w \in N(u)} w(\{u, w\})} \quad \forall u, v \in V$$

( $\implies$ ) Let  $P$  be reversible with stationary distribution  $\pi$ . Let  $G = (V, E)$  have edge weights  $w(\{x, y\}) = \pi_x P_{x,y} = \pi_y P_{y,x}$ , by reversibility. Then, for  $u, v \in V$ ,

$$\frac{w(\{u, v\})}{\sum_{w \in N(u)} w(\{u, w\})} = \frac{\pi_u P_{u,v}}{\sum_{w \in N(u)} \pi_u P_{u,w}} = P_{u,v}$$

( $\impliedby$ ) Fix  $G = (V, E)$  with  $w : E \rightarrow [0, \infty)$ . Suppose  $P$  is given by a random walk on  $G$ :

$$P_{u,v} = \frac{w(\{u, v\})}{\sum_{w \in N(u)} w(\{u, w\})}$$

Then, for  $x \in V$ , a stationary distribution is provided by  $\pi(x) = \frac{\sum_{e \in \delta(x)} w(e)}{2 \sum_{e \in E} w(e)}$ . Notate  $D := 2 \sum_{e \in E} w(e)$ . For each  $u, v \in V$ ,

$$\begin{aligned} \pi_u P_{u,v} &= \frac{\sum_{e \in \delta(u)} w(e)}{D} \cdot \frac{w(\{u, v\})}{\sum_{w \in N(u)} w(\{u, w\})} = \frac{w(\{u, v\})}{D} \\ &= \frac{\sum_{e \in \delta(v)} w(e)}{D} \cdot \frac{w(\{v, u\})}{\sum_{e \in \delta(v)} w(e)} = \pi_v P_{v,u} \end{aligned}$$

Hence,  $P$  is reversible. □

If  $\pi$  satisfies  $\pi_u P_{u,v} = \pi_v P_{v,u} \quad \forall u, v \in V$ , then  $\pi$  is stationary for  $P$ .

PROP 1.21



PROOF.

Check that

$$\mathbb{P}_\pi(X_1 = v) = \sum_{u \in V} \pi_u P_{u,v} = \sum_{u \in V} \pi_v P_{v,u} = \pi_v \implies \pi P = \pi \quad \square$$

**Eg. 1.15** This class example is unrelated to our recent work. Consider a cipher of the alphabet, i.e. a bijection  $f : C \rightarrow A$ , where  $A = \{a, b, \dots, z\}$ , and  $C$  are a set of cipher symbols. From a collection of ciphered text, how can we determine  $f$ ?

One naive solution is to model the frequency of symbols in  $C$ , and "align" it with the frequency of the alphabet in English text. However, the mapping from letters to their frequency is not injective, so we cannot identify letters uniquely from their frequency.

Alternatively, we can let  $M : A \times A \rightarrow [0, 1]$  describe the frequency of transitions, i.e.

$M(x, y)$  = proportion of " $x$ " that are followed by a " $y$ " in English text

For instance,  $M(s, t)$  may be large, but  $M(x, j) = 0$ . Then for a given text  $s = (s_1, \dots, s_L)$  of cipher characters, and a hypothesized mapping  $f$ ,

$$\text{Plausibility}(f) = \prod_{i=1}^{L-1} M(f(s_i), f(s_{i-1}))$$

## INDEX OF DEFINITIONS

absorbing 1.26  
accessible 1.14  
aperiodic 1.19  
  
canonical decomposition 1.24  
closed 1.25  
communicate 1.15  
communication class 1.16  
conditional density 1.1  
  
hitting time 1.7  
  
initial distribution 1.9  
irreducible 1.17  
  
law of large numbers 1.10  
limiting distribution 1.12  
local time 1.23  
  
Markov chain 1.2  
Markov property 1.4  
  
period 1.18  
  
recurrent 1.21  
regular 1.13  
return time 1.20  
reversible 1.28  
  
state space 1.3  
stationary 1.11  
stochastic matrix 1.8  
  
time reversal 1.27  
time-homogeneous 1.5  
transient 1.22  
transition matrix 1.6