

## MSH ED's Intra-MSHS Transfer Criteria for Oncology Patients

Level 1	
Intra-MSHS Transfer Protocol	Clinical Criteria
Unless an oncology attending states otherwise, the ED clinician can transfer the patient and communicate to the patient that the primary oncologist approves of the transfer	<p>Solid tumor patient with a non-primary oncologic or non-acute primary oncologic reason for admission who does not meet any of the exclusion criteria in Level 3.</p> <p>Non-primary and non-acute primary oncologic reasons include, but are not limited to:</p> <ul style="list-style-type: none"> <li>○ Cardiac dysrhythmias</li> <li>○ Chest pain</li> <li>○ CHF</li> <li>○ COPD exacerbation</li> <li>○ Diabetes mellitus with complications</li> <li>○ Failure to thrive</li> <li>○ Fractures not related to mets</li> <li>○ Infections (including sepsis, UTI, PNA, viral, etc.)</li> <li>○ Skin and subcutaneous tissue infections</li> <li>○ Symptom management due to disease but not chemotherapy (including pain, nausea, vomiting, diarrhea, and constipation)</li> </ul>
Level 2	
Intra-MSHS Transfer Protocol	Clinical Criteria
<p>Will need sign-off from primary oncologist before transfer</p> <p><i>If primary oncologist is unavailable, solid tumor oncology patients can be escalated to Dr. Cardinale Smith and malignant hematology patients to Dr. Luis Isola</i></p>	<p>Any cancer patient with primary oncologic reason for admission who does not meet any of the exclusion criteria in Level 3.</p> <p>Primary oncologic reasons include, but are not limited to:</p> <ul style="list-style-type: none"> <li>○ Complications directly related to active chemotherapy, for example neutropenic sepsis</li> <li>○ Complications directly related to oncologic diagnosis, for example pathologic fracture</li> <li>○ Newly suspected metastasis referred for management</li> </ul>
Level 3	
Intra-MSHS Transfer Protocol	Clinical Criteria
Not suitable for transfer and should be flagged for immediate admission at MSH	<ul style="list-style-type: none"> <li>● New acute leukemia or other new malignant hematology diagnosis</li> <li>● Research patient</li> <li>● Patient who requires titration of vents, increased vent requirements, frequent suctioning</li> <li>● Patient who received a transplant (Allo/Auto/CAR-T) within last year</li> <li>● Patient acutely ill with unstable vital signs and severe, symptomatic lab abnormalities or cancer sequelae presenting as: <ul style="list-style-type: none"> <li>○ Acute kidney failure</li> <li>○ Acute liver failure</li> <li>○ Acute mental status change/neurologic symptoms</li> <li>○ Cord Compression</li> <li>○ Cytokine release syndrome</li> <li>○ Disseminated intravascular coagulation</li> <li>○ GI bleed requiring repeat Hgb checks more often than Q8 hours</li> <li>○ GVHD</li> <li>○ Hyperviscosity syndrome</li> <li>○ Leptomenigeal disease</li> <li>○ Multifocal PNA with SpO2 &lt;90% on room air</li> <li>○ Tumor lysis syndrome</li> </ul> </li> </ul>