



Remodulin Guidelines

Step by Step:

IV Remodulin Priming Technique

- Close roller clamp.
- Spike IV bag and hang.
- Squeeze and release middle of drip chamber until it is approximately 2/3 full.
- Open roller clamp to prime **slowly**.
- Close clamp below filter after priming.
- Program pump settings.

Initial Set up - IV Remodulin administration

- Hand Hygiene
- Scrub hub, attach syringe and withdraw until a flash of blood is visible. Place syringe to the side. (Amount in syringe is catheter volume).
- Scrub hub, flush line with 10 ml of Normal Saline; place curo on port.
- Calculate re-priming volume by multiplying catheter volume by 0.8.
- Scrub port of IV bag with alcohol, withdraw re-priming volume from IV bag.
- Scrub hub, attach syringe to IV and **slowly** re-prime.
- Scrub hub, attach new tubing to patient and start infusion.

Change of Concentration - IV Remodulin administration

To be performed after priming technique completed for new concentration bag

- Hand Hygiene.
- Disconnect patient from old IV concentration infusion.
- Scrub hub, attach syringe and withdraw until a flash of blood is visible. Place syringe to the side. (Amount in syringe is catheter volume).
- **Scrub hub, attach 3 ml syringe and withdraw 3 ml of blood waste. Discard Syringe.**
- Scrub hub, flush line with 10 ml of Normal Saline.
- Calculate re-priming volume by multiplying catheter volume by 0.8
- Scrub port of new IV bag with alcohol; withdraw re-priming volume from new IV bag.
- Scrub hub, attach syringe to IV port and **slowly** re-prime catheter volume.
- Scrub hub, attach new tubing to patient and start infusion.