

## THE PRESENT AND FUTURE

### JACC STATE-OF-THE-ART REVIEW

# Care Models for Acute Chest Pain That Improve Outcomes and Efficiency



## JACC State-of-the-Art Review

Luke P. Dawson, MBBS, MPH,<sup>a,b,c</sup> Karen Smith, PhD,<sup>a,d,e</sup> Louise Cullen, MBBS, PhD,<sup>f</sup> Ziad Nehme, PhD,<sup>d,e</sup> Jeffrey Lefkowitz, MBBS,<sup>a,b</sup> Andrew J. Taylor, MBBS, PhD,<sup>a,c</sup> Dion Stub, MBBS, PhD<sup>a,c,d</sup>

### ABSTRACT

Existing assessment pathways for acute chest pain are often resource-intensive, prolonged, and expensive. In this review, the authors describe existing chest pain pathways and current issues at the patient and system level, and provide an overview of recent advances in chest pain research that could inform improved outcomes for both patients and health systems. There are multiple avenues to improve existing models of chest pain care, including novel risk stratification pathways incorporating highly sensitive point-of-care troponin assays; new devices available before first medical contact that could allow clinicians to access vital signs and electrocardiogram data; artificial intelligence and precision medicine tools that may guide indications for further testing; and strategies to improve hospital benchmarking and performance monitoring to standardize care. Improving the speed and accuracy of chest pain diagnosis and management should be a priority for researchers and is likely to translate to substantive benefits for patients and health systems. (J Am Coll Cardiol 2022;79:2333-2348) © 2022 by the American College of Cardiology Foundation.

Acute chest pain is common and can be caused by a broad range of conditions ranging from benign to life-threatening. To exclude serious and life-threatening conditions, such as acute coronary syndromes (ACS), emergency department (ED) and hospital assessment is frequently required and can be resource-intensive, prolonged, and expensive. Although discussion commonly focusses on improving disease-based outcomes, recent chest pain guidelines have highlighted the importance of monitoring and improving symptom-based care, which provides a more comprehensive representation of health care provision at the patient level.<sup>1</sup> In this review, we aimed to: 1) describe existing chest pain

pathways and current problems at a patient and system level, such as hospital overcrowding, high costs, and overtesting; and 2) provide an overview of recent advances in chest pain research that could result in improved acute chest pain assessment models and outcomes for both patients and health systems.

### ACUTE CHEST PAIN EPIDEMIOLOGY AND OUTCOMES

Acute chest pain accounts for approximately 5%-10% of attendances to EDs and 10% of calls for assistance to emergency medical services (EMS), with >7 million presentations to ED annually in the United States.<sup>2-4</sup>



Listen to this manuscript's audio summary by Editor-in-Chief Dr Valentin Fuster on [www.jacc.org/journal/jacc](http://www.jacc.org/journal/jacc).

From the <sup>a</sup>Department of Epidemiology and Preventive Medicine, Monash University, Melbourne, Victoria, Australia; <sup>b</sup>Department of Cardiology, The Royal Melbourne Hospital, Melbourne, Victoria, Australia; <sup>c</sup>Department of Cardiology, The Alfred Hospital, Melbourne, Victoria, Australia; <sup>d</sup>Ambulance Victoria, Melbourne, Victoria, Australia; <sup>e</sup>Department of Paramedicine, Monash University, Melbourne, Victoria, Australia; and the <sup>f</sup>Emergency and Trauma Centre, Royal Brisbane and Women's Hospital, Brisbane, Queensland, Australia.

The authors attest they are in compliance with human studies committees and animal welfare regulations of the authors' institutions and Food and Drug Administration guidelines, including patient consent where appropriate. For more information, visit the [Author Center](#).

Manuscript received February 11, 2022; revised manuscript received March 30, 2022, accepted March 30, 2022.

## ABBREVIATIONS AND ACRONYMS

**ACS** = acute coronary syndromes

**CDP** = clinical decision pathway

**CMR** = cardiac magnetic resonance imaging

**CPU** = chest pain unit

**CT** = computed tomography

**CTCA** = computed tomography coronary angiography

**ECG** = electrocardiogram

**ED** = emergency department

**EMS** = emergency medical services

**MACE** = major adverse cardiac events

**MI** = myocardial infarction

**ncRNA** = noncoding RNA

**STEMI** = ST-segment elevation myocardial infarction

Demand for EMS services for chest pain is growing faster than overall EMS demand and population growth,<sup>5</sup> and lifetime prevalence of acute chest pain has been estimated at 20%-40%.<sup>6</sup> Rates of acute chest pain presentations are marginally higher among women than men.<sup>2</sup> A significant socioeconomic gradient has been described in some studies, with higher incidences of acute chest pain attendances observed among socially disadvantaged cohorts.<sup>7,8</sup> Racial and ethnic differences have been observed, including a higher incidence of acute chest pain among Black and Hispanic patients in the United States.<sup>9-11</sup>

The underlying causes of acute chest pain are broad, and after assessment, one-half of all patients are eventually discharged with a diagnosis of nonspecific pain.<sup>3,4</sup> Nonetheless, life-threatening or serious causes are present in a significant proportion of presentations (Figure 1).<sup>12</sup> Although differences are

observed across different cohorts and populations, around 25% have a cardiovascular diagnosis, whereas 20%-25% have a noncardiovascular diagnosis.<sup>4</sup> Most common cardiovascular diagnoses include acute coronary syndromes (10%), congestive heart failure (5%), and arrhythmias (5%). Noncardiovascular diagnoses include pneumonia or exacerbations of chronic obstructive pulmonary disease (10%) and gastrointestinal disorders, especially gastroesophagitis (5%). Pulmonary emboli (1%) and acute aortic pathologies (0.3%) are rare but are important conditions to exclude due to high mortality rates.<sup>3</sup>

Patient outcomes are dependent on the underlying cause and can vary from mortality rates of <1% for nonspecific pain, to 20% for acute aortic pathologies (Figure 1).<sup>12</sup> Overall mortality rates for acute chest pain presentations are approximately 1%-2%, with substantial variation according to patient demographics.<sup>4,12</sup> For patients discharged with an index admission diagnosis of nonspecific chest pain, readmission for serious cardiovascular events including ACS is approximately 3% within 6 months.<sup>13</sup> One-year mortality among patients diagnosed with nonspecific pain is similar to that of the general population.<sup>14,15</sup>

## EXISTING MODELS OF ACUTE CHEST PAIN CARE

To avoid missing serious or life-threatening causes of acute chest pain, current guidelines recommend patients or bystanders contact EMS for rapid transfer to the nearest ED for further assessment.<sup>1</sup> Although


## HIGHLIGHTS

- Acute chest pain is the most common reason for emergency medical contact, but existing assessment pathways for acute chest pain are resource intensive, lengthy, and expensive.
- Advances in point-of-care troponin assays, wearable devices, shared decision-making, noninvasive imaging strategies, artificial intelligence, and big data applications promise to improve management of patients with chest pain.
- Further development and application of these tools could reduce the system burden of chest pain assessment and improve clinical outcomes.

rapid transfer to hospital remains the priority, the role of EMS has expanded in recent years to include more complex assessment such as 12-lead electrocardiograms (ECGs) and prenotification in ST-segment elevation myocardial infarction (STEMI). Initial ED assessment involves a focused history and examination, combined with targeted investigations that usually include an ECG and troponin sampling, using a stepwise approach aimed at excluding the most serious conditions first. Much of this process is focused on identifying the risk of ACS, with high-risk patients usually admitted and frequently investigated with coronary angiography, whereas intermediate-risk patients are typically investigated with functional or noninvasive anatomical testing, and low-risk patients triaged to no testing or optional testing strategy for coronary artery disease.<sup>1</sup> Existing tools demonstrating improved clinical or system outcomes in assessment and management of acute chest pain are summarized in the following section.

**PREHOSPITAL CARE.** Thirty percent to 60% of chest pain patients present via ambulance, and several interventions in recent decades have generated improvements in prehospital care for acute chest pain.<sup>16,17</sup> Prehospital ECGs are associated with reduced mortality in undifferentiated chest pain cohorts,<sup>18</sup> possibly through early diagnosis of STEMI, with hospital prenotification and often earlier cardiac catheterization laboratory activation.<sup>19-21</sup> Contacting EMS directly is associated with improved door-to-balloon-times for STEMI,<sup>22</sup> and public health campaigns aimed at educating the community to contact EMS when acute chest pain occurs have similarly been successful in improving public engagement with

**FIGURE 1** Diagnoses and Mortality Among Acute Chest Pain Cohorts




Most common specific diagnoses and 30-day mortality rates among acute chest pain attendances by emergency medical services according to age and sex. Nonspecific pain according to male and female sex accounts for 52.1% and 50.0% in the 18-49-year age group, 47.2% and 53.8% in the 50-69-year age group, and 37.1% and 41.6% in the ≥70-year age group. Internal data from Ambulance Victoria, 2015 to 2019.<sup>12</sup> AF = atrial fibrillation; COPD = chronic obstructive pulmonary disease; GORD = gastroesophageal reflux disease; NSTEACS = non-ST-segment elevation acute coronary syndromes; SIHD = stable ischemia heart disease; STEMI = ST-segment elevation myocardial infarction; SVT = supraventricular tachycardia.

EMS in some jurisdictions.<sup>23</sup> Despite these improvements, uptake of prehospital ECGs can still be improved, with rates approximately 70% among chest pain cohorts.<sup>18</sup> Similarly, delays in ambulance off-load times to overcrowded EDs can be improved. These delays have been associated with worsened clinical outcomes and have been a concern in the United Kingdom, the United States, Canada, and Australia.<sup>24,25</sup>

**TROPONIN PROTOCOLS.** Contemporary troponin assays were developed in the 1990s and substantially improved diagnosis of myocardial infarction (MI) for patients presenting with acute chest pain. However, contemporary cardiac troponin assays were limited by low sensitivity for MI at initial presentation, requiring prolonged periods of testing over 6-12 hours, delaying diagnosis among patients with MI, and delaying discharge among patients

without MI (Figure 2).<sup>26-28</sup> Advancements in assay technology led to the development of highly sensitive cardiac troponin assays, allowing shorter periods of testing between serial troponin samplings.<sup>29</sup> Coupled with greater understanding regarding the incremental information provided by serial troponin results if troponin assays are within the normal range,<sup>30</sup> recent studies have demonstrated the safety of 3-hour,<sup>31</sup> 2-hour,<sup>32</sup> and subsequently, 1-hour rapid troponin pathways,<sup>32-34</sup> leading to improvements in hospital assessment times (Figure 3).<sup>35-37</sup> However, implementation of guideline-recommended rapid troponin pathways is not universal in clinical practice, and hospital admission times can be variable even with institution of such testing protocols, with a median ED length of stay of 2.5 to 4.6 hours among low-risk patients in the 0/1-hour arm of recent trials.<sup>37-39</sup>

**FIGURE 2** Temporal Trends in Hospital Assessment Times for Acute Chest Pain




Progression in hospital assessment times for acute chest pain presentations over the last 2 decades in major randomized clinical trials and observational studies. Year of study recruitment is shown on the x-axis in comparison to median hospital assessment times for acute chest pain presentations. Hospital times represent those for patients classified as low-risk of index acute coronary syndrome following initial assessment. Each point represents a study with colors representing the protocol used for that patient cohort (legend). Further data regarding these studies are detailed in [Supplemental Table 1](#). ADAPT = 2-Hour Accelerated Diagnostic Protocol to Assess Patients With Chest Pain Symptoms Using Contemporary Troponins as the Only Biomarker; APACE = Advantageous Predictors of Acute Coronary Syndrome Evaluation; CHEER = Chest Pain Evaluation in the Emergency Room; FASTEST = Fast Assessment of Thoracic Pain in the Emergency Department Using High-Sensitive Troponins and a Simple Risk Score; Henry Ford HEART = Henry Ford History, ECG, Age, Risk factors, and initial Troponin; HISTORIC = High-Sensitivity Cardiac Troponin on Presentation to Rule Out Myocardial Infarction; hsTn = highly sensitive troponin; ROMICAT = Rule Out Myocardial Infarction using Computer Assisted Tomography; ROMICAT-II = Rule Out Myocardial Infarction using Computer Assisted Tomography II; ROMIO = The Rapid Rule-Out of Myocardial Ischemia Observation; TRAPID-AMI = The High Sensitivity Cardiac Troponin T Assay for Rapid Rule-out of Acute Myocardial Infarction.

**EXISTING RISK STRATIFICATION TOOLS AND CLINICAL DECISION PATHWAYS.** Existing risk stratification tools were developed before the availability of highly sensitive troponin assays, with the most common scores including the HEART (History, ECG, Age, Risk factors, and initial Troponin) score, EDACS (Emergency Department Assessment of Chest Pain Score) score, and TIMI (Thrombolysis In Myocardial Infarction) score. In combination with contemporary troponin assays, multiple studies have demonstrated improved sensitivity in detecting 30-day major adverse cardiac events (MACE), and therefore, these tools are recommended when highly sensitive troponin assays are not available.<sup>1</sup> Although these scores have also been validated in the highly sensitive troponin era, their benefit over protocols that use

highly sensitive troponin assays alone is less clear, with several studies demonstrating no incremental value in diagnostic classification when such risk scores are added to highly sensitive troponin protocols.<sup>40,41</sup>

Protocolization of ED assessment for patients with acute chest pain through the use of Clinical Decision Pathways (CDPs) have been shown to reduce ED length of stay and rates of unnecessary testing in several studies. Many CDPs are based on risk stratification tools, whereas others, such as the 2020 European Society of Cardiology (ESC) guidelines, use highly sensitive troponin testing alone. The aim of most of these pathways is to classify patients into low-, intermediate-, and high-risk groups, with low-risk groups indicating a risk of ACS within 30 days of  $\leq 1\%$ .<sup>1,30</sup> Several CDPs, including the standard



and modified HEART pathways,<sup>42,43</sup> the ADAPT (2-Hour Accelerated Diagnostic Protocol to Assess Patients With Chest Pain Symptoms Using Contemporary Troponins as the Only Biomarker) pathway,<sup>36</sup> and the EDACS-ADP pathway<sup>44</sup> have demonstrated consistent reductions of 20%-45% in the need for admission in patients presenting with suspected ACS. These pathways are intended for use once alternate serious diagnoses, such as pneumonia, decompensated heart failure, acute aortic pathologies, and pulmonary emboli, have been excluded.

**CHEST PAIN UNITS AND CLINICS.** Chest pain units (CPUs) and emergency short stay units have been used to streamline assessment and management of patients with acute chest pain while awaiting serial troponin testing and risk stratification.<sup>45</sup> Germany has been especially proactive in the development and

accreditation of a CPU network, which now covers almost the entire country with 336 certified CPUs, including a CPU registry to facilitate quality assurance and research.<sup>46,47</sup> Guidelines for the setup and management of CPUs have been published by the Acute Cardiovascular Care Association and the German Society of Cardiology detailing the organizational structure, physical requirements, technical requirements, and management of patients within the CPU.<sup>45,47</sup> Multiple studies performed during the contemporary troponin assay era, including randomized trials, demonstrated that CPUs improved clinical outcomes, reduced admission rates and costs, and improved patient satisfaction.<sup>48-50</sup> Some CPUs incorporate functional testing such as stress echocardiography, which has been shown to be highly feasible,<sup>51</sup> but available data have not demonstrated

improvements in rates of postdischarge 30-day MACE, in patients undergoing noninvasive testing in CPUs before discharge.<sup>52</sup> Moreover, some studies have identified CPU overuse of functional testing in patients with very low pretest probability.<sup>53</sup> German CPU networks has generally recommended and used separate units within the ED with predefined continuously available beds for chest pain patients.<sup>45,54</sup> In other jurisdictions, the organizational and physical CPU structures have been more varied but often incorporate CPU assessment processes into the main ED without defined beds as long as technical facilities including cardiac monitoring are available. In the United States and some other countries, American College of Cardiology chest pain accreditation and certification is widely used, which is focused on quality improvement and performance monitoring rather than specific CPU organizational structures.<sup>54,55</sup> Rapid troponin testing protocols have reduced rates of admission to CPUs,<sup>56,57</sup> and in this setting, locating CPUs within the ED department may make more logistic sense for centers without established separate CPUs. Similarly, CPU location within the ED may expedite further evaluations if a noncardiac diagnosis is present. Rapid-access chest pain clinics can be an alternative or complementary service for further diagnostic workup and the facilitation of noninvasive testing for appropriate patients at low to intermediate risk of future coronary events, and are associated with earlier diagnosis, reduced unplanned ED reattendances within 30 days of discharge, and reduced overall costs.<sup>58,59</sup> Following ED assessment, postdischarge follow-up with a cardiologist is associated with improved outcomes at 1 year.<sup>60</sup>

### CHEST PAIN EVALUATION IN THE COVID-19 ERA

The COVID-19 pandemic has led to substantial challenges in chest pain management requiring consideration of a broader set of differential diagnoses and modifications at each stage of standard chest pain care processes.<sup>61</sup> Acute COVID-19 infection can present with symptoms of chest pain, often of a pleuritic or inflammatory characteristics, which can persist for months following the acute infection—with a prevalence of 22% at 2 months after presentation in 1 study.<sup>62</sup> Acute myocarditis, defined as cardiac symptoms with raised highly sensitive troponin and abnormal ECG, cardiac magnetic resonance imaging (CMR), echocardiogram, or histopathology, is a well-described complication of COVID-19.<sup>61</sup> Rates of COVID-19-related myocarditis vary widely according

to definitions, but occur in 0.5%–7.0% of patients based on CMR and autopsy studies.<sup>63,64</sup> Rates of myocarditis complicated by shock are estimated to be higher than that of non-COVID-19 viral myocarditis.<sup>65</sup> Similarly, thromboembolic complications, specifically pulmonary emboli, can occur and may present with chest pain. Patients with cardiovascular symptoms and COVID-19 should undergo comprehensive evaluation with ECG, highly sensitive troponin, and transthoracic echocardiogram testing, and with CMR recommended if initial testing is abnormal.<sup>61</sup> Ischemic testing should be performed in accordance with existing chest pain clinical guidelines.<sup>1</sup>

The subsequent development of COVID-19 vaccines has similarly resulted in concerns regarding postvaccine chest pain presentations, especially regarding risks of myocarditis following mRNA-based vaccinations. Post-mRNA vaccine myocarditis incidence varies according to age and sex, with an estimated incidence of 41 per million vaccinations among males aged <30 years and 4 per million vaccinations among females aged <30 years.<sup>66</sup> The prognosis from vaccine-related myocarditis is generally very good with >95% having a mild clinical course. Nonspecific chest pain and generalized musculoskeletal symptoms without biomarker or imaging evidence of myopericarditis is significantly more common following COVID-19 vaccination.<sup>67</sup>

**CHALLENGES AND MODIFICATIONS TO CHEST PAIN MANAGEMENT DURING COVID-19.** At the prehospital level, COVID-19 has resulted in reduced ambulance availability, saturation of emergency call lines leading to delays in dispatch, and reduced system efficiencies.<sup>68</sup> Substantial adaptations to infection control measures, including paramedic personal protective equipment, transfer processes to EDs, decontamination of ambulances, and the segregation of hospital resources, all contribute to lost efficiency.<sup>69</sup> At the peak of the pandemic, most jurisdictions instituted full personal protective equipment and N95 masks for paramedics for all attendances, resulting in increased complexity of patient retrieval, delays to EMS response times,<sup>70</sup> and greater difficulties with standard chest pain assessment processes such as prehospital ECGs. Delays to first-ECG, diagnosis, and door-to-balloon times were observed in several studies of patients with STEMI.<sup>71</sup> Screening questions, such as the presence of fever, dyspnea, or risk factors for infection, are often used to select patients that require isolation until a COVID-19 test.<sup>72</sup> However, overlap between chest pain and these symptoms in addition to concerns regarding COVID-19-related causes of chest pain means that many chest pain presentations still require initial isolation.



Follow-up for chest pain presentations more commonly occurred via telehealth, which is limited by the absence of in-person environment and physical examination.<sup>73</sup> Finally, patient presentation rates for many conditions, including ACS, were observed to decrease during periods of high community prevalence in the setting of patient concerns regarding overburdening health systems under strain or the risk of contracting COVID-19 while in hospital.<sup>74</sup>

Many of the challenges with chest pain care models in the setting of COVID-19 have improved as community prevalence has decreased with widespread vaccine availability. Access to rapid antigen testing may have reduced some delays to evaluation and management. Modifications to existing chest pain CDPs that consider the need for COVID-19 testing and isolation precautions are worthwhile to streamline care models and prioritize early ECG for rapid identification of STEMI.<sup>75</sup> Similarly, public health education to encourage patients with acute chest pain to contact EMS or present to EDs might allay some fears regarding presentation.

#### SYSTEM AND PATIENT IMPACTS OF EXISTING ASSESSMENT PATHWAYS

The patient and system impacts of assessment pathways are summarized in this section and include high financial costs among patients often classified as low risk, system overcrowding and delays in treatment and diagnosis for other patients, the impact of prolonged assessment and admission times on patients, and harms relating to overinvestigation (Figure 4).


**FINANCIAL COSTS OF ACUTE CHEST PAIN CARE.** Each aspect of acute chest pain assessment—including prehospital transfer (often requiring specialized staff), emergency assessments and subsequent cardiac investigations, and management for high- or intermediate-risk patients—is associated with costs. In 2015, an Australian study estimated mean ED and admission costs for acute chest pain attendances at AU\$13,509 (US\$9,854) for patients diagnosed with ACS, AU\$7,283 (US\$5,314) for other cardiovascular conditions, and AU\$3,331 (US\$2,430) for noncardiac conditions,<sup>76</sup> with similar estimates in other jurisdictions.<sup>77,78</sup> In the United States, mean costs in 2016 per acute chest pain attendance to ED were estimated at US\$6,325, with a total annual cost of US\$1.5 billion.<sup>79</sup> Rapid testing protocols and adherence to chest pain pathway protocols are associated with reductions in hospital costs.<sup>80-83</sup> In 1 study, the introduction of an accelerated diagnostic protocol across 16 hospitals in Australia resulted in a AU\$13.5 million (US\$9.8 million) saving through a 20%

reduction in ED length of stay (AU\$2.3 million saving) and a 13% reduction in hospital admissions (AU\$11.2 million saving).<sup>84</sup> The introduction of highly sensitive troponin assays has been associated with improved early rule-out processes, reduced need for functional testing, and a mean 20% reduction in costs.<sup>85</sup> Noninvasive anatomical or functional tests are a significant driver of high costs. However, despite these costs, some studies have suggested that early use of imaging in intermediate-risk patients may be cost effective by facilitating earlier discharge and reduced subsequent invasive angiography.<sup>86-90</sup> Benefits of early noninvasive testing in low-risk patients are less clear, especially in the setting of highly sensitive troponin assays.<sup>91,92</sup>

**IMPACTS ON HEALTH SYSTEMS.** Acute chest pain attendances are frequently associated with rates of hospital admission ranging from 40%-70%.<sup>38,39</sup> Although assessment times are improving with highly sensitive troponin assays and protocolized chest pain pathways, the burden on systems remains significant. Overcrowding in EDs in the setting of hospital access block has become increasingly common in many jurisdictions and is associated with increased costs, delays to treatment, worsened patient outcomes, cancellation of elective procedures, delays in ambulance off-load times, and in turn, delays to ambulance response times.<sup>91,93-95</sup> Acute chest pain, accounting for 10% of ED attendances, remains a significant driver of this problem, and even small improvements in ED or hospital admission times have the potential to reduce overcrowding and improve overall health care quality.<sup>36,96</sup>

**RISKS OF OVERINVESTIGATION AND OVERTRIAGE.** Low-risk categorization in acute chest pain patients generally aims to represent an ACS “miss” rate of  $\leq 1\%$ , consistent with the expectations of  $>50\%$  of clinicians that ACS diagnostic strategies should achieve a sensitivity of 99% or higher.<sup>97</sup> However, increased sensitivity needs to be balanced against the risk of harms associated with false-positive testing and overinvestigation, sometimes termed the test threshold.<sup>98</sup> In the setting of contemporary troponin assays, this has been estimated at 2% (ie, it is estimated to be worthwhile undertaking troponin testing if the pretest probability is  $>1$  in 50).<sup>98</sup> Conversely, indiscriminate troponin testing can lead to diagnostic uncertainty and unnecessary investigations.<sup>99</sup> Similarly indiscriminate noninvasive imaging investigations for low-risk patients with chest pain leads to increased costs and system burden, higher rates of invasive angiography (and associated patient risks), with little evidence to suggest improvements

**FIGURE 4** Problems With Existing Acute Chest Pain Care Models



**Left panel** shows the structure of existing acute chest pain care models including rates of presentation and final etiology. The **right panel in blue** identifies problems with existing models of care in regard to patient care and outcomes, and the burden on health systems. ACS = acute coronary syndrome; CPU = chest pain unit; ECG = electrocardiogram; ED = emergency department; EMS = emergency medical services; STEMI = ST-segment elevation myocardial infarction.

in diagnosis rates of coronary artery disease or improvements on outcomes.<sup>92,100,101</sup> Furthermore, in 1 study, higher clinician-specific hospitalization rates for acute chest pain were not associated with improvements in 30-day outcomes, suggesting that an overly conservative approach to acute chest pain management resulting in a lower threshold for admission is not necessarily safer.<sup>102</sup>

### IMPROVING ACUTE CHEST PAIN MODELS OF CARE


Novel research into improving models of acute chest pain evaluation is of great importance in both improving patient outcomes and reducing the burden and costs of lengthy acute chest pain assessment processes. There is substantial scope for improvements in speed of diagnosis and risk stratification methods given that the vast majority of patients are eventually safely discharged home. This section focuses on developing research that might safely improve both the speed of care and tools available

to diagnose and manage patients with chest pain without increasing the risk of adverse events (**Central Illustration**).

**NOVEL TECHNOLOGIES FOR RISK STRATIFICATION BEFORE MEDICAL CONTACT.** Just as parts of usual ED care are being shifted to prehospital EMS care, similar developments in technologies might provide patients and clinicians with clinical information before arrival of medical personnel. Smartwatch and mobile devices that can monitor arrhythmias, including atrial fibrillation, have seen significant advances in the last decade, and research in this area has expanded to assessment of ischemia.<sup>103</sup> Home ECG monitoring devices, such as the AliveCor device,<sup>104</sup> have demonstrated usefulness in detecting ischemia and acute ST changes.<sup>105</sup> Nine-lead ECGs are obtainable with some smartwatches, including the Apple Watch, by placing the smart watch on different positions on the body, and have been used to detect ischemia with comparable results to standard ECGs.<sup>106,107</sup> Similarly, smart phone and watch tools to



## CENTRAL ILLUSTRATION Improving Acute Chest Pain Models of Care



Dawson LP, et al. *J Am Coll Cardiol.* 2022;79(23):2333-2348.

Novel areas of research and development that might translate to improved patient and system outcomes for acute chest pain presentations. AI = artificial intelligence; ECG = electrocardiogram; EMS = emergency medical services.

detect arrhythmias have been approved by the U.S. Food and Drug Administration, which also may provide useful information to patients and clinicians before medical contact, given acute chest pain relates to arrhythmias in a proportion of cases.

Developments in smart devices also include clinical observations, such as pulse oximetry, blood pressure, and heart rate, with similar accuracy to commercial purpose-specific devices.<sup>108</sup> With these developments, it might be feasible for patients with acute chest pain to contact EMS and transmit a full set of observations, and ECG rhythm and ischemia data to assist in early risk stratification and triage decisions, and prioritize EMS attendance to patients with life-threatening conditions such as STEMI. However, it is important to note that for existing devices, variability between devices and between activity types have been observed, with 1 study demonstrating an average error rate 30% higher during activity compared with rest for wearable heart rate sensors.<sup>109</sup>

**EMS RISK STRATIFICATION AND PREHOSPITAL POINT-OF-CARE TROPONIN ASSAYS.** Improvements in prehospital care of acute chest pain, including the use of ECGs and prenotification for STEMI, have

resulted in improved outcomes among chest pain cohorts and these are now established standards of care.<sup>18</sup> Success with these interventions has prompted research into further avenues for improvement, including the use of prehospital risk stratification, use of point-of-care troponin assays, and improved ECG algorithms.

Several studies have assessed the feasibility of incorporating paramedic-based risk assessments using risk stratification scores, that might ordinarily be used by clinicians in EDs, often in combination with point-of-care troponin sampling. Overall, the process of troponin sampling and risk score calculation (eg, HEART score) has been shown to be feasibly performed by paramedics and is associated with reduced ED length of stay.<sup>110-116</sup> Studies to date have been performed using contemporary troponin assays, which may not safely rule out ACS,<sup>115</sup> but highly sensitive point-of-care troponin assays are in development by several companies and likely to be available for use in the prehospital setting in the near future. For the 30%-60% of patients arriving to ED with acute chest pain via EMS, routine use of risk stratification and point-of-care highly sensitive troponin sampling would allow patients to arrive to

ED with a completed HEART score, potentially allowing early discharge for low-risk patients within the first hour after a second troponin is sampled or consideration of alternate diagnoses. Conversely, high-risk patients with an initially elevated troponin could assist in guiding prehospital management, especially aspirin administration, intravenous access, transport urgency decisions, and decisions to directly transport patients to catheterization-capable centers, avoiding interhospital transfers. Similarly, on arrival to ED, an elevated troponin result would assist in guiding ED triage categorization, and decisions regarding early evaluation and management, and might facilitate earlier decisions regarding disposition from the ED. Some studies have suggested paramedic risk stratification and point-of-care troponin results could occur at the patient's home with only patients at intermediate or high risk transferred to hospital,<sup>116</sup> but this may be somewhat optimistic without a risk assessment process to exclude other serious conditions such as pulmonary emboli, pneumonia, and acute aortic pathologies. Further trials assessing these processes are planned.<sup>117,118</sup>

**RISK STRATIFICATION MODELS, BIG DATA, AND MACHINE LEARNING.** Existing risk stratification scores for acute chest pain are mostly variations on simple counting tools, similar to the CHA<sub>2</sub>DS<sub>2</sub>-VASc score for stroke risk in atrial fibrillation, which facilitate ease of use clinically. However, these tools do not quantify absolute risk in the way that more complex models incorporating more variables do, such as the QStroke score, which demonstrates substantially improved discrimination for stroke risk in AF patients compared with the CHA<sub>2</sub>DS<sub>2</sub>-VASc score, or the numerous mortality predictive models used for risk adjustment in cardiovascular procedural registries.<sup>119,120</sup> Electronic medical records systems are now widespread and provide opportunities to incorporate more complex models into standard clinical processes, which might more accurately quantify risk in comparison to simple counting tools such as the HEART score.<sup>121,122</sup> Greater availability of large linked clinical and administrative datasets provide similar opportunities for developing enhanced risk prediction models. Moreover, predictive models that can stratify risk of serious non-ACS causes of chest pain, may assist in determining the need for non-coronary investigations and safety of early discharge among patients with normal serial highly sensitive troponin assays.

Machine learning and artificial intelligence methods have been well-publicized across clinical medicine in recent years, and several studies have

demonstrated potentially promising applications in prediction modeling, disease phenotyping, image interpretation (including ECGs and radiology), and precision medicine.<sup>123</sup> In stable chest pain, a phenomapping-derived tool was used in the PROMISE (PROspective Multicenter Imaging Study for Evaluation of Chest Pain) and SCOT-HEART (Scottish COmputed Tomography of the HEART Trial) populations to select phenotypic neighborhoods favoring anatomical or functional testing, demonstrating a lower incidence of MACE using the tool's recommended testing strategy.<sup>124</sup> Deep learning-based artificial intelligence algorithms have demonstrated excellent discrimination for diagnosing MI using 6- or 12-lead ECGs,<sup>125</sup> with improvements in sensitivity of 52% compared with commercial interpretation software and 37% compared with experienced clinicians in 1 study.<sup>126,127</sup> Multiple studies have assessed the use of artificial intelligence in predicting diagnosis or outcomes in chest pain cohorts using artificial neural networks, random forest, support vector machine, and gradient boosting methods, and although these have generally outperformed clinicians and existing risk stratification tools, few have yet been incorporated into practice.<sup>128,129</sup> Similarly, although these methods potentially show promise, whether performance in chest pain predictive modeling improves on more conventional statistical methods such as logistic regression is not yet clear.<sup>130,131</sup>

**NOVEL BIOMARKERS.** Troponin assays predominate assessment of patients with suspected ACS, but several novel biomarkers are potentially promising and warrant mention, including cardiac myosin-binding protein C (cMyC) and noncoding RNAs (ncRNAs). cMyC, a sarcomeric protein associated with myosin and actin, can be detected earlier in blood and rises more rapidly in comparison to troponin.<sup>132</sup> In undifferentiated chest pain patients presenting to ED, cMyC at presentation provided comparable discriminatory power (area under the curve: 0.924) in comparison to a single highly sensitive troponin assay, and led to improved discrimination when combined with highly sensitive troponin assays.<sup>133</sup> ncRNAs include microRNAs, circular RNAs, and long non-coding RNAs, and although sensitivity of current ncRNAs is inferior to troponin assays, research into potential combined biomarkers using miRNA and cMyC is ongoing.<sup>134,135</sup> Furthermore, some biomarkers such as copeptin, midregional proatrial natriuretic peptide, C-terminal proendothelin-1, midregional proendrenomedullin, and procalcitonin may assist in differentiating type 2 MI from type 1 MI in chest pain cohorts.<sup>136,137</sup>

Perhaps of greater importance in the near future are improvements in point-of-care devices and troponin assays. Rapid point-of-care highly sensitive troponin assays that provide results in <15 minutes have been developed but are not yet widely available for clinical use.<sup>138</sup> Such assays are likely to improve ED length of stay, especially using the 0/1-hour troponin pathway, which would be able to classify patients as low risk with increased speed compared with awaiting results from current, slower, lab-based assays.

**UPFRONT NONINVASIVE IMAGING STRATEGIES.** A complete discussion of noninvasive imaging strategies in ED and index admission of patients with acute chest pain is outside the scope of this review, but numerous studies have assessed the impact of different imaging strategies on patient outcomes and cost effectiveness. Upfront strategies (following troponin and ECG but before invasive angiography) for several modalities, including computed tomography (CT) coronary angiography (CTCA), have been associated with improved cost effectiveness in acute chest pain cohorts.<sup>139</sup> The ACRIN-PA (CT Coronary Angiogram Versus Traditional Care in Emergency Department Assessment of Potential ACS) and ROMICAT-II (Multicenter Study to Rule Out Myocardial Infarction by Cardiac Computed Tomography) trials demonstrated that CTCA reduced ED length of stay and increased ED discharge rates in comparison to inpatient functional testing.<sup>140,141</sup> However, a routine noninvasive imaging strategy before discharge has not been demonstrated to improve early outcomes, and in the ROMICAT-II trial, was associated with increased downstream testing and radiation exposure.<sup>142</sup> Moreover, the BEACON (Better Evaluation of Acute Chest Pain with Computed Tomography Angiography) trial, which was performed in the highly sensitive troponin era, found that although overall costs were lower due to reduced outpatient testing, there was no added benefit of additional testing over highly sensitive troponin assays in identifying significant coronary disease.<sup>143</sup> Therefore, the use of these tests needs to be carefully selected and may be most suited to intermediate-risk rather than low-risk patients.<sup>91,92,139</sup>

Although the use of imaging in low- and intermediate-risk patients has been extensively studied, potential new applications for developing techniques might include upfront imaging to assess selected high-risk patients and those with elevated troponin values. Invasive angiography has been the gold-standard investigation for high-risk patients for over 2 decades and allows treatment with

percutaneous coronary intervention at the time if required. However, up to 50% of patients undergoing angiography for non-STEMI do not require revascularization, and 5%-10% are better managed with bypass surgery—both groups that might avoid complications associated with invasive angiography if noninvasive workup could be done safely and accurately without delaying revascularization.<sup>144</sup> An initial imaging-guided strategy in non-STEMI was associated with a 44% reduction in invasive angiography use for patients initially undergoing CTCA and an 13% reduction in patients undergoing CMR in comparison to conventional angiography first pathways, with no increase in 1-year MACE.<sup>145</sup> Similar findings were identified in the VERDICT (Very Early Versus Deferred Invasive Evaluation Using Computerized Tomography in Patients With Acute Coronary Syndromes) trial, which demonstrated high diagnostic accuracy to rule out clinically significant coronary disease using upfront CTCA in patients with non-ST-segment elevation acute coronary syndromes, potentially avoiding unnecessary invasive testing in some patients.<sup>146</sup> Nonetheless, invasive angiography remains the best approach for high-risk patients, and further large studies would be required before a shift toward noninvasive strategies could be recommended. Similarly, further data assessing whether CT imaging (including fractional flow reserve and perfusion) can be used to accurately guide revascularization decisions, rather than simply acting as a gate-keeping test to angiography, would also be required.

**PATIENT-CLINICIAN SHARED DECISION-MAKING.** Clinician concern regarding missed diagnoses often results in a risk averse approach to chest pain workup including admission to CPUs and further diagnostic testing of low-risk patients. This can result in increased downstream procedures and higher costs without necessarily improving outcomes. A growing body of evidence has demonstrated that patient-clinician shared decision-making results in greater patient education, lower rates of CPU admission, lower rates of unnecessary diagnostic testing, and is not associated with higher rates of MACE.<sup>147,148</sup> Shared decision-making aids are useful across a variety of decision contexts, and can reinforce the importance of outpatient follow-up, effectively educate patients regarding their individual risk, and facilitate timely discharge.<sup>149</sup> Recent chest pain guidelines recommend decision aids for both low-risk patients to facilitate risk communication, and intermediate-risk patients to assist in shared decisions surrounding further inpatient or outpatient

investigations and the need for admission, observation, or discharge.<sup>1</sup>

**BENCHMARKING AND MONITORING HOSPITAL PERFORMANCE.** Clinical quality registries have proliferated over the last 2 decades, and many countries and localities now monitor hospital and operator performance in cardiovascular procedures through such registries.<sup>150</sup> In the United States, the Chest Pain-MI registry has been developed to standardize chest pain and MI hospital-based care,<sup>151</sup> but few registries using a symptom-based rather than disease-based (ie, MI or ACS) approach have been developed elsewhere. Chest pain center accreditation is offered by the American College of Cardiology in the United States with the goal of helping hospitals to identify and improve care for patients with suspected ACS, with reaccreditation required every 3-5 years, which may be an effective mechanism for improving outcomes.<sup>152,153</sup> This accreditation process has been taken up widely in the United States in addition to some other countries, including China.<sup>55</sup> As availability of electronic record systems, administrative databases, and linkage processes improve, benchmarking and monitoring of hospital (and prehospital) performance by measuring chest pain-specific key performance indicators and risk-adjusted outcomes (as is done for percutaneous coronary intervention registries) might assist in standardizing hospital care and identifying outlier hospitals where care processes can be improved.<sup>151</sup> A similar approach might also be applicable to avoid excess costs and overtesting among low-risk chest pain presentations. For example, risk-adjusted key performance indicators might be developed using linked dataset that monitor institutional costs, rates of imaging, and adherence to CDPs for acute chest pain presentations.

## CONCLUSIONS

Patients with acute chest pain are common and current assessment pathways frequently include prolonged ED and hospital admissions with a focus on exclusion of ACS with high certainty, contributing to health care costs, hospital overcrowding, and sometimes over-investigation. There are multiple avenues to improve existing models of chest pain evaluation including novel risk stratification pathways incorporating highly sensitive point-of-care troponin assays, in both the hospital and prehospital setting; public education surrounding new technologies such as smartwatches' ongoing development into indications for further testing including the use of developing artificial intelligence and precision medicine methods; increased uptake of CDPs, CPUs, and accreditation programs; and improved hospital benchmarking and performance monitoring for acute chest pain attendances to standardize care. Improving the speed and accuracy of chest pain diagnosis and management is likely to translate to significant benefits for patients and health systems.

## FUNDING SUPPORT AND AUTHOR DISCLOSURES

Dr Dawson is supported by National Health and Medical Research Council of Australia (NHMRC) and National Heart Foundation (NHF) postgraduate scholarships. Dr Nehme is supported by an NHMRC/NHF early career fellowship. Dr Stub is supported by an NHF Fellowship (#105793). Dr Taylor is supported by an NHMRC Investigator grant. The authors have reported that they have no relationships relevant to the contents of this paper to disclose.

**ADDRESS FOR CORRESPONDENCE:** Dr Dion Stub, Department of Cardiology, Alfred Health, 55 Commercial Road, Prahran, Victoria 3004, Australia. E-mail: [d.stub@alfred.org.au](mailto:d.stub@alfred.org.au).

## REFERENCES

- Gulati M, Levy PD, Mukherjee D, et al. 2021 AHA/ACC/AASE/CHEST/SAEM/SCCT/SCMR guideline for the evaluation and diagnosis of chest pain: executive summary: a report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *J Am Coll Cardiol*. 2021;78(22):2218-2261. <https://doi.org/10.1016/j.jacc.2021.07.052>
- National Center for Health Statistics. National Hospital Ambulatory Medical Care Survey: 2018 Emergency Department Summary Tables. 2018. Accessed January 15, 2022. [https://www.cdc.gov/nchs/data/nhamcs/web\\_tables/2018-ed-web-tables-508.pdf](https://www.cdc.gov/nchs/data/nhamcs/web_tables/2018-ed-web-tables-508.pdf)
- Hsia RY, Hale Z, Tabas JA. A national study of the prevalence of life-threatening diagnoses in patients with chest pain. *JAMA Intern Med*. 2016;176:1029-1032.
- Pedersen CK, Stengaard C, Friesgaard K, et al. Chest pain in the ambulance; prevalence, causes and outcome - a retrospective cohort study. *Scand J Trauma Resusc Emerg Med*. 2019;27:84.
- Andrew E, Nehme Z, Cameron P, Smith K. Drivers of increasing emergency ambulance demand. *Prehosp Emerg Care*. 2020;24:385.
- Ruigómez A, Rodríguez LA, Wallander MA, Johansson S, Jones R. Chest pain in general practice: incidence, comorbidity and mortality. *Fam Pract*. 2006;23:167-174.
- Walters K, Rait G, Hardoon S, Kalaitzaki E, Petersen I, Nazareth I. Socio-demographic variation in chest pain incidence and subsequent coronary heart disease in primary care in the United Kingdom. *Eur J Prev Cardiol*. 2014;21:566-575.
- Dawson LP, Andrew E, Nehme Z, et al. Association of socioeconomic status with outcomes and care quality in patients presenting with undifferentiated chest pain in the setting of universal health care coverage. *J Am Heart Assoc*. 2022;11(7):e024923. <https://doi.org/10.1161/JAHA.121.024923>
- Alrwan A, Eworuke E. Are discrepancies in waiting time for chest pain at emergency departments between African Americans and whites improving over time? *J Emerg Med*. 2016;50:349-355.
- Graham G. Racial and ethnic differences in acute coronary syndrome and myocardial infarction within the United States: from demographics to outcomes. *Clin Cardiol*. 2016;39:299-306.
- López L, Wilper AP, Cervantes MC, Betancourt JR, Green AR. Racial and sex differences in emergency department triage assessment

and test ordering for chest pain, 1997-2006. *Acad Emerg Med*. 2010;17:801-808.

12. Andrew E, Cox S, Smith K. Linking ambulance records with hospital and death index data to evaluate patient outcomes. *Int J Gen Med*. 2022;15:567-572.

13. Kwok CS, Brown DL, Van Spall HGC, et al. Non-specific chest pain and subsequent serious cardiovascular readmissions. *Int J Cardiol*. 2019;291:1-7.

14. Fagring AJ, Lappas G, Kjellgren KI, Welin C, Manhem K, Rosengren A. Twenty-year trends in incidence and 1-year mortality in Swedish patients hospitalised with non-AMI chest pain. Data from 1987-2006 from the Swedish hospital and death registries. *Heart*. 2010;96:1043-1049.

15. Ilankovan N, Mickle H, Diederichsen A, et al. Clinical features and prognosis of patients with acute non-specific chest pain in emergency and cardiology departments after the introduction of high-sensitivity troponins: a prospective cohort study. *BMJ Open*. 2017;7:e018636.

16. Mnataganian G, Hiller JE, Braitberg G, et al. Sex disparities in the assessment and outcomes of chest pain presentations in emergency departments. *Heart*. 2020;106:111-118.

17. Van Severen E, Willemsen R, Vandervoort P, Sabbe M, Dinant GJ, Buntinx F. How do patients with chest pain access emergency department care? *Eur J Emerg Med*. 2017;24:423-427.

18. Rawshani N, Rawshani A, Gelang C, et al. Association between use of pre-hospital ECG and 30-day mortality: a large cohort study of patients experiencing chest pain. *Int J Cardiol*. 2017;248:77-81.

19. Jin J, Yu J, Wang J. Impact of improved pre-hospital emergency medical service system on the time management of chest pain patients in the emergency department. *Am J Transl Res*. 2021;13:7743-7755.

20. Anroedh SS, Kardys I, Akkerhuis KM, et al. e-Transmission of ECGs for expert consultation results in improved triage and treatment of patients with acute ischaemic chest pain by ambulance paramedics. *Neth Heart J*. 2018;26:562-571.

21. Patel M, Dunford JV, Aguilar S, et al. Pre-hospital electrocardiography by emergency medical personnel: effects on scene and transport times for chest pain and ST-segment elevation myocardial infarction patients. *J Am Coll Cardiol*. 2012;60:806-811.

22. Fernández-Bergés D. Data from the German Chest Pain Unit Registry: the well known gap between knowledge and practice. *Int J Cardiol*. 2017;246:16-17.

23. Nehme Z, Cameron PA, Akram M, et al. Effect of a mass media campaign on ambulance use for chest pain. *Med J Aust*. 2017;206:30-35.

24. Kingswell C, Shaban RZ, Crilly J. Concepts, antecedents and consequences of ambulance ramping in the emergency department: a scoping review. *Australas Emerg Nurs J*. 2017;20:153-160.

25. Li M, Vanberkel P, Carter AJE. A review on ambulance offload delay literature. *Health Care Manag Sci*. 2019;22:658-675.

26. Morrow DA, Antman EM, Tanasijevic M, et al. Cardiac troponin I for stratification of early outcomes and the efficacy of enoxaparin in unstable angina: a TIMI-11B substudy. *J Am Coll Cardiol*. 2000;36:1812-1817.

27. Macrae AR, Kavsak PA, Lustig V, et al. Assessing the requirement for the 6-hour interval between specimens in the American Heart Association Classification of Myocardial Infarction in Epidemiology and Clinical Research Studies. *Clin Chem*. 2006;52:812-818.

28. Kavsak PA, MacRae AR, Lustig V, et al. The impact of the ESC/ACC redefinition of myocardial infarction and new sensitive troponin assays on the frequency of acute myocardial infarction. *Am Heart J*. 2006;152:118-125.

29. Reichlin T, Hochholzer W, Bassetti S, et al. Early diagnosis of myocardial infarction with sensitive cardiac troponin assays. *N Engl J Med*. 2009;361:858-867.

30. Neumann JT, Twerenbold R, Ojeda F, et al. Application of high-sensitivity troponin in suspected myocardial infarction. *N Engl J Med*. 2019;380:2529-2540.

31. Wildi K, Nelles B, Twerenbold R, et al. Safety and efficacy of the 0 h/3 h protocol for rapid rule out of myocardial infarction. *Am Heart J*. 2016;181:16-25.

32. Boeddinghaus J, Twerenbold R, Nestelberger T, et al. Clinical validation of a novel high-sensitivity cardiac troponin I assay for early diagnosis of acute myocardial infarction. *Clin Chem*. 2018;64:1347-1360.

33. Twerenbold R, Neumann JT, Sörensen NA, et al. Prospective validation of the 0/1-h algorithm for early diagnosis of myocardial infarction. *J Am Coll Cardiol*. 2018;72:620-632.

34. Neumann JT, Sörensen NA, Schwemer T, et al. Diagnosis of myocardial infarction using a high-sensitivity troponin I 1-hour algorithm. *JAMA Cardiol*. 2016;1:397-404.

35. Bandstein N, Ljung R, Lundbäck M, Johansson M, Holzmann MJ. Trends in admissions for chest pain after the introduction of high-sensitivity cardiac troponin T. *Int J Cardiol*. 2017;240:1-7.

36. Than M, Aldous S, Lord SJ, et al. A 2-hour diagnostic protocol for possible cardiac chest pain in the emergency department: a randomized clinical trial. *JAMA Intern Med*. 2014;174:51-58.

37. Twerenbold R, Costabel JP, Nestelberger T, et al. Outcome of applying the ESC 0/1-hour algorithm in patients with suspected myocardial infarction. *J Am Coll Cardiol*. 2019;74:483-494.

38. Chew DP, Lambrakis K, Blyth A, et al. A randomized trial of a 1-hour troponin t protocol in suspected acute coronary syndromes: the Rapid Assessment of Possible Acute Coronary Syndrome in the Emergency Department With High-Sensitivity Troponin T Study (RAPID-TnT). *Circulation*. 2019;140:1543-1556.

39. Stoyanov KM, Hund H, Biener M, et al. RAPID-CPU: a prospective study on implementation of the ESC 0/1-hour algorithm and safety of

discharge after rule-out of myocardial infarction. *Eur Heart J Acute Cardiovasc Care*. 2020;9:39-51.

40. Fanaroff AC, Rymer JA, Goldstein SA, Simel DL, Newby LK. Does this patient with chest pain have acute coronary syndrome? The rational clinical examination systematic review. *JAMA*. 2015;314:1955-1965.

41. Khan E, Lambrakis K, Blyth A, et al. Classification performance of clinical risk scoring in suspected acute coronary syndrome beyond a rule-out troponin profile. *Eur Heart J Acute Cardiovasc Care*. 2021;10(9):1038-1047. <https://doi.org/10.1093/ehjacc/zuab040>

42. Mahler SA, Riley RF, Hiestand BC, et al. The HEART Pathway randomized trial: identifying emergency department patients with acute chest pain for early discharge. *Circ Cardiovasc Qual Outcomes*. 2015;8:195-203.

43. McCord J, Cabrera R, Lindahl B, et al. Prognostic utility of a modified HEART score in chest pain patients in the emergency department. *Circ Cardiovasc Qual Outcomes*. 2017;10(2):e003101. <https://doi.org/10.1161/CIRCOUTCOMES.116.003101>

44. Than MP, Pickering JW, Aldous SJ, et al. Effectiveness of EDACS versus ADAPT accelerated diagnostic pathways for chest pain: a pragmatic randomized controlled trial embedded within practice. *Ann Emerg Med*. 2016;68:93-102.e1.

45. Claeys MJ, Ahrens I, Sinnaeve P, et al. Editor's choice-the organization of chest pain units: position statement of the Acute Cardiovascular Care Association. *Eur Heart J Acute Cardiovasc Care*. 2017;6:203-211.

46. Münzel T, Heusch G. Chest pain unit network in Germany: its effect on patients with acute coronary syndromes. *J Am Coll Cardiol*. 2017;69:2459-2460.

47. Post F, Gori T, Giannitsis E, et al. Criteria of the German Society of Cardiology for the establishment of chest pain units: update 2014. *Clin Res Cardiol*. 2015;104:918-928.

48. Goodacre S, Nicholl J, Dixon S, et al. Randomised controlled trial and economic evaluation of a chest pain observation unit compared with routine care. *BMJ*. 2004;328:254.

49. Goodacre SW, Quinney D, Revell S, Morris F, Capewell S, Nicholl J. Patient and primary care physician satisfaction with chest pain unit and routine care. *Acad Emerg Med*. 2004;11:827-833.

50. Keller T, Post F, Tzikas S, et al. Improved outcome in acute coronary syndrome by establishing a chest pain unit. *Clin Res Cardiol*. 2010;99:149-155.

51. Shah BN, Balaji G, Alhajiri A, Ramzy IS, Ahmadvazir S, Senior R. Incremental diagnostic and prognostic value of contemporary stress echocardiography in a chest pain unit: mortality and morbidity outcomes from a real-world setting. *Circ Cardiovasc Imaging*. 2013;6:202-209.

52. Howell SJ, Prasad P, Vipparla NS, Venugopal S, Amsterdam EA. Usefulness of predischARGE cardiac testing in low risk women and men for safe, rapid discharge from a chest pain unit. *Am J Cardiol*. 2019;123:1772-1775.



53. Jade J, Huggan P, Stephenson D. Who gets admitted to the Chest Pain Unit (CPU) and how do we manage them? Improving the use of the CPU in Waikato DHB, New Zealand. *BMJ Qual Improv Rep*. 2015;4(1). <https://doi.org/10.1136/bmjquality.201506670.w2735>
54. Breuckmann F, Burt DR, Melching K, et al. Chest pain centers: a comparison of accreditation programs in Germany and the United States. *Crit Pathw Cardiol*. 2015;14:67-73.
55. Hao Y, Liu J, Liu J, et al. Rationale and design of the Improving Care for Cardiovascular Disease in China (CCC) project: a national effort to prompt quality enhancement for acute coronary syndrome. *Am Heart J*. 2016;179:107-115.
56. Husain I, Mahler SA, Hiestand BC, Miller CD, Stoppyra JP. The impact of accelerated diagnostic protocol implementation on chest pain observation unit utilization. *Crit Pathw Cardiol*. 2021;21(1):7-10. <https://doi.org/10.1097/HPC.0000000000000254>
57. Terry N, Franks N, Moran T, Pitts S, Osborne A, Ross MA. The changing role of chest pain in the emergency department observation unit. *Crit Pathw Cardiol*. 2021;20:119-125.
58. Black JA, Cheng K, Flood JA, et al. Evaluating the benefits of a rapid access chest pain clinic in Australia. *Med J Aust*. 2019;210:321-325.
59. Kozor R, Mooney J, Lowe H, et al. Rapid access chest pain clinics: an Australian cost-benefit study. *Heart Lung Circ*. 2022;31(2):177-182. <https://doi.org/10.1016/j.hlc.2021.05.106>
60. Czarnecki A, Chong A, Lee DS, et al. Association between physician follow-up and outcomes of care after chest pain assessment in high-risk patients. *Circulation*. 2013;127:1386-1394.
61. Gluckman T, Bhavne N, Allen LA, et al. 2022 ACC expert consensus decision pathway on cardiovascular sequelae of COVID-19 in adults: myocarditis and other myocardial involvement, post-acute sequelae of SARS-CoV-2 infection, and return to play: a report of the American College of Cardiology Solution Set Oversight Committee. *J Am Coll Cardiol*. 2022;79(17):1717-1756.
62. Carfi A, Bernabei R, Landi F. Persistent symptoms in patients after acute COVID-19. *JAMA*. 2020;324:603-605.
63. Halushka MK, Vander Heide RS. Myocarditis is rare in COVID-19 autopsies: cardiovascular findings across 277 postmortem examinations. *Cardiovasc Pathol*. 2021;50:107300.
64. Moulson N, Petek BJ, Drezner JA, et al. SARS-CoV-2 cardiac involvement in young competitive athletes. *Circulation*. 2021;144:256-266.
65. Rathore SS, Rojas GA, Sondhi M, et al. Myocarditis associated with Covid-19 disease: a systematic review of published case reports and case series. *Int J Clin Pract*. 2021;75:e14470.
66. Gargano JW, Wallace M, Hadler SC, et al. Use of mRNA COVID-19 vaccine after reports of myocarditis among vaccine recipients: update from the Advisory Committee on Immunization Practices - United States, June 2021. *MMWR Morb Mortal Wkly Rep*. 2021;70:977-982.
67. Kadali RAK, Janagama R, Peruru S, Malayala SV. Side effects of BNT162b2 mRNA COVID-19 vaccine: a randomized, cross-sectional study with detailed self-reported symptoms from healthcare workers. *Int J Infect Dis*. 2021;106:376-381.
68. Nadarajah R, Gale CP. Collateral cardiovascular damage during the COVID-19 pandemic. *Nat Rev Cardiol*. 2022;19:81-82.
69. Fernandez AR, Crowe RP, Bourn S, et al. COVID-19 preliminary case series: characteristics of EMS encounters with linked hospital diagnoses. *Prehosp Emerg Care*. 2021;25:16-27.
70. Andrew E, Nehme Z, Stephenson M, Walker T, Smith K. The impact of the COVID-19 pandemic on demand for emergency ambulances in Victoria, Australia. *Prehosp Emerg Care*. Published online on July 16, 2021.
71. McLaren JTT, Taher AK, Chartier LB. Flattening the other curve: Reducing emergency department STEMI delays during the COVID-19 pandemic. *Am J Emerg Med*. 2021;49:367-372.
72. Quah LJ, Tan BKK, Fua TP, et al. Reorganising the emergency department to manage the COVID-19 outbreak. *Int J Emerg Med*. 2020;13:32.
73. Bloom JE, Dawson LP, Seneviratne Epa D, et al. Perceptions of telehealth and cardiac care during the COVID-19 pandemic. *Intern Med J*. 2021;51:151-152.
74. De Filippo O, D'Ascenzo F, Angelini F, et al. Reduced rate of hospital admissions for ACS during Covid-19 outbreak in Northern Italy. *N Engl J Med*. 2020;383:88-89.
75. Huber K, Goldstein P. Covid-19: implications for prehospital, emergency and hospital care in patients with acute coronary syndromes. *Eur Heart J Acute Cardiovasc Care*. 2020;9:222-228.
76. Cullen L, Greenslade J, Merollini K, et al. Cost and outcomes of assessing patients with chest pain in an Australian emergency department. *Med J Aust*. 2015;202:427-432.
77. van Dongen DN, Ottervanger JP, Tolsma R, et al. In-hospital healthcare utilization, outcomes, and costs in pre-hospital-adjudicated low-risk chest-pain patients. *Appl Health Econ Health Policy*. 2019;17:875-882.
78. Vester MPM, Eindhoven DC, Bonten TN, et al. Utilization of diagnostic resources and costs in patients with suspected cardiac chest pain. *Eur Heart J Qual Care Clin Outcomes*. 2021;7(6):583-590. <https://doi.org/10.1093/ehjqcco/qcaa064>
79. Aalam AA, Alsabbab A, Pines JM. National trends in chest pain visits in US emergency departments (2006-2016). *Emerg Med J*. 2020;37:696-699.
80. Bhatti Y, Stevenson A, Weerasuriya S, Khan S. Reducing avoidable chest pain admissions and implementing high-sensitivity troponin testing. *BMJ Open Qual*. 2019;8:e000629.
81. Riley RF, Miller CD, Russell GB, et al. Cost analysis of the History, ECG, Age, Risk factors, and initial Troponin (HEART) Pathway randomized control trial. *Am J Emerg Med*. 2017;35:77-81.
82. Cheng Q, Greenslade JH, Parsonage WA, et al. Change to costs and lengths of stay in the emergency department and the Brisbane protocol: an observational study. *BMJ Open*. 2016;6:e009746.
83. Jülicher P, Greenslade JH, Parsonage WA, Cullen L. The organisational value of diagnostic strategies using high-sensitivity troponin for patients with possible acute coronary syndromes: a trial-based cost-effectiveness analysis. *BMJ Open*. 2017;7:e013653.
84. Parsonage WA, Milburn T, Ashover S, et al. Implementing change: evaluating the Accelerated Chest pain Risk Evaluation (ACRE) project. *Med J Aust*. 2017;207:201-205.
85. Twerenbold R, Jaeger C, Rubini Gimenez M, et al. Impact of high-sensitivity cardiac troponin on use of coronary angiography, cardiac stress testing, and time to discharge in suspected acute myocardial infarction. *Eur Heart J*. 2016;37:3324-3332.
86. Papachristidis A, Vaughan GF, Denny SJ, et al. Comparison of NICE and ESC proposed strategies on new onset chest pain and the contemporary clinical utility of pretest probability risk score. *Open Heart*. 2020;7(1):e001081. <https://doi.org/10.1136/openhrt-2019-001081>
87. Bertoldi EG, Stella SF, Rohde LEP, Polanczyk CA. Cost-effectiveness of anatomical and functional test strategies for stable chest pain: public health perspective from a middle-income country. *BMJ Open*. 2017;7:e012652.
88. Chrysant GS, Fillmore S, McClain R. A cardiac computed tomography first strategy to evaluate chest pain in a rural setting: outcomes and cost implications. *Coron Artery Dis*. 2019;30:413-417.
89. Hulten E, Goehler A, Bittencourt MS, et al. Cost and resource utilization associated with use of computed tomography to evaluate chest pain in the emergency department: the Rule Out Myocardial Infarction using Computer Assisted Tomography (ROMICAT) study. *Circ Cardiovasc Qual Outcomes*. 2013;6:514-524.
90. Karády J, Mayrhofer T, Ivanov A, et al. Cost-effectiveness analysis of anatomic vs functional index testing in patients with low-risk stable chest pain. *JAMA Netw Open*. 2020;3:e2028312.
91. Foy AJ, Liu G, Davidson WR Jr, Sciamanna C, Leslie DL. Comparative effectiveness of diagnostic testing strategies in emergency department patients with chest pain: an analysis of downstream testing, interventions, and outcomes. *JAMA Intern Med*. 2015;175:428-436.
92. Sandhu AT, Heidenreich PA, Bhattacharya J, Bundorf MK. Cardiovascular testing and clinical outcomes in emergency department patients with chest pain. *JAMA Intern Med*. 2017;177:1175-1182.
93. Pines JM, Pollack CV Jr, Diercks DB, Chang AM, Shofer FS, Hollander JE. The association between emergency department crowding and adverse cardiovascular outcomes in patients with chest pain. *Acad Emerg Med*. 2009;16:617-625.
94. Sprivilis PC, Da Silva JA, Jacobs IG, Frazer AR, Jelinek GA. The association between hospital overcrowding and mortality among patients admitted via Western Australian emergency departments. *Med J Aust*. 2006;184:208-212.
95. Hoot NR, Aronsky D. Systematic review of emergency department crowding: causes, effects, and solutions. *Ann Emerg Med*. 2008;52:126-136.



96. Vigen R, Diercks DB, Hashim IA, et al. Association of a novel protocol for rapid exclusion of myocardial infarction with resource use in a US safety net hospital. *JAMA Netw Open*. 2020;3:e203359.
97. Than M, Herbert M, Flaws D, et al. What is an acceptable risk of major adverse cardiac event in chest pain patients soon after discharge from the emergency department?: a clinical survey. *Int J Cardiol*. 2013;166:752-754.
98. Kline JA, Johnson CL, Pollack CV Jr, et al. Pretest probability assessment derived from attribute matching. *BMC Med Inform Decis Mak*. 2005;5:26.
99. Shah ASV, Sandoval Y, Noaman A, et al. Patient selection for high sensitivity cardiac troponin testing and diagnosis of myocardial infarction: prospective cohort study. *BMJ*. 2017;359:j4788.
100. Hermann LK, Newman DH, Pleasant WA, et al. Yield of routine provocative cardiac testing among patients in an emergency department-based chest pain unit. *JAMA Intern Med*. 2013;173:1128-1133.
101. Carpenter CR, Raja AS, Brown MD. Over-testing and the downstream consequences of overtreatment: implications of "preventing over-diagnosis" for emergency medicine. *Acad Emerg Med*. 2015;22:1484-1492.
102. Natsui S, Sun BC, Shen E, et al. Higher emergency physician chest pain hospitalization rates do not lead to improved patient outcomes. *Circ Cardiovasc Qual Outcomes*. 2021;14:e006297.
103. Perez MV, Mahaffey KW, Hedlin H, et al. Large-scale assessment of a smartwatch to identify atrial fibrillation. *N Engl J Med*. 2019;381:1909-1917.
104. Barbagelata A, Bethea CF, Severance HW, et al. Smartphone ECG for evaluation of ST-segment elevation myocardial infarction (STEMI): design of the ST LEUIS International Multicenter Study. *J Electrocardiol*. 2018;51:260-264.
105. Lim YW, Levis J, Pursnani SK. Detection of ST-elevation myocardial infarction via at-home ECG monitoring device. *Am J Med*. 2021;134:1242-1243.
106. Spaccarotella CAM, Polimeni A, Migliarino S, et al. Multichannel electrocardiograms obtained by a smartwatch for the diagnosis of ST-segment changes. *JAMA Cardiol*. 2020;5:1176-1180.
107. Drexler M, Elsner C, Gabelmann V, Gori T, Münzel T. Apple Watch detecting coronary ischaemia during chest pain episodes or an apple a day may keep myocardial infarction away. *Eur Heart J*. 2020;41:2224.
108. Pipek LZ, Nascimento RFV, Acencio MMP, Teixeira LR. Comparison of SpO<sub>2</sub> and heart rate values on Apple Watch and conventional commercial oximeters devices in patients with lung disease. *Sci Rep*. 2021;11:18901.
109. Bent B, Goldstein BA, Kibbe WA, Dunn JP. Investigating sources of inaccuracy in wearable optical heart rate sensors. *NPJ Digit Med*. 2020;3:18.
110. Stopyra JP, Snively AC, Ashburn NP, et al. EMS blood collection from patients with acute chest pain reduces emergency department length of stay. *Am J Emerg Med*. 2021;47:248-252.
111. DuCharme B, Macchi Bires A, Montanye E, et al. Effect of prehospital blood draws on length of stay for chest pain patients in the emergency department: a pilot study. *Crit Care Nurs Q*. 2019;42:208-214.
112. Stopyra JP, Snively AC, Scheidler JF, et al. Point-of-care troponin testing during ambulance transport to detect acute myocardial infarction. *Prehosp Emerg Care*. 2020;24:751-759.
113. Martín-Rodríguez F, Sanz-García A, Castro-Portillo E, et al. Prehospital troponin as a predictor of early clinical deterioration. *Eur J Clin Invest*. 2021;51:e13591.
114. Sagel D, Vlaar PJ, van Roosmalen R, et al. Prehospital risk stratification in patients with chest pain. *Emerg Med J*. 2021;38:814-819.
115. Cooper JG, Ferguson J, Donaldson LA, et al. The Ambulance Cardiac Chest Pain Evaluation in Scotland Study (ACCESS): a prospective cohort study. *Ann Emerg Med*. 2021;77:575-588.
116. Ishak M, Ali D, Fokkert MJ, et al. Fast assessment and management of chest pain patients without ST-elevation in the pre-hospital gateway (Famous Triage): ruling out a myocardial infarction at home with the modified HEART score. *Eur Heart J Acute Cardiovasc Care*. 2018;7:102-110.
117. Aarts GWA, Camaro C, van Geuns RJ, et al. Acute rule-out of non-ST-segment elevation acute coronary syndrome in the (pre)hospital setting by HEART score assessment and a single point-of-care troponin: rationale and design of the ARTICA randomised trial. *BMJ Open*. 2020;10:e034403.
118. Ten Berg JM. Pre-hospital Evaluation of Chest Pain Patients With sUSpected Non ST-segment eLevation myocARdial Infarction Using the HEART-score With a Troponin Point-of-care Test (POPular HEART). *ClinicalTrials.gov*. Accessed January 15, 2022. <https://www.clinicaltrials.gov/ct2/show/NCT04851418>
119. Hippisley-Cox J, Coupland C, Brindle P. Derivation and validation of QStroke score for predicting risk of ischaemic stroke in primary care and comparison with other risk scores: a prospective open cohort study. *BMJ*. 2013;346:f2573.
120. McNamara RL, Kennedy KF, Cohen DJ, et al. Predicting in-hospital mortality in patients with acute myocardial infarction. *J Am Coll Cardiol*. 2016;68:626-635.
121. Goldstein BA, Navar AM, Pencina MJ. Risk prediction with electronic health records: the importance of model validation and clinical context. *JAMA Cardiol*. 2016;1:976-977.
122. Goldstein BA, Navar AM, Pencina MJ, Ioannidis JP. Opportunities and challenges in developing risk prediction models with electronic health records data: a systematic review. *J Am Med Inform Assoc*. 2017;24:198-208.
123. Corral-Acero J, Margara F, Marciniak M, et al. The 'Digital Twin' to enable the vision of precision cardiology. *Eur Heart J*. 2020;41:4556-4564.
124. Oikonomou EK, Van Dijk D, Parise H, et al. A phenomapping-derived tool to personalize the selection of anatomical vs. functional testing in evaluating chest pain (ASSIST). *Eur Heart J*. 2021;42:2536-2548.
125. Cho Y, Kwon JM, Kim KH, et al. Artificial intelligence algorithm for detecting myocardial infarction using six-lead electrocardiography. *Sci Rep*. 2020;10:20495.
126. Al-Zaiti S, Besomi L, Bouzid Z, et al. Machine learning-based prediction of acute coronary syndrome using only the pre-hospital 12-lead electrocardiogram. *Nat Commun*. 2020;11:3966.
127. Bigler MR, Seiler C. Detection of myocardial ischemia by intracoronary ECG using convolutional neural networks. *PLoS One*. 2021;16:e0253200.
128. Stewart J, Lu J, Goudie A, et al. Applications of machine learning to undifferentiated chest pain in the emergency department: a systematic review. *PLoS One*. 2021;16:e0252612.
129. Than MP, Pickering JW, Sandoval Y, et al. Machine learning to predict the likelihood of acute myocardial infarction. *Circulation*. 2019;140:899-909.
130. Lynam AL, Dennis JM, Owen KR, et al. Logistic regression has similar performance to optimised machine learning algorithms in a clinical setting: application to the discrimination between type 1 and type 2 diabetes in young adults. *Diagn Progn Res*. 2020;4:6.
131. Liu N, Chee ML, Koh ZX, et al. Utilizing machine learning dimensionality reduction for risk stratification of chest pain patients in the emergency department. *BMC Med Res Methodol*. 2021;21:74.
132. Marber MS, Mills NL, Morrow DA, Mueller C. Cardiac myosin-binding protein C as a biomarker of acute myocardial infarction. *Eur Heart J Acute Cardiovasc Care*. 2021;10:963-965.
133. Kaier TE, Twerenbold R, Puelacher C, et al. Direct comparison of cardiac myosin-binding protein C with cardiac troponins for the early diagnosis of acute myocardial infarction. *Circulation*. 2017;136:1495-1508.
134. Schulte C, Barwari T, Joshi A, et al. Comparative analysis of circulating noncoding RNAs versus protein biomarkers in the detection of myocardial injury. *Circ Res*. 2019;125:328-340.
135. Agwa SHA, Elzahwy SS, El Meteini MS, et al. ABHD4-regulating RNA panel: novel biomarkers in acute coronary syndrome diagnosis. *Cells*. 2021;10(6):1512. <https://doi.org/10.3390/cells10061512>
136. Horiuchi Y, Wettersten N, Patel MP, et al. Biomarkers enhance discrimination and prognosis of type 2 myocardial infarction. *Circulation*. 2020;142:1532-1544.
137. Raskovalova T, Twerenbold R, Collinson PO, et al. Diagnostic accuracy of combined cardiac troponin and copeptin assessment for early rule-out of myocardial infarction: a systematic review and meta-analysis. *Eur Heart J Acute Cardiovasc Care*. 2014;3:18-27.
138. Pickering JW, Young JM, George PM, et al. Validity of a novel point-of-care troponin assay for single-test rule-out of acute myocardial infarction. *JAMA Cardiol*. 2018;3:1108-1112.

139. Nieman K, Hoffmann U. Cardiac computed tomography in patients with acute chest pain. *Eur Heart J*. 2015;36:906–914.
140. Litt HI, Gatsonis C, Snyder B, et al. CT angiography for safe discharge of patients with possible acute coronary syndromes. *N Engl J Med*. 2012;366:1393–1403.
141. Hoffmann U, Truong QA, Schoenfeld DA, et al. Coronary CT angiography versus standard evaluation in acute chest pain. *N Engl J Med*. 2012;367:299–308.
142. Reinhardt SW, Lin CJ, Novak E, Brown DL. Noninvasive cardiac testing vs clinical evaluation alone in acute chest pain: a secondary analysis of the ROMICAT-II randomized clinical trial. *JAMA Intern Med*. 2018;178:212–219.
143. Dedic A, Lubbers MM, Schaap J, et al. Coronary CT angiography for suspected ACS in the era of high-sensitivity troponins: randomized multicenter study. *J Am Coll Cardiol*. 2016;67:16–26.
144. Case BC, Yerasi C, Wang Y, et al. Admissions rate and timing of revascularization in the United States in patients with non-ST-elevation myocardial infarction. *Am J Cardiol*. 2020;134:24–31. <https://doi.org/10.1016/j.amjcard.2020.08.010>
145. Smulders MW, Kietselaer B, Wildberger JE, et al. Initial imaging-guided strategy versus routine care in patients with Non-ST-segment elevation myocardial infarction. *J Am Coll Cardiol*. 2019;74:2466–2477.
146. Linde JJ, Kelbæk H, Hansen TF, et al. Coronary CT angiography in patients with non-ST-segment elevation acute coronary syndrome. *J Am Coll Cardiol*. 2020;75:453–463.
147. Hess EP, Knoedler MA, Shah ND, et al. The chest pain choice decision aid: a randomized trial. *Circ Cardiovasc Qual Outcomes*. 2012;5:251–259.
148. Hess EP, Hollander JE, Schaffer JT, et al. Shared decision making in patients with low risk chest pain: prospective randomized pragmatic trial. *BMJ*. 2016;355:i6165.
149. Stacey D, Légaré F, Lewis K, et al. Decision aids for people facing health treatment or screening decisions. *Cochrane Database Syst Rev*. 2017;4:Cd001431.
150. Dawson LP, Biswas S, Lefkovits J, et al. Characteristics and quality of national cardiac registries: a systematic review. *Circ Cardiovasc Qual Outcomes*. 2021;14:e007963.
151. Nathan AS, Xiang Q, Wojdyla D, et al. Performance of hospitals when assessing disease-based mortality compared with procedural mortality for patients with acute myocardial infarction. *JAMA Cardiol*. 2020;5:765–772.
152. Winchester DE, Osborne A, Peacock WF, et al. Closing gaps in essential chest pain care through accreditation. *J Am Coll Cardiol*. 2020;75:2478–2482.
153. El-Sayed Ahmed MM, Thomas M, Jacob S, et al. Triple bridge of mechanical circulatory support to heart transplantation listing: a case report. *SAGE Open Med Case Rep*. 2019;7:2050313x19834816.

---

**KEY WORDS** acute chest pain, clinical outcomes, health systems, models of care

---

**APPENDIX** For supplemental tables, please see the online version of this paper.