

Instructions for ordering the Buprenorphine (SUBUTEX) Starter Pack in the MSH ED

Key points:

1. Go to Order Set (not discharge prescription)
2. Click box for Starter Pack
3. Call Social Work to pick up the prescription

Instructions:

1. Go to orders and search “buprenorphine”

The screenshot displays the MSH ED Orders interface. The top navigation bar includes tabs for Chart Review, Triage, Orders, ED Notes, DC/Transfer, and Sidebar Sum... The Orders tab is active, showing a search for "buprenorphine" in the Manage Orders section. The search results are displayed in a table with columns for Name, User Version Name, Type, and Lab(s). The results show "Buprenorphine" as an Order Set and "Buprenorphine Conf U" as a Lab test.

Orders Interface:

- Top navigation: Chart Review, Triage, Orders, ED Notes, DC/Transfer, Sidebar Sum...
- Order Sets: Suggested (3), Critical Care Orders, ED COVID Admissions, Intubation
- Search: buprenorphine
- Order Sets List: Buprenorphine (Order Set)
- Order Sets Details: Buprenorphine (Order Set)

Order and Order Set Search:

Search: BUPRENORPHINE

Order Sets & Panels:

Name	User Version Name	Type
Buprenorphine		Order Set

Medications: (No results found)

Procedures:

Name	Type	Pref List	Px Code	Lab(s)
Buprenorphine Conf U	Lab	MSH ED LA...	105020	LABCORP

Buttons: Select And Stay, Accept, Cancel

2. To order the starter pack, click “opioid use disorder DISCHARGE PRESCRIPTION Starter Pack – Buprenorphine HCVL (SUBUTEX) SL tablet”

Order Sets

- Provide a bridge prescription to last until the patient's scheduled appointment with their prescriber. Effective December 29, 2022, the DEA no longer requires an X-waiver to prescribe buprenorphine for opioid use disorder treatment.
- Inpatient Buprenorphine Treatment Protocol for Hospitalized Patients

▼ **Medications**

▼ **Buprenorphine HCL (SUBUTEX) SL tablet**

- ☒ Buprenorphine HCL (SUBUTEX) SL tablet
 - ☒ buprenorphine HCL (SUBUTEX) SL tablet
 - 2 mg (0.0315 mg/kg), sublingual, DAILY, Include Now
 - Do NOT initiate buprenorphine for the first time in someone who recently took other opioids and is not currently in withdrawal. This may cause precipitated withdrawal. For patients who wish to initiate buprenorphine treatment and are currently receiving full agonist opioids, please consult the REACH-IN team at inpatientbupe@mountsinai.org or Epic chat Drs. Michael Herscher and Linda Wang.
 - ☒ Opioid Use Disorder DISCHARGE PRESCRIPTION Starter Pack - Buprenorphine HCL (SUBUTEX) SL tablet
 - The 7-day discharge prescription starter pack is for patients who will have difficulty filling prescription post-discharge to ensure continuity of treatment.
 - It will be given to the patient at bedside before discharge and will NOT be sent to a community pharmacy.
 - To discharge a patient with a 7-day starter pack, contact SW covering your area to process request through the patient prescription drug program.
 - ☒ buprenorphine HCL (SUBUTEX) 8 mg tablet, sublingual
 - Take 1 to 3 tablets daily, as directed by provider. Max 3 tablets per day, Disp-21 tablet, R-0, E-Prescribing, Maximum MME/Day: Unknown for this order

⚠ Maximum MME cannot be calculated for this prescription. Enter discrete sig details to calculate maximum MME.

3. Call Social Work to pick up the prescription and deliver it to the patient.
4. If you do not click off on “opioid use disorder DISCHARGE PRESCRIPTION Starter Pack – Buprenorphine HCVL (SUBUTEX) SL tablet” you can order a regular Subutex prescription.

Order Sets

Orders

Clear All Orders

Buprenorphine

Remove Order Sets

Do NOT initiate buprenorphine for the first time in someone who recently took other opioids (e.g. <6 hours for short-acting opioids) and is not currently in withdrawal. This may cause precipitated withdrawal. Exception to this rule is starting buprenorphine via low dose initiation or microinduction.

The ordering team must ensure the following in advance of discharge:

- if the patient is newly started on buprenorphine, ensure they have a scheduled appointment within 1-2 weeks post-discharge with a provider who can continue their buprenorphine.
- Provide a bridge prescription to last until the patient's scheduled appointment with their prescriber. Effective December 29, 2022, the DEA no longer requires an X-waiver to prescribe buprenorphine for opioid use disorder treatment.

- Inpatient Buprenorphine Treatment Protocol for Hospitalized Patients

Medications

Buprenorphine HCL (SUBUTEX) SL tablet

☒ Buprenorphine HCL (SUBUTEX) SL tablet

☒ buprenorphine HCL (SUBUTEX) SL tablet
 2 mg (0.0315 mg/kg), sublingual, DAILY, Include Now
 Do NOT initiate buprenorphine for the first time in someone who recently took other opioids and is not currently in withdrawal. This may cause precipitated withdrawal. For patients who wish to initiate buprenorphine treatment and are currently receiving full agonist opioids, please consult the REACH-IN team at inpatientbupe @mountsinai.org or Epic chat Drs. Michael Herscher and Linda Wang.

☐ Opioid Use Disorder DISCHARGE PRESCRIPTION Starter Pack - Buprenorphine HCL (SUBUTEX) SL tablet

Manage Orders

Order Sets

Options

Place orders or order sets

New

Standard

Next

Orders from Order Sets

Buprenorphine

☒ buprenorphine HCL (SUBUTEX) SL tablet
 2 mg (0.0315 mg/kg), sublingual, DAILY, Include Now
 Do NOT initiate buprenorphine for the first time in someone who recently took other opioids and is not currently in withdrawal. This may cause precipitated withdrawal. For patients who wish to initiate buprenorphine treatment and are currently receiving full agonist opioids, please consult the REACH-IN team at inpatientbupe @mountsinai.org or Epic chat Drs. Michael Herscher and Linda Wang.

Remove All

Save Work

Sign

buprenorphine HCL (SUBUTEX) SL tablet

Accept Cancel

Order Instructions:

Do NOT initiate buprenorphine for the first time in someone who recently took other opioids and is not currently in withdrawal...

Report:

Lab Test Results

Component	Time Elapsed	Value	Range	Status
QTc	1050 days (01/03/21 1939)	434	ms	Final result

Reference Links:

Buprenorphine Treatment Protocol

Lexi-Comp

Lexi-Comp Peds

Protocol Document

http://mshsintranet.mountsinai.org/uploadedFiles/Protocols_Reference/Inpatient%20Buprenorphine%20Treatment%20Protocol...

For patients being continued on outpatient buprenorphine treatment, is the patient currently receiving or needs to receive opioid agonists in the hospital?

Yes

No

Dose:

4

mg

2 mg

4 mg

6 mg

8 mg

12 mg

16 mg

Calculated dose:

4 mg (0.063 mg/kg) = 2 tablet

Route:

sublingual

Frequency:

DAILY

DAILY

Starting

11/20/2023

Today

Tomorrow

For

Doses

Hours

Days

First Dose

Include Now

As Scheduled

First Dose: Today 1730

Final Dose: Until Discontinued

11/20	11/21	11/22	11/23	11/24	...
1730	1000	1000	1000	1000	

Admin Instructions:

Do NOT crush or chew.

Priority:

STAT

Routine

STAT

Class:

E-Prescribing

Fax

Historical Med

Inpatient Discharge To Outside Facility

No Print

Normal

Office Procedures/Supplies

Next Required

Accept Cancel