

# Algebraic Data Types for Hazel

Eric Griffis  
egriffis@umich.edu

## 1 Syntax

$\text{HTyp} \quad \tau ::= \emptyset \mid \tau \rightarrow \tau \mid \alpha \mid \mu\pi.\tau \mid +\{C_i(\tau_i)\}_{C_i \in \mathcal{C}} \mid +\{C(\tau); \dots\} \mid \langle \rangle \mid \langle \alpha \rangle$   
 $\text{HTypPat} \quad \pi ::= \alpha \mid \langle \rangle$   
 $\text{HExp} \quad e ::= \emptyset \mid x \mid \lambda x:\tau.e \mid e(e) \mid e:\tau \mid \text{inj}_C(e) \mid \text{roll}(e) \mid \text{unroll}(e)$   
 $\quad \mid \langle \rangle^u \mid \langle e \rangle^u \mid \langle e \rangle^{u\blacktriangleright}$   
 $\text{IHExp} \quad d ::= \emptyset \mid x \mid \lambda x:\tau.d \mid d(d) \mid \text{inj}_C^\tau(d) \mid \text{roll}^{\mu\alpha.\tau}(d) \mid \text{unroll}(d)$   
 $\quad \mid d\langle \tau \Rightarrow \tau \rangle \mid d\langle \tau \Rightarrow \langle \rangle \nRightarrow \tau \rangle \mid \langle \rangle_\sigma^u \mid \langle d \rangle_\sigma^u \mid \langle d \rangle_\sigma^{u\blacktriangleright}$   
 $\text{HTag} \quad C ::= \mathbf{C} \mid \langle \rangle^u \mid \langle \mathbf{C} \rangle^u$

### 1.1 Context Extension

We write  $\Theta, \pi$  to denote the extension of type variable context  $\Theta$  with optional type variable name  $\pi$ .

$$\Theta, \pi = \begin{cases} \Theta, \alpha & \pi = \alpha \\ \Theta & \pi = \langle \rangle \end{cases}$$

## 2 Static Semantics

$\boxed{[\tau/\pi]\tau' = \tau''} \quad \tau'' \text{ is obtained by substituting } \tau \text{ for } \pi \text{ in } \tau'$

$$\begin{array}{lll}
[\tau/\langle \rangle]\tau' & = & \tau' \\
[\tau/\alpha]\emptyset & = & \emptyset \\
[\tau/\alpha](\tau_1 \rightarrow \tau_2) & = & [\tau/\alpha]\tau_1 \rightarrow [\tau/\alpha]\tau_2 \\
[\tau/\alpha]\alpha & = & \tau \\
[\tau/\alpha]\alpha_1 & = & \tau' & \text{when } \alpha \neq \alpha_1 \\
[\tau/\alpha]\mu\alpha_1.\tau_2 & = & \mu\alpha_1.[\tau/\alpha]\tau_2 & \text{when } \alpha \neq \alpha_1 \text{ and } \alpha_1 \notin \text{FV}(\tau) \\
[\tau/\alpha]\mu\langle \rangle.\tau_2 & = & \mu\langle \rangle.[\tau/\alpha]\tau_2 \\
[\tau/\alpha]+\{C_i(\tau_i)\}_{C_i \in \mathcal{C}} & = & +\{C_i([\tau/\alpha]\tau_i)\}_{C_i \in \mathcal{C}} \\
[\tau/\alpha]+\{C(\tau'); \dots\} & = & +\{C([\tau/\alpha]\tau'); \dots\} \\
[\tau/\alpha]\langle \rangle & = & \langle \rangle \\
[\alpha'/\alpha]\langle \alpha \rangle & = & \langle \alpha' \rangle \\
[\alpha'/\alpha]\langle \alpha' \rangle & = & \langle \alpha' \rangle & \text{when } \alpha \neq \alpha'
\end{array}$$

$\boxed{\Theta \vdash \tau \text{ valid}}$   $\tau$  is a valid type

$$\begin{array}{c}
\text{TVUNIT} \\
\hline
\Theta \vdash \emptyset \text{ valid}
\end{array}
\quad
\begin{array}{c}
\text{TVARR} \\
\hline
\frac{\Theta \vdash \tau_1 \text{ valid} \quad \Theta \vdash \tau_2 \text{ valid}}{\Theta \vdash \tau_1 \rightarrow \tau_2 \text{ valid}}
\end{array}
\quad
\begin{array}{c}
\text{TVVAR} \\
\hline
\frac{\alpha \in \Theta}{\Theta \vdash \alpha \text{ valid}}
\end{array}
\quad
\begin{array}{c}
\text{TVREC} \\
\hline
\frac{\Theta, \pi \vdash \tau \text{ valid}}{\Theta \vdash \mu\pi.\tau \text{ valid}}
\end{array}
\quad
\begin{array}{c}
\text{TVSUM1} \\
\hline
\frac{\{\Theta \vdash \tau_i \text{ valid}\}_{C_i \in \mathcal{C}}}{\Theta \vdash +\{C_i(\tau_i)\}_{C_i \in \mathcal{C}} \text{ valid}}
\end{array}$$

$$\begin{array}{c}
\text{TVSUM2} \\
\hline
\frac{\Theta \vdash \tau \text{ valid}}{\Theta \vdash +\{C(\tau); \dots\} \text{ valid}}
\end{array}
\quad
\begin{array}{c}
\text{TVEHOLE} \\
\hline
\frac{}{\Theta \vdash \langle \rangle \text{ valid}}
\end{array}
\quad
\begin{array}{c}
\text{TVNEHOLE} \\
\hline
\frac{\alpha \notin \Theta}{\Theta \vdash \langle \alpha \rangle \text{ valid}}
\end{array}$$

$\boxed{\tau \sim \tau'}$   $\tau$  and  $\tau'$  are consistent

$$\begin{array}{c}
\text{TCREFL} \\
\hline
\tau \sim \tau
\end{array}
\quad
\begin{array}{c}
\text{TCEHOLE1} \\
\hline
\langle \rangle \sim \tau
\end{array}
\quad
\begin{array}{c}
\text{TCEHOLE2} \\
\hline
\tau \sim \langle \rangle
\end{array}
\quad
\begin{array}{c}
\text{TCNEHOLE1} \\
\hline
\langle \alpha \rangle \sim \tau
\end{array}
\quad
\begin{array}{c}
\text{TCNEHOLE2} \\
\hline
\tau \sim \langle \alpha \rangle
\end{array}
\quad
\begin{array}{c}
\text{TCARR} \\
\hline
\frac{\tau_1 \sim \tau'_1 \quad \tau_2 \sim \tau'_2}{\tau_1 \rightarrow \tau_2 \sim \tau'_1 \rightarrow \tau'_2}
\end{array}$$

$$\begin{array}{c}
\text{TCREC} \\
\hline
\frac{\tau \sim \tau'}{\mu\pi.\tau \sim \mu\pi.\tau'}
\end{array}
\quad
\begin{array}{c}
\text{TCRECHOLE1} \\
\hline
\frac{\alpha \notin \text{FV}(\tau) \quad \tau \sim \tau'}{\mu\langle \rangle.\tau \sim \mu\alpha.\tau'}
\end{array}
\quad
\begin{array}{c}
\text{TCRECHOLE2} \\
\hline
\frac{\alpha \notin \text{FV}(\tau) \quad \tau \sim \tau'}{\mu\alpha.\tau \sim \mu\langle \rangle.\tau'}
\end{array}$$

$$\begin{array}{c}
\text{TCSUM1} \\
\hline
\frac{|\mathcal{C}| = |\mathcal{C}'| \quad \{C_i \sim C'_i\}_{i < |\mathcal{C}|} \quad \{\tau_i \sim \tau'_i\}_{i < |\mathcal{C}|}}{+\{C_i(\tau_i)\}_{C_i \in \mathcal{C}} \sim +\{C'_i(\tau'_i)\}_{C_i \in \mathcal{C}'}}
\end{array}
\quad
\begin{array}{c}
\text{TCSUM2} \\
\hline
\frac{\tau \sim \tau'}{+\{C(\tau); \dots\} \sim +\{C(\tau'); \dots\}}
\end{array}$$

$$\begin{array}{c}
\text{TCSUM12} \\
\hline
\frac{C_j \in \mathcal{C} \quad \tau \sim \tau_j}{+\{C_i(\tau_i)\}_{C_i \in \mathcal{C}} \sim +\{C_j(\tau); \dots\}}
\end{array}
\quad
\begin{array}{c}
\text{TCSUM21} \\
\hline
\frac{C_j \in \mathcal{C} \quad \tau \sim \tau_j}{+\{C_j(\tau); \dots\} \sim +\{C_i(\tau_i)\}_{C_i \in \mathcal{C}}}
\end{array}$$

$\boxed{C \sim C'}$   $C$  and  $C'$  are consistent

$$\begin{array}{c}
\text{TAGCREFL} \\
\hline
\overline{C \sim C}
\end{array}
\quad
\begin{array}{c}
\text{TAGCEHOLE1} \\
\hline
\overline{\langle \rangle^u \sim C}
\end{array}
\quad
\begin{array}{c}
\text{TAGCEHOLE2} \\
\hline
\overline{C \sim \langle \rangle^u}
\end{array}
\quad
\begin{array}{c}
\text{TAGCNEHOLE1} \\
\hline
\overline{\langle C \rangle^u \sim C'}
\end{array}
\quad
\begin{array}{c}
\text{TAGCNEHOLE2} \\
\hline
\overline{C \sim \langle C' \rangle^u}
\end{array}$$

## 2.1 Bidirectional Typing

We call  $[\mu\pi.\tau/\pi]\tau$  the *unrolling* of recursive type  $\mu\pi.\tau$ .

**Theorem 1** (Synthetic Type Validity). *If  $\Gamma \vdash e \Rightarrow \tau$  then  $\emptyset \vdash \tau \text{ valid}$ .*

**Theorem 2** (Consistency Preserves Validity). *If  $\Theta \vdash \tau \text{ valid}$  and  $\tau \sim \tau'$  then  $\Theta \vdash \tau' \text{ valid}$ .*

$\boxed{\tau \blacktriangleright_{\rightarrow} \tau_1 \rightarrow \tau_2}$   $\tau$  has matched arrow type  $\tau_1 \rightarrow \tau_2$

$$\begin{array}{c}
\text{MAHOLE} \\
\hline
\langle \rangle \blacktriangleright_{\rightarrow} \langle \rangle \rightarrow \langle \rangle
\end{array}
\quad
\begin{array}{c}
\text{MAARR} \\
\hline
\tau_1 \rightarrow \tau_2 \blacktriangleright_{\rightarrow} \tau_1 \rightarrow \tau_2
\end{array}$$

$\boxed{\tau \blacktriangleright_{\mu} \mu\pi.\tau'}$   $\tau$  has matched recursive type  $\mu\pi.\tau'$

$$\begin{array}{c}
\text{MRREC} \\
\hline
\mu\pi.\tau \blacktriangleright_{\mu} \mu\pi.\tau
\end{array}
\quad
\begin{array}{c}
\text{MRHOLE} \\
\hline
\langle \rangle \blacktriangleright_{\mu} \mu\langle \rangle.\langle \rangle
\end{array}$$

$\boxed{\tau \blacktriangleright_+ + \{C_i(\tau_i)\}_{C_i \in \mathcal{C}}}$   $\tau$  has matched finite sum type  $+ \{C_i(\tau_i)\}_{C_i \in \mathcal{C}}$

$$\begin{array}{c}
\text{MSFINITE} \quad \frac{}{+ \{C_i(\tau_i)\}_{C_i \in \mathcal{C}} \blacktriangleright_+ + \{C_i(\tau_i)\}_{C_i \in \mathcal{C}}} \quad \text{MSELIDED} \quad \frac{}{+ \{C(\tau); \dots\} \blacktriangleright_+ + \{C(\tau)\}} \\
\\
\boxed{\Gamma \vdash e \Rightarrow \tau} \quad e \text{ synthesizes type } \tau \\
\\
\text{SUNIT} \quad \frac{}{\Gamma \vdash \emptyset \Rightarrow \emptyset} \quad \text{SVAR} \quad \frac{x : \tau \in \Gamma}{\Gamma \vdash x \Rightarrow \tau} \quad \text{SVARFREE} \quad \frac{x \notin \text{dom}(\Gamma)}{\Gamma \vdash \langle x \rangle^u \Rightarrow \langle \rangle} \quad \text{SLAM} \quad \frac{\emptyset \vdash \tau \text{ valid} \quad \Gamma, x : \tau \vdash e \Rightarrow \tau'}{\Gamma \vdash \lambda x : \tau. e \Rightarrow \tau \rightarrow \tau'} \\
\\
\text{SAPP} \quad \frac{\Gamma \vdash e_1 \Rightarrow \tau_1 \quad \tau_1 \blacktriangleright \rightarrow \tau_2 \rightarrow \tau \quad \Gamma \vdash e_2 \Leftarrow \tau_2}{\Gamma \vdash e_1(e_2) \Rightarrow \tau} \quad \text{SAPPNOTARR} \quad \frac{\Gamma \vdash e_1 \Rightarrow \tau_1 \quad \tau_1 \approx \langle \rangle \rightarrow \langle \rangle \quad \Gamma \vdash e_2 \Leftarrow \langle \rangle}{\Gamma \vdash \langle e_1 \rangle^u \blacktriangleright (e_2) \Rightarrow \langle \rangle} \\
\\
\text{SASC} \quad \frac{\emptyset \vdash \tau \text{ valid} \quad \Gamma \vdash e \Leftarrow \tau}{\Gamma \vdash e : \tau \Rightarrow \tau} \quad \text{SROLLERR} \quad \frac{\Gamma \vdash e \Leftarrow \langle \rangle}{\Gamma \vdash \langle \text{roll}(e) \rangle^u \Rightarrow \mu \langle \rangle. \langle \rangle} \quad \text{SUNROLL} \quad \frac{\Gamma \vdash e \Rightarrow \tau \quad \tau \blacktriangleright_\mu \mu \pi. \tau'}{\Gamma \vdash \text{unroll}(e) \Rightarrow [\mu \pi. \tau' / \pi] \tau'} \\
\\
\text{SUNROLLNOTREC} \quad \frac{\Gamma \vdash e \Rightarrow \tau \quad \tau \approx \mu \langle \rangle. \langle \rangle}{\Gamma \vdash \text{unroll}(\langle e \rangle^u \blacktriangleright) \Rightarrow \langle \rangle} \quad \text{SINJ} \quad \frac{\Gamma \vdash e \Rightarrow \tau}{\Gamma \vdash \text{inj}_C(e) \Rightarrow + \{C(\tau); \dots\}} \quad \text{SEHOLE} \quad \frac{}{\Gamma \vdash \langle \rangle^u \Rightarrow \langle \rangle} \quad \text{SNEHOLE} \quad \frac{}{\Gamma \vdash \langle e \rangle^u \Rightarrow \langle \rangle} \\
\\
\boxed{\Gamma \vdash e \Leftarrow \tau} \quad e \text{ analyzes against type } \tau \\
\\
\text{AROLL} \quad \frac{\tau \blacktriangleright_\mu \mu \pi. \tau' \quad \Gamma \vdash e \Leftarrow [\mu \pi. \tau' / \pi] \tau'}{\Gamma \vdash \text{roll}(e) \Leftarrow \tau} \quad \text{AROLLNOTREC} \quad \frac{\tau \approx \mu \langle \rangle. \langle \rangle \quad \Gamma \vdash e \Leftarrow \langle \rangle}{\Gamma \vdash \langle \text{roll}(e) \rangle^u \Leftarrow \tau} \quad \text{AINJHOLE} \quad \frac{}{\Gamma \vdash \text{inj}_C(e) \Leftarrow \langle \rangle} \\
\\
\text{AINJ} \quad \frac{\tau \blacktriangleright_+ + \{C_i(\tau_i)\}_{C_i \in \mathcal{C}} \quad C_j \in \mathcal{C} \quad \Gamma \vdash e \Leftarrow \tau_j}{\Gamma \vdash \text{inj}_{C_j}(e) \Leftarrow \tau} \quad \text{AINJTAGER} \quad \frac{\tau \blacktriangleright_+ + \{C_i(\tau_i)\}_{C_i \in \mathcal{C}} \quad C \notin \mathcal{C} \quad \Gamma \vdash e \Leftarrow \langle \rangle}{\Gamma \vdash \langle \text{inj}_C(e) \rangle^u \Leftarrow \tau} \\
\\
\text{AINJUNEXPECTEDARG} \quad \frac{\tau \blacktriangleright_+ + \{C_i(\tau_i)\}_{C_i \in \mathcal{C}} \quad C_j \in \mathcal{C} \quad \tau_j = \emptyset \quad e \neq \emptyset \quad \Gamma \vdash e \Leftarrow \langle \rangle}{\Gamma \vdash \langle \text{inj}_{C_j}(e) \rangle^u \Leftarrow \tau} \\
\\
\text{AINJEXPECTEDARG} \quad \frac{\tau \blacktriangleright_+ + \{C_i(\tau_i)\}_{C_i \in \mathcal{C}} \quad C_j \in \mathcal{C} \quad \tau_j \neq \emptyset}{\Gamma \vdash \langle \text{inj}_{C_j}(\emptyset) \rangle^u \Leftarrow \tau} \quad \text{ASUBSUME} \quad \frac{\Gamma \vdash e \Rightarrow \tau' \quad \tau' \sim \tau}{\Gamma \vdash e \Leftarrow \tau}
\end{array}$$

## 2.2 Typed Elaboration

**Theorem 3** (Synthetic Typed Elaboration Validity). *If  $\Gamma \vdash e \Rightarrow \tau \rightsquigarrow d \dashv \Delta$  then  $\emptyset \vdash \tau \text{ valid}$ .*

$\boxed{\Gamma \vdash e \Rightarrow \tau \rightsquigarrow d \dashv \Delta}$   $e$  synthesizes type  $\tau$  and elaborates to  $d$

$$\begin{array}{c}
\text{ESUNIT} \quad \frac{}{\Gamma \vdash \emptyset \Rightarrow \emptyset \rightsquigarrow \emptyset \dashv \emptyset} \quad \text{ESVAR} \quad \frac{x : \tau \in \Gamma}{\Gamma \vdash x \Rightarrow \tau \rightsquigarrow x \dashv \emptyset} \quad \text{ESVARFREE} \quad \frac{x \notin \text{dom}(\Gamma)}{\Gamma \vdash \langle x \rangle^u \Rightarrow \langle \rangle \rightsquigarrow \langle x \rangle_{\text{id}(\Gamma)}^u \dashv u :: \langle \rangle [\Gamma]} \\
\\
\text{ESLAM} \quad \frac{\emptyset \vdash \tau \text{ valid} \quad \Gamma, x : \tau \vdash e \Rightarrow \tau' \rightsquigarrow d \dashv \Delta}{\Gamma \vdash \lambda x : \tau. e \Rightarrow \tau \rightarrow \tau' \rightsquigarrow \lambda x : \tau. d \dashv \Delta}
\end{array}$$

$$\frac{\text{ESAPP} \quad \Gamma \vdash e_1 \Rightarrow \tau_1 \quad \tau_1 \blacktriangleright \rightarrow \tau_2 \rightarrow \tau \quad \Gamma \vdash e_1 \Leftarrow \tau_2 \rightarrow \tau \rightsquigarrow d_1 : \tau'_1 \dashv \Delta_1 \quad \Gamma \vdash e_2 \Leftarrow \tau_2 \rightsquigarrow d_2 : \tau'_2 \dashv \Delta_2}{\Gamma \vdash e_1(e_2) \Rightarrow \tau \rightsquigarrow (d_1 \langle \tau'_1 \Rightarrow \tau_1 \rangle) (d_2 \langle \tau'_2 \Rightarrow \tau_2 \rangle) \dashv \Delta_1 \cup \Delta_2}$$

$$\frac{\text{ESAPPNOTARR} \quad \Gamma \vdash e_1 \Rightarrow \tau_1 \rightsquigarrow d_1 \dashv \Delta_1 \quad \tau_1 \approx \langle \rangle \rightarrow \langle \rangle \quad \Gamma \vdash e_2 \Leftarrow \langle \rangle \rightsquigarrow d_2 : \tau'_2 \dashv \Delta_2}{\Gamma \vdash \langle e_1 \rangle^{u \blacktriangleright} (e_2) \Rightarrow \langle \rangle \rightsquigarrow \langle d_1 \rangle_{\text{id}(\Gamma)}^{u \blacktriangleright} (d_2 \langle \tau'_2 \Rightarrow \langle \rangle \rangle) \dashv \Delta_1 \cup \Delta_2, u :: \langle \rangle \rightarrow \langle \rangle [\Gamma]}$$

$$\frac{\text{ESAsc} \quad \emptyset \vdash \tau \text{ valid} \quad \Gamma \vdash e \Leftarrow \tau \rightsquigarrow d : \tau' \dashv \Delta}{\Gamma \vdash e : \tau \Rightarrow \tau \rightsquigarrow d \langle \tau' \Rightarrow \tau \rangle \dashv \Delta}$$

$$\frac{\text{ESROLLERR} \quad \Gamma \vdash e \Leftarrow \langle \rangle \rightsquigarrow d : \tau \dashv \Delta}{\Gamma \vdash \langle \text{roll}(e) \rangle^u \Rightarrow \mu \langle \rangle . \langle \rangle \rightsquigarrow \langle \text{roll}^{\mu \langle \rangle . \langle \rangle} (d \langle \tau \Rightarrow \langle \rangle \rangle) \rangle_{\text{id}(\Gamma)}^u \dashv \Delta, u :: \mu \langle \rangle . \langle \rangle [\Gamma]}$$

$$\frac{\text{ESUNROLL} \quad \Gamma \vdash e \Rightarrow \tau \rightsquigarrow d \dashv \Delta \quad \tau \blacktriangleright_{\mu} \mu \pi . \tau'}{\Gamma \vdash \text{unroll}(e) \Rightarrow [\mu \pi . \tau' / \pi] \tau' \rightsquigarrow \text{unroll}(d \langle \tau \Rightarrow \mu \pi . \tau' \rangle) \dashv \Delta}$$

$$\frac{\text{ESUNROLLNOTREC} \quad \Gamma \vdash e \Rightarrow \tau \rightsquigarrow d \dashv \Delta \quad \tau \approx \mu \langle \rangle . \langle \rangle}{\Gamma \vdash \text{unroll}(\langle e \rangle^{u \blacktriangleright}) \Rightarrow \langle \rangle \rightsquigarrow \text{unroll}(\langle d \rangle_{\text{id}(\Gamma)}^{u \blacktriangleright}) \dashv \Delta, u :: \mu \langle \rangle . \langle \rangle [\Gamma]}$$

$$\frac{\text{ESINJ} \quad \Gamma \vdash e \Rightarrow \tau \rightsquigarrow d \dashv \Delta \quad \tau' = +\{C(\tau); \dots\}}{\Gamma \vdash \text{inj}_C(e) \Rightarrow \tau' \rightsquigarrow \text{inj}_C^{\tau'}(d) \dashv \Delta} \quad \frac{\text{ESEHOLE}}{\Gamma \vdash \langle \rangle^u \Rightarrow \langle \rangle \rightsquigarrow \langle \rangle_{\text{id}(\Gamma)}^u \dashv u :: \langle \rangle [\Gamma]}$$

$$\frac{\text{ESNEHOLE} \quad \Gamma \vdash e \Rightarrow \tau \rightsquigarrow d \dashv \Delta}{\Gamma \vdash \langle e \rangle^u \Rightarrow \langle \rangle \rightsquigarrow \langle d \rangle_{\text{id}(\Gamma)}^u \dashv \Delta, u :: \langle \rangle [\Gamma]}$$

$$\boxed{\Gamma \vdash e \Leftarrow \tau_1 \rightsquigarrow d : \tau_2 \dashv \Delta} \quad e \text{ analyzes against type } \tau_1 \text{ and elaborates to } d \text{ of consistent type } \tau_2$$

$$\frac{\text{EAROLL} \quad \tau \blacktriangleright_{\mu} \mu \pi . \tau' \quad \Gamma \vdash e \Leftarrow [\mu \pi . \tau' / \pi] \tau' \rightsquigarrow d : \tau'' \dashv \Delta}{\Gamma \vdash \text{roll}(e) \Leftarrow \tau \rightsquigarrow \text{roll}^{\mu \pi . \tau'}(d \langle \tau'' \Rightarrow [\mu \pi . \tau' / \pi] \tau' \rangle) : \mu \pi . \tau' \dashv \Delta}$$

$$\frac{\text{EAROLLNOTREC} \quad \tau \approx \mu \langle \rangle . \langle \rangle \quad \Gamma \vdash e \Leftarrow \langle \rangle \rightsquigarrow d : \tau' \dashv \Delta}{\Gamma \vdash \langle \text{roll}(e) \rangle^u \Leftarrow \tau \rightsquigarrow \langle \text{roll}^{\mu \langle \rangle . \langle \rangle} (d) \rangle_{\text{id}(\Gamma)}^u : \mu \langle \rangle . \langle \rangle \dashv \Delta, u :: \mu \langle \rangle . \langle \rangle [\Gamma]}$$

$$\frac{\text{EAINJHOLE} \quad \Gamma \vdash e \Leftarrow \langle \rangle \rightsquigarrow d : \tau \dashv \Delta \quad \tau' = +\{C(\tau); \dots\}}{\Gamma \vdash \text{inj}_C(e) \Leftarrow \langle \rangle \rightsquigarrow \text{inj}_C^{\tau'}(d) : \tau' \dashv \Delta}$$

$$\frac{\text{EAINJ} \quad \tau \blacktriangleright_+ \tau' \quad \tau' = +\{C_i(\tau_i)\}_{C_i \in \mathcal{C}} \quad C_j \in \mathcal{C} \quad \Gamma \vdash e \Leftarrow \tau_j \rightsquigarrow d : \tau'_j \dashv \Delta}{\Gamma \vdash \text{inj}_{C_j}(e) \Leftarrow \tau \rightsquigarrow \text{inj}_{C_j}^{\tau'}(d \langle \tau'_j \Rightarrow \tau_j \rangle) : \tau \dashv \Delta}$$

$$\begin{array}{c}
\text{EAINJTAGERR} \\
\frac{\tau \blacktriangleright_+ + \{C_i(\tau_i)\}_{C_i \in \mathcal{C}} \quad C \notin \mathcal{C} \quad \Gamma \vdash e \Leftarrow \emptyset \rightsquigarrow d : \tau' \dashv \Delta \quad \tau'' = +\{C(\tau'); \dots\}}{\Gamma \vdash (\text{inj}_C(e))^u \Leftarrow \tau \rightsquigarrow (\text{inj}_C^{\tau''}(d))_{\text{id}(\Gamma)}^u : \tau \dashv \Delta, u :: \tau[\Gamma]}
\\[10pt]
\text{EAINJUNEXPECTEDARG} \\
\frac{\tau \blacktriangleright_+ + \{C_i(\tau_i)\}_{C_i \in \mathcal{C}} \quad C_j \in \mathcal{C} \quad \tau_j = \emptyset \quad e \neq \emptyset \quad \Gamma \vdash e \Leftarrow \emptyset \rightsquigarrow d : \tau'_j \dashv \Delta \quad \tau' = +\{C_j(\tau'_j); \dots\}}{\Gamma \vdash (\text{inj}_{C_j}(e))^u \Leftarrow \tau \rightsquigarrow (\text{inj}_{C_j}^{\tau'_j}(d))_{\text{id}(\Gamma)}^u : \tau \dashv \Delta, u :: \tau[\Gamma]}
\\[10pt]
\text{EAINJEXPECTEDARG} \\
\frac{\tau \blacktriangleright_+ + \{C_i(\tau_i)\}_{C_i \in \mathcal{C}} \quad C_j \in \mathcal{C} \quad \tau_j \neq \emptyset \quad \tau' = +\{C_j(\emptyset); \dots\}}{\Gamma \vdash (\text{inj}_{C_j}(\emptyset))^u \Leftarrow \tau \rightsquigarrow (\text{inj}_{C_j}^{\tau'}(\emptyset))_{\text{id}(\Gamma)}^u : \tau \dashv \Delta, u :: \tau[\Gamma]}
\\[10pt]
\text{EASUBSUME} \quad \text{EAEHOLE} \\
\frac{e \neq \emptyset^u \quad e \neq (e')^u \quad \Gamma \vdash e \Rightarrow \tau' \rightsquigarrow d \dashv \Delta \quad \tau \sim \tau'}{\Gamma \vdash e \Leftarrow \tau \rightsquigarrow d : \tau' \dashv \Delta} \quad \frac{}{\Gamma \vdash \emptyset^u \Leftarrow \tau \rightsquigarrow \emptyset_{\text{id}(\Gamma)}^u : \tau \dashv u :: \tau[\Gamma]}
\\[10pt]
\text{EANEHOLE} \\
\frac{\Gamma \vdash e \Rightarrow \tau' \rightsquigarrow d \dashv \Delta}{\Gamma \vdash (e)^u \Leftarrow \tau \rightsquigarrow (d)_{\text{id}(\Gamma)}^u : \tau \dashv u :: \tau[\Gamma]}
\end{array}$$

## 2.3 Type Assignment

$$\boxed{\Delta; \Gamma \vdash d : \tau} \quad d \text{ is assigned type } \tau$$

$$\begin{array}{c}
\text{TAVAR} \quad \text{TALAM} \quad \text{TAAAPP} \\
\frac{x : \tau \in \Gamma}{\Delta; \Gamma \vdash x : \tau} \quad \frac{\emptyset \vdash \tau \text{ valid} \quad \Delta; \Gamma, x : \tau \vdash d : \tau'}{\Delta; \Gamma \vdash \lambda x : \tau. d : \tau \rightarrow \tau'} \quad \frac{\Delta; \Gamma \vdash d_1 : \tau_2 \rightarrow \tau \quad \Delta; \Gamma \vdash d_2 : \tau_2}{\Delta; \Gamma \vdash d_1(d_2) : \tau}
\\[10pt]
\text{TAROLL} \quad \text{TAUNROLL} \\
\frac{\emptyset \vdash \mu\pi.\tau \text{ valid} \quad \Delta; \Gamma \vdash d : [\mu\pi.\tau/\pi]\tau}{\Delta; \Gamma \vdash \text{roll}^{\mu\pi.\tau}(d) : \mu\pi.\tau} \quad \frac{\Delta; \Gamma \vdash d : \mu\pi.\tau}{\Delta; \Gamma \vdash \text{unroll}(d) : [\mu\pi.\tau/\pi]\tau}
\\[10pt]
\text{TAINJ} \quad \text{TAEHOLE} \\
\frac{\tau \blacktriangleright_+ + \{C_i(\tau_i)\}_{C_i \in \mathcal{C}} \quad C_j \in \mathcal{C} \quad \Delta; \Gamma \vdash d : \tau_j}{\Delta; \Gamma \vdash \text{inj}_{C_j}^{\tau}(d) : \tau} \quad \frac{u :: \tau[\Gamma'] \in \Delta \quad \Delta; \Gamma \vdash \sigma : \Gamma'}{\Delta; \Gamma \vdash \emptyset_{\sigma}^u : \tau}
\\[10pt]
\text{TANEHOLE} \quad \text{TAMHOLE} \\
\frac{\Delta; \Gamma \vdash d : \tau' \quad u :: \tau[\Gamma'] \in \Delta \quad \Delta; \Gamma \vdash \sigma : \Gamma'}{\Delta; \Gamma \vdash (d)_{\sigma}^u : \tau} \quad \frac{\Delta; \Gamma \vdash d : \tau' \quad u :: \tau[\Gamma'] \in \Delta \quad \Delta; \Gamma \vdash \sigma : \Gamma'}{\Delta; \Gamma \vdash (d)_{\sigma}^{u\blacktriangleright} : \tau}
\\[10pt]
\text{TACAST} \quad \text{TAFAILEDCAST} \\
\frac{\Delta; \Gamma \vdash d : \tau_1 \quad \tau_1 \sim \tau_2}{\Delta; \Gamma \vdash d \langle \tau_1 \Rightarrow \tau_2 \rangle : \tau_2} \quad \frac{\Delta; \Gamma \vdash d : \tau_1 \quad \tau_1 \text{ ground} \quad \tau_2 \text{ ground} \quad \tau_1 \neq \tau_2}{\Delta; \Gamma \vdash d \langle \tau_1 \Rightarrow \emptyset \nRightarrow \tau_2 \rangle : \tau_2}
\end{array}$$

### 3 Dynamic Semantics

$\boxed{\tau \text{ ground}}$   $\tau$  is a ground type

$$\frac{\text{GARR}}{\emptyset \rightarrow \emptyset \text{ ground}}$$

$$\frac{\text{GREC}}{\mu \emptyset . \emptyset \text{ ground}}$$

$$\frac{\text{GSUM} \quad \tau \blacktriangleright_+ + \{C_i(\tau_i)\}_{C_i \in \mathcal{C}} \quad \{\tau_i = \emptyset \vee \tau_i = \emptyset\}_{C_i \in \mathcal{C}}}{\tau \text{ ground}}$$

$\boxed{\tau \blacktriangleright_{\text{ground}} \tau'}$   $\tau$  has matched ground type  $\tau'$

$$\frac{\text{MGARR} \quad \tau_1 \rightarrow \tau_2 \neq \emptyset \rightarrow \emptyset}{\tau_1 \rightarrow \tau_2 \blacktriangleright_{\text{ground}} \emptyset \rightarrow \emptyset}$$

$$\frac{\text{MGREC} \quad \tau \neq \emptyset}{\mu \pi . \tau \blacktriangleright_{\text{ground}} \mu \emptyset . \emptyset}$$

$$\frac{\text{MGSUM1} \quad \{(\tau_i = \emptyset \implies \tau'_i = \emptyset) \wedge (\tau_i \neq \emptyset \implies \tau'_i = \emptyset)\}_{C_i \in \mathcal{C}} \quad +\{C_i(\tau_i)\}_{C_i \in \mathcal{C}} \neq +\{C_i(\tau'_i)\}_{C_i \in \mathcal{C}}}{+\{C_i(\tau_i)\}_{C_i \in \mathcal{C}} \blacktriangleright_{\text{ground}} +\{C_i(\tau'_i)\}_{C_i \in \mathcal{C}}}$$

$$\frac{\text{MGSUM2} \quad \tau \neq \emptyset \quad \tau \neq \emptyset}{+\{C(\tau); \dots\} \blacktriangleright_{\text{ground}} +\{C(\emptyset); \dots\}}$$

$\boxed{d \text{ final}}$   $d$  is final

$$\frac{\text{FBOXEDVAL} \quad d \text{ boxedval}}{d \text{ final}}$$

$$\frac{\text{FINDET} \quad d \text{ indet}}{d \text{ final}}$$

$\boxed{d \text{ val}}$   $d$  is a value

$$\frac{\text{VUNIT}}{\emptyset \text{ val}}$$

$$\frac{\text{VLAM}}{\lambda x : \tau . d \text{ val}}$$

$$\frac{\text{VROLL} \quad d \text{ val}}{\text{roll}^{\mu \pi . \tau}(d) \text{ val}}$$

$$\frac{\text{VINJ} \quad d \text{ val}}{\text{inj}_C^\tau(d) \text{ val}}$$

$\boxed{d \text{ boxedval}}$   $d$  is a boxed value

$$\frac{\text{BVVAL} \quad d \text{ val}}{d \text{ boxedval}}$$

$$\frac{\text{BVROLL} \quad d \text{ boxedval}}{\text{roll}^{\mu \pi . \tau}(d) \text{ boxedval}}$$

$$\frac{\text{BVINJ} \quad d \text{ boxedval}}{\text{inj}_C^\tau(d) \text{ boxedval}}$$

$$\frac{\text{BVARRCAST} \quad \tau_1 \rightarrow \tau_2 \neq \tau_3 \rightarrow \tau_4 \quad d \text{ boxedval}}{d \langle \tau_1 \rightarrow \tau_2 \Rightarrow \tau_3 \rightarrow \tau_4 \rangle \text{ boxedval}}$$

$$\frac{\text{BVRECCAST} \quad \mu \pi . \tau \neq \mu \pi' . \tau' \quad d \text{ boxedval}}{d \langle \mu \pi . \tau \Rightarrow \mu \pi' . \tau' \rangle \text{ boxedval}}$$

$$\frac{\text{BVSUMCAST} \quad \tau \blacktriangleright_+ + \{C_i(\tau_i)\}_{C_i \in \mathcal{C}} \quad \tau' \blacktriangleright_+ + \{C'_i(\tau'_i)\}_{C'_i \in \mathcal{C}'} \quad \tau \neq \tau' \quad d \text{ boxedval}}{d \langle \tau \Rightarrow \tau' \rangle \text{ boxedval}}$$

$$\frac{\text{BVHOLECAST} \quad d \text{ boxedval} \quad \tau \text{ ground}}{d \langle \tau \Rightarrow \emptyset \rangle \text{ boxedval}}$$

$\boxed{d \text{ indet}}$   $d$  is indeterminate

$$\frac{\text{IEHOLE}}{\emptyset_\sigma^u \text{ indet}} \quad \frac{\text{INEHOLE} \quad d \text{ final}}{\langle d \rangle_\sigma^u \text{ indet}}$$

$$\frac{\text{IAPP} \quad d_1 \neq d'_1 \langle \tau_1 \rightarrow \tau_2 \Rightarrow \tau_3 \rightarrow \tau_4 \rangle \quad d_1 \text{ indet} \quad d_2 \text{ final}}{d_1(d_2) \text{ indet}}$$

$$\frac{\text{IROLL} \quad d \text{ indet}}{\text{roll}^{\mu \pi . \tau}(d) \text{ indet}}$$

$$\frac{\text{IUNROLL} \quad d \text{ indet}}{\text{unroll}(d) \text{ indet}}$$

$$\frac{\text{IINJ} \quad d \text{ indet}}{\text{inj}_C^\tau(d) \text{ indet}}$$

$$\frac{\text{IINJHOLE} \quad C \neq \mathbf{C} \quad d \text{ final}}{\text{inj}_C^\tau(d) \text{ indet}}$$

$$\frac{\text{ICASTGROUNDHOLE} \quad d \text{ indet} \quad \tau \text{ ground}}{d \langle \tau \Rightarrow \emptyset \rangle \text{ indet}}$$

$$\begin{array}{c}
\text{ICASTHOLEGROUND} \\
\frac{d \neq d' \langle \tau' \Rightarrow \emptyset \rangle \quad d \text{ indet} \quad \tau \text{ ground}}{d \langle \emptyset \Rightarrow \tau \rangle \text{ indet}}
\end{array}
\quad
\begin{array}{c}
\text{ICASTARR} \\
\frac{\tau_1 \rightarrow \tau_2 \neq \tau_3 \rightarrow \tau_4 \quad d \text{ indet}}{d \langle \tau_1 \rightarrow \tau_2 \Rightarrow \tau_3 \rightarrow \tau_4 \rangle \text{ indet}}
\end{array}
\quad
\begin{array}{c}
\text{ICASTREC} \\
\frac{\mu\pi.\tau \neq \mu\pi'.\tau' \quad d \text{ indet}}{d \langle \mu\pi.\tau \Rightarrow \mu\pi'.\tau' \rangle \text{ indet}}
\end{array}$$

$$\begin{array}{c}
\text{ICASTSUM} \\
\frac{\tau \blacktriangleright_{+=} + \{C_i(\tau_i)\}_{C_i \in \mathcal{C}} \quad \tau' \blacktriangleright_{+=} + \{C'_i(\tau'_i)\}_{C_i \in \mathcal{C}} \quad \tau \neq \tau' \quad d \text{ indet}}{d \langle \tau \Rightarrow \tau' \rangle \text{ indet}}
\end{array}$$

$$\begin{array}{c}
\text{IFAILEDCAST} \\
\frac{d \text{ final} \quad \tau_1 \text{ ground} \quad \tau_2 \text{ ground} \quad \tau_1 \approx \tau_2}{d \langle \tau_1 \Rightarrow \emptyset \rangle \not\Rightarrow \tau_2 \rangle \text{ indet}}
\end{array}$$

$\boxed{d \longrightarrow d'}$   $d$  takes an instruction transition to  $d'$

$$\begin{array}{c}
\text{ITAPP} \\
\frac{[d_2 \text{ final}]}{(\lambda x:\tau.d_1)(d_2) \longrightarrow [d_2/x]d_1}
\end{array}$$

$$\begin{array}{c}
\text{ITUNROLL} \\
\frac{[d \text{ final}]}{\text{unroll}(\text{roll}^{\mu\pi.\tau}(d)) \longrightarrow d}
\end{array}$$

$$\begin{array}{c}
\text{ITAPPCAST} \\
\frac{[d_1 \text{ final}] \quad [d_2 \text{ final}] \quad \tau_1 \rightarrow \tau_2 \neq \tau'_1 \rightarrow \tau'_2}{d_1 \langle \tau_1 \rightarrow \tau_2 \Rightarrow \tau'_1 \rightarrow \tau'_2 \rangle (d_2) \longrightarrow (d_1(d_2 \langle \tau'_1 \Rightarrow \tau_1 \rangle)) \langle \tau_2 \Rightarrow \tau'_2 \rangle}
\end{array}$$

$$\begin{array}{c}
\text{ITUNROLLCAST} \\
\frac{[d \text{ final}] \quad \mu\pi.\tau \neq \mu\pi'.\tau'}{\text{unroll}(d \langle \mu\pi.\tau \Rightarrow \mu\pi'.\tau' \rangle) \longrightarrow \text{unroll}(d) \langle [\mu\pi.\tau/\pi]\tau \Rightarrow [\mu\pi'.\tau'/\pi']\tau' \rangle}
\end{array}
\quad
\begin{array}{c}
\text{ITCASTID} \\
\frac{[d \text{ final}]}{d \langle \tau \Rightarrow \tau \rangle \longrightarrow d}
\end{array}$$

$$\begin{array}{c}
\text{ITCASTSUCCEED} \\
\frac{[d \text{ final}] \quad \tau \text{ ground}}{d \langle \tau \Rightarrow \emptyset \Rightarrow \tau \rangle \longrightarrow d}
\end{array}$$

$$\begin{array}{c}
\text{ITCASTFAIL} \\
\frac{[d \text{ final}] \quad \tau_1 \neq \tau_2 \quad \tau_1 \text{ ground} \quad \tau_2 \text{ ground}}{d \langle \tau_1 \Rightarrow \emptyset \Rightarrow \tau_2 \rangle \longrightarrow d \langle \tau_1 \Rightarrow \emptyset \rangle \not\Rightarrow \tau_2 \rangle}
\end{array}$$

$$\begin{array}{c}
\text{ITGROUND} \\
\frac{[d \text{ final}] \quad \tau \blacktriangleright_{\text{ground}} \tau'}{d \langle \tau \Rightarrow \emptyset \rangle \longrightarrow d \langle \tau \Rightarrow \tau' \Rightarrow \emptyset \rangle}
\end{array}$$

$$\begin{array}{c}
\text{ITEXPAND} \\
\frac{[d \text{ final}] \quad \tau \blacktriangleright_{\text{ground}} \tau'}{d \langle \emptyset \Rightarrow \tau \rangle \longrightarrow d \langle \emptyset \Rightarrow \tau' \Rightarrow \tau \rangle}
\end{array}$$

$$\begin{array}{l}
\text{EvalCtx } \mathcal{E} ::= \circ \mid \mathcal{E}(d) \mid d(\mathcal{E}) \mid \text{roll}^{\mu\pi.\tau}(\mathcal{E}) \mid \text{unroll}(\mathcal{E}) \mid \text{inj}_C^\tau(\mathcal{E}) \mid \langle \mathcal{E} \rangle_\sigma^u \mid \langle \mathcal{E} \rangle_\sigma^u \blacktriangleright \\
\mid \mathcal{E} \langle \tau \Rightarrow \tau \rangle \mid \mathcal{E} \langle \tau \Rightarrow \emptyset \rangle \not\Rightarrow \tau \rangle
\end{array}$$

$\boxed{d = \mathcal{E}\{d'\}}$   $d$  is obtained by placing  $d'$  at the mark in  $\mathcal{E}$

$$\begin{array}{c}
\text{FHOUTER} \quad \text{FHAPP1} \quad \text{FHAPP2} \quad \text{FHROLL} \\
\frac{}{d = \circ\{d\}} \quad \frac{d_1 = \mathcal{E}\{d'_1\}}{d_1(d_2) = \mathcal{E}(d_2)\{d'_1\}} \quad \frac{\textcolor{red}{[d_1 \text{ final}]} \quad d_2 = \mathcal{E}\{d'_2\}}{d_1(d_2) = d_1(\mathcal{E})\{d'_2\}} \quad \frac{d = \mathcal{E}\{d'\}}{\text{roll}^{\mu\pi.\tau}(d) = \text{roll}^{\mu\pi.\tau}(\mathcal{E})\{d'\}} \\
\\
\text{FHUNROLL} \quad \text{FHINJ} \quad \text{FNEHOLEINSIDE} \quad \text{FMHOLEINSIDE} \\
\frac{d = \mathcal{E}\{d'\}}{\text{unroll}(d) = \text{unroll}(\mathcal{E})\{d'\}} \quad \frac{d = \mathcal{E}\{d'\}}{\text{inj}_C^\tau(d) = \text{inj}_C^\tau(\mathcal{E})\{d'\}} \quad \frac{d = \mathcal{E}\{d'\}}{\langle d \rangle_\sigma^u = \langle \mathcal{E} \rangle_\sigma^u\{d'\}} \quad \frac{d = \mathcal{E}\{d'\}}{\langle d \rangle_\sigma^{u\blacktriangleright} = \langle \mathcal{E} \rangle_\sigma^{u\blacktriangleright}\{d'\}} \\
\\
\text{FHCASINSIDE} \quad \text{FHFAILEDCAST} \\
\frac{d = \mathcal{E}\{d'\}}{d\langle \tau_1 \Rightarrow \tau_2 \rangle = \mathcal{E}\langle \tau_1 \Rightarrow \tau_2 \rangle\{d'\}} \quad \frac{d = \mathcal{E}\{d'\}}{d\langle \tau_1 \Rightarrow \emptyset \nRightarrow \tau_2 \rangle = \mathcal{E}\langle \tau_1 \Rightarrow \emptyset \nRightarrow \tau_2 \rangle\{d'\}}
\end{array}$$

$\boxed{d \mapsto d'}$   $d$  steps to  $d'$

$$\begin{array}{c}
\text{STEP} \\
\frac{d = \mathcal{E}\{d_0\} \quad d_0 \longrightarrow d'_0 \quad d' = \mathcal{E}\{d'_0\}}{d \mapsto d'}
\end{array}$$