

Open Regions Foliated by Annulus Orbits

Hazel Brenner

Summer 2023

1 Introduction

The following is a result about the structure of compact three-manifolds admitting fixed point-free $\text{Homeo}_0(S^1)$ actions having only one dimensional and annulus orbits. A fact we will make repeated use of to prove these results is the following:

Remark 1. Let $\pi_{SO(2)}$ denote the projection along the orbits of the rotation subgroup to the base orbifold. Then, since M^3 and the base orbifold are compact Hausdorff spaces, by a classical fact from point-set topology, $\pi_{SO(2)}$ is a closed map.

2 Global structure

Consider the subset of M consisting of 1-dimensional orbits. The set $\text{fix}(G_0)$ is closed, and $\pi_{SO(2)}$ is a closed map, so $\pi_{SO(2)}(\text{fix}(G_0))$ is closed in the base surface. By continuity then $\pi_{SO(2)}^{-1}(\pi_{SO(2)}(\text{fix}(G_0)))$, which is the union of all of the one-dimensional orbits, is closed. The complement is then an open subset of M which decomposes into annulus orbits. It is natural to ask about the structure of this open region. Below are two important structural results:

3 Foliation by Annuli

Proposition 1. Suppose X is a component of the open subset of M^3 in which all orbits are annuli. Then the decomposition of X into orbits is a C^0 foliation by annuli.

Proof sketch. Take a short curve segment through x , build two one parameter families of homeos, giving local product structure (do this explicitly, use flow of a bump vector field on circle). \square

flesh this out with details

4 2D Orbits Compactify Nicely

As a small point of notation, recall the following definition from point set topology:

Definition 1. The *frontier* of a set X , denoted $\text{fr}(X)$ is defined as $\overline{X} - X$

When X is a 2-dimensional orbit, we use the following notation for convenience.

Definition 2. Let A be an annulus orbit, fix a model annulus $f : A \rightarrow S^1 \times I$. Then, denote by $\text{fr}_+(A)$ the set of frontier points which can be accumulated by sequences with increasing I coordinate and $\text{fr}_-(A)$ analogously. Note that $\text{fr}_+(A) \cup \text{fr}_-(A) = \text{fr}(A)$, but they are not necessarily distinct. We can compatibly label subsets of the frontier of a G_0 invariant interval in an annulus A . For notational consistency, when X is a Möbius band orbit, let $\text{fr}_+(X) = \text{fr}_-(X) = \text{fr}(X)$.

For the following two proofs, we want to make use of a kind of canonical coordinates on interval orbits of G_0 .

Remark 2. [?] Suppose A is an annulus orbit of $\text{Homeo}_0(S^1)$ and $I \subset A$ is an interval orbit of G_0 . Then, we can define two homeomorphisms $\varphi, \psi : (0, 1) \rightarrow I$ by $\varphi(\theta) = \text{fix}(G_0 \cap G_\theta)$ and $\psi(\theta) = \text{fix}(G_0 \cap G_{-\theta})$. One of these maps will send monotonic sequences that converge to 1 to sequences which converge to something in $\text{fr}_+(I)$, we will denote the inverse of this map as f_I , which we will refer to as the *canonical homeomorphism between I and $S^1 - \{0\}$* . Note, the same construction works for the one interval orbit in a Möbius band orbit.

The following results are true for 2-dimensional orbits of any $\text{Homeo}_0(S^1)$ action without fixed points on a closed 3-manifold.

Proposition 2. Suppose X is an 2-dimensional orbit, then $\text{fr}(X)$ consists of 1-dimensional orbits.

Proof. Trivially, $\text{fr}(X)$ contains no 3-dimensional orbit.

Suppose toward a contradiction that $\text{fr}(X)$ contains some 2-dimensional orbit X' . Now, consider the restriction of the $\text{Homeo}_0(S^1)$ action to the point stabilizer subgroup G_0 . There is some interval orbit I in X . Since $\pi_{SO(2)}$ is continuous and closed,

$$\pi_{SO(2)}(\bar{I}) = \overline{\pi_{SO(2)}(I)} = \overline{\pi_{SO(2)}(A)} = \pi_{SO(2)}(\bar{A}).$$

In particular, this means that $\text{fr}(I)$ must nontrivially intersect X' . The closure of every G_0 orbit in X' contains at least one interval orbit I' , so $I' \subseteq \text{fr}(I)$. In particular, there is some sequence of points $\{x_n\} \subset I$ converging to a point $x \in I'$.

From the remark, there are continuous bijections $f : I \rightarrow S^1 - \{0\}$ and $g : I' \rightarrow S^1 - \{0\}$ given by mapping the unique fixed point of $G_\theta \cap G_0$ on I to $\theta \in S^1 - \{0\}$ and similarly for I' . Then, up to a subsequence, $\{f(x_n)\}$ is a monotonic sequence converging to 0. So we can choose a monotonic element $\phi \in G_0$ satisfying $\phi(f(x_k)) = f(x_{k+1})$. In particular, we must have that $\{\rho(\phi)(x_n)\}$ converges to x , but by monotonicity, $\rho(\phi)(x) \neq x$, violating continuity, which is a contradiction. Thus $\text{fr}(A) \cap X = \emptyset$ \square

Note that, in particular if X is an annulus orbit in a component foliated by annuli, this implies that $\text{fr}(X)$ is a subset of the frontier of the component.

We can in fact say something stronger though.

Proposition 3. Suppose X is a 2-dimensional orbit. Then, $\text{fr}_+(X)$ and $\text{fr}_-(X)$ are (not necessarily distinct) one-dimensional orbits.


Proof. Suppose that σ and σ' are invariant circles in (WLOG) $\text{fr}_+(X)$ with respective G_0 fixed points x and x' . Let I be a G_0 -invariant interval in X . By the same reasoning as the previous result, $x, x' \in \text{fr}(I)$, so let $\{a_k\}$ and $\{b_k\}$ be sequences in I representing x and x' . Since both sequences are monotonic in I , there is a homeomorphism $f \in G_0$ such that $\rho(f)(a_k) = b_k$. By continuity, this implies that $\{b_k\}$ converges to $\rho(f)(x)$, which is x , since $x \in \text{fix}(G_0)$. Thus, $x = x'$; moreover, $\sigma = \sigma'$ since every fixed point of G_0 is part of a unique 1-dimensional orbit of $\text{Homeo}_0(S^1)$. \square

This result can be rephrased in a slightly more concrete way.

Corollary 1. When X is an annulus orbit, \bar{X} is either an invariant closed annulus with boundary circles $\text{fr}_+(A)$ and $\text{fr}_-(A)$ or an invariant T^2 when $\text{fr}_-(A) = \text{fr}_+(A)$.

When X is a Möbius band orbit, \bar{X} is an invariant closed Möbius band, with invariant boundary circle $\text{fr}(X)$.

Notes

 flesh this out with details 1