


Cardiac arrhythmia data set

- Detect the presence of cardiac arrhythmia based on various ECG data and other attributes such as age and weight ($n = 451, p = 194$).



H. A. Guvenir et al. (1998). *UCI Machine Learning Repository: Arrhythmia Data Set*. URL: <https://archive.ics.uci.edu/ml/datasets/Arrhythmia>

Cardiac arrhythmia data set - Model fit


- Fit an l-prior probit model using Canonical and FBM kernels. The full data set takes about 35 seconds.

```
R> mod <- iprior(y, X, kernel = "FBM")
```


- Compare against popular classifiers: 1) k -nearest neighbours; 2) support vector machine; 3) Gaussian process classification; 4) random forests; 5) nearest shrunk centroids (Tibshirani et al. 2003); and 6) L-1 penalised logistic regression.
- Experiment set-up:
 - ▶ Form training set by sub-sampling $n_{\text{sub}} \in \{50, 100, 200\}$ data points.
 - ▶ Use remaining data as test set.
 - ▶ Fit model on training set and obtain test error rates.
 - ▶ Repeat 100 times.

T. I. Cannings and R. J. Samworth (2017). "Random-projection ensemble classification". *J. R. Stat. Soc. Ser. B: Stat. Methodol (w. discussion)*, to appear

Cardiac arrhythmia data set - Results



Meta-analysis of smoking cessation



- Data from 27 separate smoking cessation studies, where participants subjected to nicotine gum treatment or placed in control group.
- Some summary statistics:

	Min.	Avg.	Max.	Prop. quit	Odds quit
Control	20	101	617	0.207	0.261
Treated	21	117	600	0.320	0.470

- Raw odds ratio: 1.801.
- Random-effects analysis using a multilevel logistic model estimates this odds ratio as 1.768.

A. Skrondal and S. Rabe-Hesketh (2004). *Generalized Latent Variable Modeling: Multilevel, Longitudinal, and Structural Equation Models*. Chapman & Hall/CRC, §9.5

Meta-analysis of smoking cessation - model

- Let $i = 1, \dots, n_j$ index the patients in study group $j \in 1, \dots, 27$.
- Denote y_{ij} as the binary response variable indicating Quit (1) or Remain (0), and x_{ij} as patient ij 's treatment group indicator.
- Model binary data using l-probit model

$$\begin{aligned}\Phi^{-1}(p_{ij}) &= f(x_{ij}, j) \\ &= f_1(x_{ij}) + f_2(j) + f_{12}(x_{ij}, j)\end{aligned}$$

with $f_1, f_2 \in$ Pearson RKHS, and $f_{12} \in$ ANOVA RKHS.

	Model	Lower bound	Brier score	No. of RKHS param.
1	f_1	-3210.79	0.0311	1
2	$f_1 + f_2$	-3097.24	0.0294	2
3	$f_1 + f_2 + f_{12}$	-3091.21	0.0294	2

Meta-analysis of smoking cessation - results

