

A beginner's guide to variational inference

Haziq Jamil

Social Statistics
London School of Economics and Political Science

1 February 2018

Social Statistics Meeting

<http://socialstats.haziqj.ml>

Outline

① Introduction

② Examples

③ Discussion

- Exponential families

- Zero-forcing vs Zero-avoiding

- Quality of approximation

- Advanced topics

① Introduction

② Examples

③ Discussion

① Introduction

② Examples

③ Discussion

Exponential families

- For the mean-field variational method, suppose that each complete conditional is in the exponential family:

$$p(\mathbf{z}^{(j)} | \mathbf{z}_{-j}, \mathbf{y}) = h(\mathbf{z}^{(j)}) \exp(\eta_j(\mathbf{z}_{-j}, \mathbf{y}) \cdot \mathbf{z}^{(j)} - A(\eta_j)).$$

Exponential families

- For the mean-field variational method, suppose that each complete conditional is in the exponential family:

$$p(\mathbf{z}^{(j)} | \mathbf{z}_{-j}, \mathbf{y}) = h(\mathbf{z}^{(j)}) \exp(\eta_j(\mathbf{z}_{-j}, \mathbf{y}) \cdot \mathbf{z}^{(j)} - A(\eta_j)).$$

- Then, from (??),

$$\begin{aligned} \tilde{q}_j(\mathbf{z}^{(j)}) &\propto \exp(E_{-j}[\log p(\mathbf{z}^{(j)} | \mathbf{z}_{-j}, \mathbf{y})]) \\ &= \exp(\log h(\mathbf{z}^{(j)}) + E[\eta_j(\mathbf{z}_{-j}, \mathbf{y})] \cdot \mathbf{z}^{(j)} - E[A(\eta_j)]) \\ &\propto h(\mathbf{z}^{(j)}) \exp(E[\eta_j(\mathbf{z}_{-j}, \mathbf{y})] \cdot \mathbf{z}^{(j)}) \end{aligned}$$

is also in the same exponential family.

Exponential families

- For the mean-field variational method, suppose that each complete conditional is in the exponential family:

$$p(\mathbf{z}^{(j)} | \mathbf{z}_{-j}, \mathbf{y}) = h(\mathbf{z}^{(j)}) \exp(\eta_j(\mathbf{z}_{-j}, \mathbf{y}) \cdot \mathbf{z}^{(j)} - A(\eta_j)).$$

- Then, from (??),

$$\begin{aligned} \tilde{q}_j(\mathbf{z}^{(j)}) &\propto \exp(E_{-j}[\log p(\mathbf{z}^{(j)} | \mathbf{z}_{-j}, \mathbf{y})]) \\ &= \exp(\log h(\mathbf{z}^{(j)}) + E[\eta_j(\mathbf{z}_{-j}, \mathbf{y})] \cdot \mathbf{z}^{(j)} - E[A(\eta_j)]) \\ &\propto h(\mathbf{z}^{(j)}) \exp(E[\eta_j(\mathbf{z}_{-j}, \mathbf{y})] \cdot \mathbf{z}^{(j)}) \end{aligned}$$

is also in the same exponential family.

- C.f. Gibbs conditional densities.

Exponential families

- For the mean-field variational method, suppose that each complete conditional is in the exponential family:

$$p(\mathbf{z}^{(j)} | \mathbf{z}_{-j}, \mathbf{y}) = h(\mathbf{z}^{(j)}) \exp(\eta_j(\mathbf{z}_{-j}, \mathbf{y}) \cdot \mathbf{z}^{(j)} - A(\eta_j)).$$


- Then, from (??),

$$\begin{aligned}\tilde{q}_j(\mathbf{z}^{(j)}) &\propto \exp(E_{-j}[\log p(\mathbf{z}^{(j)} | \mathbf{z}_{-j}, \mathbf{y})]) \\ &= \exp(\log h(\mathbf{z}^{(j)}) + E[\eta_j(\mathbf{z}_{-j}, \mathbf{y})] \cdot \mathbf{z}^{(j)} - E[A(\eta_j)]) \\ &\propto h(\mathbf{z}^{(j)}) \exp(E[\eta_j(\mathbf{z}_{-j}, \mathbf{y})] \cdot \mathbf{z}^{(j)})\end{aligned}$$

is also in the same exponential family.


- C.f. Gibbs conditional densities.
- ISSUE:** What if not in exponential family? Importance sampling or Metropolis sampling.

Non-convexity of ELBO



- CAVI only guarantees converges to a local optimum.
- Multiple local optima may exist.

Non-convexity of ELBO



- CAVI only guarantees converges to a local optimum.
- Multiple local optima may exist.

Zero-forcing vs Zero-avoiding

- Back to the KL divergence:

$$\text{KL}(q\|p) = \int \log \frac{q(\mathbf{z})}{p(\mathbf{z}|\mathbf{y})} q(\mathbf{z}) d\mathbf{z}$$


- $\text{KL}(q\|p)$ is large when $p(\mathbf{z}|\mathbf{y})$ is close to zero, unless $q(\mathbf{z})$ is also close to zero (*zero-forcing*).
- **ISSUE:** What about other measures of closeness? For instance,

$$\text{KL}(p\|q) = \int \log \frac{p(\mathbf{z}|\mathbf{y})}{q(\mathbf{z}|\mathbf{y})} p(\mathbf{z}|\mathbf{y}) d\mathbf{z}.$$


- This gives the Expectation Propagation (EP) algorithm.
- It is *zero-avoiding*, because $\text{KL}(p\|q)$ is small when both $p(\mathbf{z}|\mathbf{y})$ and $q(\mathbf{z})$ are non-zero.

Zero-forcing vs Zero-avoiding (cont.)


— $p(z)$




Zero-forcing vs Zero-avoiding (cont.)




Zero-forcing vs Zero-avoiding (cont.)



Zero-forcing vs Zero-avoiding (cont.)




Distortion of higher order moments



- Consider $\mathbf{z} = (z_1, z_2)^\top \sim N_2(\boldsymbol{\mu}, \boldsymbol{\Psi}^{-1})$, $\text{Cov}(z_1, z_2) \neq 0$.

Distortion of higher order moments




- Consider $\mathbf{z} = (z_1, z_2)^\top \sim N_2(\boldsymbol{\mu}, \boldsymbol{\Psi}^{-1})$, $\text{Cov}(z_1, z_2) \neq 0$.
- Approximating $p(\mathbf{z})$ by $q(\mathbf{z}) = q(z_1)q(z_2)$ yields

$$\tilde{q}(z_1) = N(z_1|\mu_1, \boldsymbol{\Psi}_{11}^{-1}) \quad \text{and} \quad \tilde{q}(z_2) = N(z_2|\mu_2, \boldsymbol{\Psi}_{22}^{-1})$$

and by definition, $\text{Cov}(z_1, z_2) = 0$ under \tilde{q} .

Distortion of higher order moments



- Consider $\mathbf{z} = (z_1, z_2)^\top \sim N_2(\boldsymbol{\mu}, \boldsymbol{\Psi}^{-1})$, $\text{Cov}(z_1, z_2) \neq 0$.
- Approximating $p(\mathbf{z})$ by $q(\mathbf{z}) = q(z_1)q(z_2)$ yields

$$\tilde{q}(z_1) = N(z_1|\mu_1, \boldsymbol{\Psi}_{11}^{-1}) \quad \text{and} \quad \tilde{q}(z_2) = N(z_2|\mu_2, \boldsymbol{\Psi}_{22}^{-1})$$

and by definition, $\text{Cov}(z_1, z_2) = 0$ under \tilde{q} .

- This leads to underestimation of variances (widely reported in the literature—Zhao and Marriott 2013).

Quality of approximation

- Variational inference converges to a different optimum than ML, except for certain models (Gunawardana and Byrne 2005).

Quality of approximation


- Variational inference converges to a different optimum than ML, except for certain models (Gunawardana and Byrne 2005).
- But not much can be said about the quality of approximation.

Quality of approximation

- Variational inference converges to a different optimum than ML, except for certain models (Gunawardana and Byrne 2005).
- But not much can be said about the quality of approximation.
- Statistical properties not well understood—what is its statistical profile relative to the exact posterior?

Quality of approximation

- Variational inference converges to a different optimum than ML, except for certain models (Gunawardana and Byrne 2005).
- But not much can be said about the quality of approximation.
- Statistical properties not well understood—what is its statistical profile relative to the exact posterior?
- Speed trumps accuracy?



Advanced topics

- Local variational bounds
 - ▶ Not using the mean-field assumption.
 - ▶ Instead, find a bound for the marginalising integral \mathcal{I} .
 - ▶ Used for Bayesian logistic regression as follows:

$$\mathcal{I} = \int \text{expit}(\mathbf{x}^\top \beta) p(\beta) \, \mathrm{d}\beta \geq \int f(\mathbf{x}^\top \beta, \xi) p(\beta) \, \mathrm{d}\beta.$$

Advanced topics

- Local variational bounds
 - ▶ Not using the mean-field assumption.
 - ▶ Instead, find a bound for the marginalising integral \mathcal{I} .
 - ▶ Used for Bayesian logistic regression as follows:

$$\mathcal{I} = \int \text{expit}(\mathbf{x}^\top \beta) p(\beta) \, \mathrm{d}\beta \geq \int f(\mathbf{x}^\top \beta, \xi) p(\beta) \, \mathrm{d}\beta.$$

- Stochastic variational inference
 - ▶ VI on its own doesn't offer much computational advantages.
 - ▶ Use ideas from stochastic optimisation—gradient based improvement of ELBO from subsamples of the data.
 - ▶ Scales to massive data.

Advanced topics

- Local variational bounds
 - ▶ Not using the mean-field assumption.
 - ▶ Instead, find a bound for the marginalising integral \mathcal{I} .
 - ▶ Used for Bayesian logistic regression as follows:

$$\mathcal{I} = \int \text{expit}(\mathbf{x}^\top \beta) p(\beta) \, \mathrm{d}\beta \geq \int f(\mathbf{x}^\top \beta, \xi) p(\beta) \, \mathrm{d}\beta.$$

- Stochastic variational inference
 - ▶ VI on its own doesn't offer much computational advantages.
 - ▶ Use ideas from stochastic optimisation—gradient based improvement of ELBO from subsamples of the data.
 - ▶ Scales to massive data.
- Black box variational inference
 - ▶ Beyond exponential families and model-specific derivations.

End

Thank you!

Slides and source code are made available at: <http://socialstats.haziqj.ml>

References I

- Gunawardana, A. and W. Byrne (2005). “Convergence theorems for generalized alternating minimization procedures”. *Journal of machine learning research* 6.Dec, pp. 2049–2073.
- Zhao, H. and P. Marriott (2013). “Diagnostics for Variational Bayes approximations”. *arXiv: 1309.5117*.

④ Additional material