

Non-Hermitian Skin Effect: A Primer

Sagnik Seth,^{1,*} Bihan Bhowmick,^{2,†} and Himadri Barman^{3,‡}

¹*IISER Kolkata, Kolkata, India*

²*IISER Pune, Pune, India*

³*Department of Physics, Zhejiang University, Hangzhou 310027, China*

Hermiticity is a crucial condition for the observables to be real (hence measurable) in our conventional quantum physics. However, the conventional quantum physics deals with the ideal scenario where the system is isolated from its environment. As soon as we take account of the interaction with the environment, we enter the generic world of non-equilibrium quantum physics where Hermiticity is not a required condition. The recent hunt of topological insulators for fault-tolerant quantum computation has opened the gate for studying the physics of non-Hermiticity into those materials. It has been recently discovered that non-Hermiticity induces an exotic phenomenon dubbed the *skin effect* which entails asymmetric localization of quantum states. In our article, we provide an introduction to this phenomenon and physics behind this with a flavor of students-friendly pedagogy.

CONTENTS

I. Introduction	3
II. Hatano-Nelson model: A flavor of non-Hermiticity in a lattice	3
III. SSH model	3
IV. Reciprocal space representation	4
V. Digression: Berry phase	5
VI. Digression: Two level system	6
VII. Back to the SSH model	6
Dispersion:	6
Other Nomenclatures:	7
VIII. Winding number (ν): The rubber band analogy	7
Formula 4	8
IX. Edge states	8
X. Non-Hermitian Topological Insulators	9
XI. Exceptional points	9
XII. Edge states	9
XIII. Non-Bloch Band Theory	9
XIV. Project proposal	9
References	10
References	10

I. INTRODUCTION

II. HATANO-NELSON MODEL: A FLAVOR OF NON-HERMITICITY IN A LATTICE

The Hatano-Nelson model (non-disordered) is the simplest tight-binding lattice model where the left and right-moving hopping amplitudes are unequal (*non-reciprocal*). The Hamiltonian for spinless particles in one dimension reads

$$\hat{H}_{\text{HN}} = \sum_{i=0}^{L-1} [t_R c_i^\dagger c_{i+1} + t_L c_{i+1}^\dagger c_i], \quad (1)$$

where L is the lattice size; t_R and t_L are the right and left hopping amplitudes; c_i and c_i^\dagger are the particle annihilation and creation operators at lattice site i , respectively.

We can express the annihilation operator in terms of Fourier components as

$$c_i \equiv \frac{1}{\sqrt{L}} \sum_{k=0}^{L-1} e^{ikx_i} c_k. \quad (2)$$

(where x_i is the coordinate or position of site i) and rewrite the Hamiltonian in the Fourier (k) space:

$$\begin{aligned} \hat{H}_{\text{HN}} &= \sum_{i=0}^{L-1} \sum_{k=0}^{L-1} \sum_{k'=0}^{L-1} [t_R c_k^\dagger c_k e^{ik'((x_{i+1}-x_i))} + t_L c_k^\dagger c_k e^{ik'(x_i-x_{i+1})}] \\ &= \sum_i \sum_k [t_R c_k^\dagger c_k e^{ik((x_{i+1}-x_i))} + t_L c_k^\dagger c_k e^{i(x_i-x_{i+1})}] \quad [\text{Dropped details of summation indices for clarity}] \\ &= [t_R e^{ika} + t_L e^{-ika}] c_k^\dagger c_k \end{aligned} \quad (3)$$

where we utilize $x_{i+1} - x_i = a$, a being the lattice constant (spacing between two consecutive lattice sites).




FIG. 1: Hatano-Nelson chain with non-reciprocal hopping parameters t_R and t_L .

III. SSH MODEL

Let us start with the simplest topological model Hamiltonian known as the Su-Schrieffer-Heeger (SSH) model which has been used to describe polyacetylene or bi-partite/dimerized (Peierls) lattice ¹. The Hamiltonian reads

$$\begin{aligned} \hat{H}_{\text{SSH}} &= \sum_{I=1}^L [t + (-1)^I \delta] c_I^\dagger c_{I+1} \\ &= \sum_{i=0}^{L-1} (t + \delta) c_{A i}^\dagger c_{B i} + \sum_i (t - \delta) c_{A i+1}^\dagger c_{B i} + \text{h.c.} \\ &\equiv \sum_i [t_+ c_{A i}^\dagger c_{B i} + t_- c_{A i+1}^\dagger c_{B i} + \text{h.c.}] . \end{aligned} \quad (1)$$

Here the lattice consists of two different sublattice sites A and B inside each primitive cell as they associate with two different hopping amplitudes, namely $t_- = t - \delta$ (representing longer single bond having lesser kinetic energy or hopping for $\delta > 0$) and $t_+ = t + \delta$ (representing shorter double bond having higher kinetic energy or hopping for $\delta > 0$). t_+ and t_- are dubbed *intracell* and *intercell* hopping respectively.




FIG. 2: Su-Schrieffer-Heeger (SSH) chains in two different configurations.

Now the Hamiltonian can be written in the matrix form $\hat{H} = \Psi^\dagger \mathbf{H} \Psi$ with the basis (or spinor) $\Psi^\dagger = [c_{A1}^\dagger c_{B1}^\dagger c_{A2}^\dagger \cdots]$. Then the Hamiltonian matrix becomes

$$\mathbf{H} = \begin{bmatrix} 0 & t_+ & 0 & \cdots & 0 \\ t_+^* & 0 & t_- & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & t_{L-1}^* & 0 \end{bmatrix} \quad (2)$$

where the $(L-1)$ -th hopping amplitude t_{L-1} is either t_+ or t_- depending on the value of L (even or odd).

Note that the above matrix represents the open boundary condition (OBC) where there exists no left-side hopping from cell 1 (considering cell 2 is on the right of cell 1) and no right-side hopping from cell L . However, if the periodic boundary condition (PBC) is applied, i.e. left-side hopping from cell 1 is the last cell L and right-side hopping from cell L finds the cell 1 back, then the Hamiltonian becomes

$$\mathbf{H} = \begin{bmatrix} 0 & t_+ & 0 & \cdots & t_L^* \\ t_+^* & 0 & t_- & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ t_L & 0 & 0 & t_{L-1} & 0 \end{bmatrix} \quad (3)$$

To see a demonstration, take a simple example of a 4-site (2-cell) lattice (two A -sites + two B -sites) with OBC:

$$\mathbf{H}_{4 \times 4} = \begin{bmatrix} 0 & t_+ & 0 & 0 \\ t_+^* & 0 & t_- & 0 \\ 0 & t_-^* & 0 & t_+ \\ 0 & 0 & t_+^* & 0 \end{bmatrix} \quad (4)$$

Then

$$\begin{aligned} \hat{H}_{4 \times 4} &= \Psi^\dagger \mathbf{H}_{4 \times 4} \Psi \\ &= \begin{bmatrix} c_{A1}^\dagger & c_{B1}^\dagger & c_{A2}^\dagger & c_{B2}^\dagger \end{bmatrix} \begin{bmatrix} 0 & t_+ & 0 & 0 \\ t_+^* & 0 & t_- & 0 \\ 0 & t_-^* & 0 & t_+ \\ 0 & 0 & t_+^* & 0 \end{bmatrix} \begin{bmatrix} c_{A1} \\ c_{B1} \\ c_{A2} \\ c_{B2} \end{bmatrix} \\ &= \begin{bmatrix} c_{A1}^\dagger & c_{B1}^\dagger & c_{A2}^\dagger & c_{B2}^\dagger \end{bmatrix} \begin{bmatrix} t_+ c_{B1} \\ t_+^* c_{A1} + t_- c_{A2} \\ t_-^* c_{B1} + t_+ c_{B2} \\ t_+^* c_{A2} \end{bmatrix} \\ &= t_+ c_{A1}^\dagger c_{B1} + t_+^* c_{B1}^\dagger c_{A1} + t_- c_{B1}^\dagger c_{A2} \\ &\quad + t_-^* c_{A2}^\dagger c_{B1} + t_+ c_{A2}^\dagger c_{B2} + t_+^* c_{B2}^\dagger c_{A2}. \end{aligned} \quad (5)$$

IV. RECIPROCAL SPACE REPRESENTATION

Now we want to write the Hamiltonian in the reciprocal (k) space. Again we can define the Fourier component of an annihilation operator:

$$c_{\alpha k} \equiv \frac{1}{\sqrt{L}} \sum_{i=1}^L e^{-ik \cdot x_i} c_{\alpha i}; \quad \alpha = A, B. \quad (1)$$

[Note: We have total $2L$ lattice sites consisting of L number of A and B sublattices.]

Considering lattice coordinate at site i to be x_i with lattice spacing a (i.e., nearest neighbor site's position $x_{i+1} = x_i + a$), we can rewrite the Hamiltonian in Eq. (1):

$$\begin{aligned}
\hat{H} &= \sum_i \sum_{kk'} \left[\left\{ t_+ c_{Ak}^\dagger c_{Bk} e^{ikx_i} e^{-ik'x_i} + \text{h.c.} \right\} \right. \\
&\quad \left. + \left\{ t_- c_{Ak}^\dagger c_{Bk} e^{ik(x_i+a)} e^{-ik'x_i} + \text{h.c.} \right\} \right] \\
&= \sum_{kk'} \left[\left\{ t_+ c_{Ak}^\dagger c_{Bk} \delta_{kk'} + \text{h.c.} \right\} \right. \\
&\quad \left. + \left\{ t_- c_{Ak}^\dagger c_{Bk} \delta_{kk'} e^{-ik'x_i} + \text{h.c.} \right\} \right] \\
&\quad [\text{use } \sum_i e^{i(k-k')x_i} = \delta_{kk'}] \\
&= \sum_k [\{ t_+ c_{Ak}^\dagger c_{Bk} + t_+^* c_{Bk}^\dagger c_{Ak} \} \\
&\quad + \{ t_- c_{Ak}^\dagger c_{Bk} e^{-ika} + t_-^* c_{Bk}^\dagger c_{Ak} e^{ika} \}] \\
&\equiv \sum_k H_k.
\end{aligned} \tag{2}$$

Thus we get the momentum (k) dependent Hamiltonian (for simplicity, we consider the hopping parameters to be real, i.e. $t_+^* = t_+$, $t_-^* = t_-$)

$$H_k = t_+ (c_{Ak}^\dagger c_{Bk} + c_{Bk}^\dagger c_{Ak}) + t_- (c_{Ak}^\dagger c_{Bk} e^{-ika} + c_{Bk}^\dagger c_{Ak} e^{ika}) \tag{3}$$

$$= (t_+ + t_- e^{-ika}) c_{Ak}^\dagger c_{Bk} + (t_+ + t_- e^{ika}) c_{Bk}^\dagger c_{Ak}. \tag{4}$$

Like what we did in the direct lattice representation case, we choose a 2-dimensional spinor $\Psi_k^\dagger \equiv [c_{Ak}^\dagger \ c_{Bk}^\dagger]$ and write H_k in a 2×2 matrix form: $H_k = \Psi_k^\dagger \mathbf{H}_k \Psi_k$ where

$$\mathbf{H}_k \equiv \begin{bmatrix} 0 & t_+ + t_- e^{-ika} \\ t_+ + t_- e^{ika} & 0 \end{bmatrix}. \tag{5}$$

Then Eq. (4) can be written in the following form:

$$\mathbf{H}_k = d_x \sigma_x + d_y \sigma_y = \vec{d}(k) \cdot \vec{\sigma} \tag{6}$$

where $d_x(k) \equiv t_+ + t_- \cos ka$, $d_y(k) \equiv t_- \sin ka$; $\sigma_x = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ and $\sigma_y = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}$.

V. DIGRESSION: BERRY PHASE

The Bloch wave function is not unique: $|u_{\mathbf{k}}\rangle \rightarrow e^{i\phi(\mathbf{k})} |u_{\mathbf{k}}\rangle$ is invariant under the gauge transformation $\mathbf{A} \rightarrow \mathbf{A} + \nabla_{\mathbf{k}}\phi(\mathbf{k})$ where

$$\mathbf{A} = i\langle u_{\mathbf{k}} | \nabla_{\mathbf{k}} u_{\mathbf{k}} \rangle \tag{1}$$

is known as the *Berry connection* (analogous to the magnetic vector potential). For any close loop C in the \mathbf{k} -space, the Berry phase can be defined as

$$\gamma_C = \oint_C \mathbf{A} \cdot d\mathbf{k} = \int_S \mathcal{F} d^2\mathbf{k} \tag{2}$$

where $\mathcal{F} \equiv \nabla \times \mathbf{A}$ is known as the *Berry curvature*.

For a TLS Hamiltonian $H(\mathbf{k}) = \vec{d}(\mathbf{k}) \cdot \vec{\sigma}$, Berry showed? [*to be shown]

$$\gamma_C = \Omega/2 \tag{3}$$

where Ω is the solid angle swept out by $\hat{d}(\mathbf{k})$.

VI. DIGRESSION: TWO LEVEL SYSTEM

In general a two-level Hamiltonian can be written as

$$\hat{H}_0 = E_1|1\rangle\langle 1| + E_2|2\rangle\langle 2| \quad (1)$$

The Hamiltonian in the matrix form becomes

$$H_0 = \begin{bmatrix} E_1 & 0 \\ 0 & E_2 \end{bmatrix} \quad (2)$$

Now if we turn on an interaction V between state $|1\rangle$ and $|2\rangle$, we get an interacting Hamiltonian

$$\hat{H} = E_1|1\rangle\langle 1| + E_2|2\rangle\langle 2| + V|1\rangle\langle 2| + V^*|2\rangle\langle 1| \quad (3)$$

which gets the matrix form

$$\mathbf{H} = \begin{bmatrix} E_1 & V \\ V^* & E_2 \end{bmatrix} \quad (4)$$

The eigenvalue equation becomes

$$(E_1 - E)(E_2 - E) - |V|^2 = 0 \quad (5)$$

which yields two eigenvalues E_+ and E_- given by

$$E_{\pm} = (E_1 + E_2) \pm \sqrt{(E_1 - E_2)^2 + 4|V|^2}. \quad (6)$$

Suppose we have the following Hamiltonian

$$\mathbf{H}(k) = \begin{bmatrix} h_0 + h_z & h_x - ih_y \\ h_x + ih_y & h_0 - h_z \end{bmatrix} \quad (7)$$

Then the Hamiltonian can be written as

$$H(k) = h^\mu(k)\sigma_\mu = h_0(k)\mathbb{1} + \vec{h}(k) \cdot \vec{\sigma} \quad (8)$$

where $\mathbb{1}$ is the 2×2 unity matrix.

With $\text{Tr}(H) = 2h_0$ and $\det(H) = h_0^2 - h^2$, one finds the eigenvalues

$$E_{\pm} = h_0 \pm h(k) \quad (9)$$

where

$$h(k) = \|\vec{h}(k)\| = \sqrt{h_x^2 + h_y^2 + h_z^2}. \quad (10)$$

The corresponding normalized eigenvectors are (up to a phase factor)

$$u_{\pm}(k) = \frac{1}{\sqrt{1 + (h_z + E_{\pm}^2)/(h_x^2 + h_y^2)}} \begin{bmatrix} E_{\pm}/(h_x + ih_y) \\ 1 \end{bmatrix}. \quad (11)$$

VII. BACK TO THE SSH MODEL

Dispersion:

Now if we compare our k -space SSH Hamiltonian in Eq. (6), we notice h_0 and $\vec{h}(k) = \vec{d}(k)$. Therefore from Eq. (9)

$$E_k = \pm |\mathbf{d}(k)| = \pm \sqrt{(t_+ + t_- \cos ka)^2 + t_-^2 \sin^2 ka} \quad (1)$$

$$\begin{aligned} &= \pm \sqrt{(t_+ - t_-)^2 + 4t_+ t_- \cos^2(ka/2)} \\ &= \pm \sqrt{(4\delta^2 + 4t_+ t_- \cos^2(ka/2))}. \end{aligned} \quad (2)$$

with $d_x(k) \equiv t_+ + t_- \cos ka$, $d_y(k) \equiv t_- \sin ka$.

Other Nomenclatures:

In many literatures, the intracell hopping $t_+ = t + \delta$ is denoted by v and intercell hopping $t_- = t - \delta$ is denoted by w . Now onward, we shall follow this notation and according to it,

$$d_x(k) \equiv v + w \cos ka; \quad d_y(k) \equiv w \sin ka \quad (3)$$


and

$$E(k) = \pm \sqrt{v^2 + w^2 + 2vw \cos ka}.$$

From Eq. (3), we get

$$(d_x - v)^2 + d_y^2 = w^2 \quad (4)$$

which describes a circle of radius w , centered at $(v, 0)$.



:

2

VIII. WINDING NUMBER (ν): THE RUBBER BAND ANALOGY


Winding number is a many times a curve winds around a specific point z_0 .


Our case: $z_0 = (d_x = 0, d_y = 0)$ $\nu = 1$ is **topologically non-trivial!**

The winding number can be mathematically formulated as [**Check derivation!**]:

$$\nu = \frac{1}{2\pi} \int_{-\pi}^{\pi} dk \left(\hat{\mathbf{d}}(k) \times \frac{d}{dk} \hat{\mathbf{d}}(k) \right)_z. \quad (1)$$

When the intracell hopping dominates over the intercell hopping (i.e. $v > w$), the winding number $\nu = 0$. On the other hand, when $v < w$, $\nu = 1$. to change ν , we need to change the path of $\mathbf{d}(k)$





Formula 4

Since $E(k) = \pm|\mathbf{d}(k)|$, we have

$$\begin{aligned} \ln E(k) &= \ln |\mathbf{d}(k)| + i \arg(\mathbf{d}(k)) . \\ \Rightarrow \partial_k \ln E(k) &= \partial_k \ln |\mathbf{d}(k)| + i \partial_k \arg(\mathbf{d}(k)) . \end{aligned} \quad (2)$$

Now we know

$$\nu = \frac{1}{2\pi} \oint_C \partial_k \arg(\mathbf{d}(k)) \quad [\text{see above}] . \quad (3)$$

Also,

$$\nu = \frac{1}{2\pi i} \oint_C dk \partial_k \ln E_k . \quad (4)$$

Also,


$$\nu = \frac{i}{2\pi} \int_{C_\beta} dq q^{-1} . \quad (5)$$

IX. EDGE STATES

2 degenerate $E = 0$ bands

- Edge states function as connectors/interfaces between 2 topological distinct phases [trivial ($\nu = 0$) and non-trivial ($\nu = 1$)] \Rightarrow Bulk-boundary correspondence (BBC).

1. Shen: Topological Insulator and Dirac Equation in Condensed Matter
2. http://optics.szfki.kfki.hu/~asboth/topins_course/2015-09-24-ELTE-Topins.pdf
- 3.



X. NON-HERMITIAN TOPOLOGICAL INSULATORS

Hamiltonian in direct space

$$\begin{aligned} \hat{H} = & B \sum_i (t_1 - \gamma/2) c_{Ai}^\dagger c_{Bi} + \sum_i (t_1 + \gamma/2) c_{Bi}^\dagger c_{Ai} \\ & + \sum_i t_2 [c_{A i+1}^\dagger c_{Bi} + c_{Bi}^\dagger c_{A i+1}] + \sum_i t_3 [c_{Ai}^\dagger c_{B i+1} + c_{B i+1}^\dagger c_{Ai}]. \end{aligned} \quad (1)$$

2

XI. EXCEPTIONAL POINTS

Complex eigenenergies

XII. EDGE STATES

Edge states no longer symmetric


- Even the bulk gets ‘edged’: [Non-Hermitian Skin Effect \(NHSE\)](#)!

XIII. NON-BLOCH BAND THEORY

XIV. PROJECT PROPOSAL

Proposal A:

- Extensively study various non-Hermitian SSH models (for both diagonal and off-diagonal non-Hermiticity).
- Add interaction to it (e.g. Hubbard type) and study.



- Extend dimensionality and study.
- Maybe design or perform an experiment.

Proposal B:

- Extend the 2-site non-Hermitian problem to a lattice model.
- Employ sophisticated methods such as DMRG, MPS.

REFERENCES

* sag123nik456@gmail.com

† bihan.bhowmick@students.iiserpune.ac.in

‡ reducedpc@gmail.com

¹ W. P. Su, J. R. Schrieffer, and A. J. Heeger, *Phys. Rev. Lett.* **42**, 1698 (1979).

² Courtesy: Asbóth *et al.*, [arXiv:1509.02295v1](https://arxiv.org/abs/1509.02295).