Table 1: Comparison of three compensation strategies across five cancers under four modality-availability settings for the v1 prototype retrieval (protocol `patient-fixed-padmask-v2`). Entries are test C-indices averaged over five seeds; *Overall* is the equally weighted mean of the four settings of a cohort. Proto. v1 denotes the unconditional Top-1 patient retrieval of v1.

	BRCA			UCEC			LUAD			BLCA			LGG		
Scenario	No comp.	Mean	Proto. v1	No comp.	Mean	Proto. v1	No comp.	Mean	Proto. v1	No comp.	Mean	Proto. v1	No comp.	Mean	Proto. v1
RNA masked	0.683842	0.684751	**0.685886**	0.647229	**0.655464**	0.650612	0.580660	0.580239	**0.581011**	**0.593546**	0.589105	0.589938	0.767219	**0.790959**	0.782195
Text masked	**0.655002**	0.619121	0.628545	0.589272	0.577743	**0.597596**	**0.596630**	0.595156	0.576027	**0.575989**	0.574913	0.550069	**0.705866**	0.664113	0.651553
Both masked	0.596026	0.614750	**0.624276**	0.576319	0.575072	**0.596751**	0.585469	**0.588803**	0.573535	0.574046	**0.574774**	0.553088	0.576190	**0.635266**	0.630159
None	**0.691359**	0.686068	0.687930	0.651502	**0.656888**	0.650879	0.582064	**0.583819**	0.583257	**0.589972**	**0.589972**	**0.589972**	0.796825	**0.799517**	0.795997
Overall	0.656557	0.651172	**0.656659**	0.616081	0.616292	**0.623959**	0.586206	**0.587004**	0.578457	**0.583388**	0.582191	0.570767	0.711525	**0.722464**	0.714976

Scenarios: RNA masked, Text masked and Both masked remove the RNA profile, the pathology report or both for all test patients (`rna_100`, `text_100`, `both_100`); None (`none`) adds no artificial mask but retains naturally occurring missingness and is therefore not a complete-data condition. Overall is an average and not a fifth experimental scenario. Strategies: No comp., no compensation (m0real), missing modalities are masked; Mean, mean imputation (m1), a missing modality is replaced by the element-wise training mean of the same cohort; Proto. v1, the v1 retrieval (`retrieval`): the training patient with the highest cosine similarity between row-order-flattened, centred WSI features over commonly valid rows is selected (Top-1), and its raw RNA/Text features replace the missing ones with full validity weight, without routing, shrinkage or down-weighting. Bold marks the highest unrounded value among the three strategies within a cohort and row; ties are all bold. The three strategies use the same 25 frozen networks; no model was retrained and no statistical test was performed. Macro-average over five cohorts, five seeds and four settings: Proto. v1 0.628964, Mean 0.631825, No comp. 0.630751; Proto. v1 was higher than both references in 2 of five cohorts (BRCA and UCEC).