A Coarse Introduction to Algebraic Geometry

A mini talk given on discord by .hchan

Sept 2026

Table of Contents

1 Quick Review

2 Rings of Functions

3 Building the Dictionary

4 The Pièce de Résistance

Quick Review

Rings

Definition

A *ring* is a set R with two binary operations $+$ and $\cdot$ such that

- $(R, +)$ is an abelian group with identity 0,
- $(R, \cdot)$ is associative and commutative, with identity 1,
- $a(b + c) = ab + ac$ for all $a, b, c \in R$.

A *ring homomorphism* $\varphi : R \to S$ preserves $+$, $\cdot$ and 1.

Definition

An *ideal* of R is an additive subgroup $I \subseteq R$ such that $rI \subseteq I$ for all $r \in R$. It is *maximal* if $I \neq R$ and no ideal lies strictly between I and R.

Rings (cont.)

Definition

If $I \subseteq R$ is an ideal, the *quotient ring* R/I is the set of cosets $a + I$, with

$$(a + I) + (b + I) = (a + b) + I, \qquad (a + I)(b + I) = ab + I.$$

These operations are well defined precisely because I is an ideal.

Theorem (First Isomorphism Theorem)

If $\varphi : R \to S$ is a ring homomorphism, then $\ker \varphi$ is an ideal of R, and

$$R/\ker \varphi \cong \operatorname{im} \varphi.$$

Topologies

Definition

Let X be a set. A *topology* $\mathcal{T}$ on X is a collection of subsets of X, called the *open sets*, such that the following three properties are satisfied:

1. $\varnothing, X \in \mathcal{T}$.
2. If $U_1, \ldots, U_n \in \mathcal{T}$ then so is $\bigcap_{i=1}^{n} U_i$; i.e. ($\mathcal{T}$ is closed under *finite intersection*).
3. If $\{U_\alpha\}_{\alpha \in \Lambda} \subseteq \mathcal{T}$ then so is $\bigcup_{\alpha \in \Lambda} U_\alpha$; i.e. ($\mathcal{T}$ is closed under *arbitrary union*).

For simplicity we will often refer to X itself as a topological space.

Rings of Functions

Rings of functions on a space

Suppose that X is a set and R is any ring.

Then the set of all functions from X to R naturally inherits the structure of a ring, via $(f+g)(x) := f(x)+g(x)$ and $(f \cdot g)(x) = f(x) \cdot g(x)$.
The additive identity is the zero function and the multiplicative identity is the one function.

We may restrict this ring to only contain functions with specific properties, e.g. continuous, differentiable, when it makes sense.
We will call these rings *rings of functions*.

Rings of functions on a space (cont.)

Often times we are interested in the cases where the functions have some additional structure. Indeed, it turns out that many areas of mathematics can be described as the study of certain rings of functions:

- In point-set topology, we are interested in studying the ring of continuous functions on a topological space.
- In measure theory, we are interested in studying the ring of bounded measurable functions on $\mathbb{R}^n$.
- In differential geometry, we are interested in studying the ring of smooth (i.e. infinitely differentiable) functions on smooth manifolds.
- In complex analysis, we are interested in studying the ring of holomorphic (complex-differentiable) functions on $\mathbb{C}$.
- In algebraic geometry, we are interested in studying the ring of polynomial functions on ??? (We will answer what this is later).

A first example

Example

Let us consider the ring of continuous real functions on $[0,1]$; i.e. $R = C([0,1])$.

If $p \in [0,1]$, let $\mathfrak{m}_p := \{f \in R : f(p) = 0\}$. It is easily verified that this is an ideal, and in fact a maximal ideal. This is known as the maximal ideal at p.

Theorem

Every maximal ideal of R is given by the maximal ideal at p for some $p \in [0,1]$.

A first example (cont.)

Proof.

Let $\mathfrak{m}$ be a maximal ideal of R. We proceed in 3 steps.

1. If $f \in \mathfrak{m}$, then there must exist some $x \in [0,1]$ such that $f(x) = 0$.
   Indeed, if this is not true then $1/f$ is a continuous function on $[0,1]$, such that f is a unit. But then no maximal ideal can contain a unit.
2. If $f, g \in \mathfrak{m}$, then $Z(f) \cap Z(g) \neq \varnothing$.
   Indeed, if this is not true then $f^2 + g^2$ is nonzero everywhere and is a unit, which is not possible.

A first example (cont.)

Proof.

Let $\mathfrak{m}$ be a maximal ideal of R. We proceed in 3 steps.

1. If $f \in \mathfrak{m}$, then there must exist some $x \in [0,1]$ such that $f(x) = 0$.
   Indeed, if this is not true then $1/f$ is a continuous function on $[0,1]$, such that f is a unit. But then no maximal ideal can contain a unit.
2. If $f, g \in \mathfrak{m}$, then $Z(f) \cap Z(g) \neq \varnothing$.
   Indeed, if this is not true then $f^2 + g^2$ is nonzero everywhere and is a unit, which is not possible.

A first example (cont.)

Proof.

Let $\mathfrak{m}$ be a maximal ideal of R. We proceed in 3 steps.

1. If $f \in \mathfrak{m}$, then there must exist some $x \in [0,1]$ such that $f(x) = 0$.
   Indeed, if this is not true then $1/f$ is a continuous function on $[0,1]$, such that f is a unit. But then no maximal ideal can contain a unit.
2. If $f, g \in \mathfrak{m}$, then $Z(f) \cap Z(g) \neq \varnothing$.
   Indeed, if this is not true then $f^2 + g^2$ is nonzero everywhere and is a unit, which is not possible.

A first example (cont..)

Proof (cont.)

3. There must exist $p \in [0,1]$ such that $f(p) = 0$ for all $f \in \mathfrak{m}$.
   Indeed, if this is not true then for every x we may find an f_x such that $f_x(x) \neq 0$. By continuity we may find an open neighbourhood U_x of x on which f_x is nonzero.
   Proceeding in this way we may find an open cover of $[0,1]$, and by compactness we obtain a finite subcover $\{U_i\}$.
   But then the element $\sum f_i^2 \in \mathfrak{m}$ is a unit, and we have reached a contradiction.

□

A Remark.

The theorem is not true if you replace $[0, 1]$ with $(0, 1)$.
Indeed, let $I = \{f \in R : f \equiv 0 \text{ on some neighbourhood of } 0\}$. This is a proper ideal that cannot be contained in $\mathfrak{m}_p$ for any p.

- This tells us that not all topological spaces and not all rings of functions are well-suited for the algebro-geometric treatment. For those in the know, this is the motivation for the Zariski topology.

Building the Dictionary

Quotients

Let us further build the dictionary between topological spaces and rings, and see what kind of topological properties or operations can be detected by its ring of functions.

Let R_X be a ring of functions on some set X. If $S \subsetneq X$ is some nonempty subset, we may naturally obtain a ring of functions R_S on S by restricting the domain of the functions in R_X; whence a ring homomorphism $\pi : R_X \to R_S$ given by $f \mapsto f|_S$.

Theorem

R_S is isomorphic to the quotient of R_X by the ideal $V(S) := \{f \in R_X : f|_S \equiv 0\}$.

Proof.

π is surjective by definition.
Now suppose that $f \in \ker \pi$. Then $f|_S = \pi(f) = 0$. Conversely if $f|_S \equiv 0$ then $\pi(f) = 0$. □

Quotients

Let us further build the dictionary between topological spaces and rings, and see what kind of topological properties or operations can be detected by its ring of functions.

Let R_X be a ring of functions on some set X. If $S \subsetneq X$ is some nonempty subset, we may naturally obtain a ring of functions R_S on S by restricting the domain of the functions in R_X; whence a ring homomorphism $\pi : R_X \to R_S$ given by $f \mapsto f|_S$.

Theorem

R_S is isomorphic to the quotient of R_X by the ideal $V(S) := \{f \in R_X : f|_S \equiv 0\}$.

Proof.

π is surjective by definition.
Now suppose that $f \in \ker \pi$. Then $f|_S = \pi(f) = 0$. Conversely if $f|_S \equiv 0$ then $\pi(f) = 0$. □

Quotients

Let us further build the dictionary between topological spaces and rings, and see what kind of topological properties or operations can be detected by its ring of functions.

Let R_X be a ring of functions on some set X. If $S \subsetneq X$ is some nonempty subset, we may naturally obtain a ring of functions R_S on S by restricting the domain of the functions in R_X; whence a ring homomorphism $\pi : R_X \to R_S$ given by $f \mapsto f|_S$.

Theorem

R_S is isomorphic to the quotient of R_X by the ideal $V(S) := \{f \in R_X : f|_S \equiv 0\}$.

Proof.

π is surjective by definition.
Now suppose that $f \in \ker \pi$. Then $f|_S = \pi(f) = 0$. Conversely if $f|_S \equiv 0$ then $\pi(f) = 0$. □

Connectedness

If X is a topological space, we say that a subset of X is *disconnected* if it can be written as the union of two disjoint nonempty open sets A, B. In which case we may write $X = A \sqcup B$. The subset is said to be connected otherwise.
Let R_X denote the ring of continuous functions from X to $\mathbb{R}$.

Theorem

$X = A \sqcup B$ if and only if $R_X \simeq R_A \times R_B$.

Proof.

$(\Rightarrow)$: Suppose that $X = A \sqcup B$. Let f be the function that is 1 on A and 0 on B. Then $R_X \simeq \langle f \rangle \times \langle 1 - f \rangle$.
$(\Leftarrow)$: Suppose that $R_X \simeq R_A \times R_B$. Let f be the function that corresponds to $(1, 0)$ under this isomorphism. Then $A = f^{-1}(1)$ and $B = f^{-1}(0)$. □

Connectedness

If X is a topological space, we say that a subset of X is *disconnected* if it can be written as the union of two disjoint nonempty open sets A, B. In which case we may write $X = A \sqcup B$. The subset is said to be connected otherwise.
Let R_X denote the ring of continuous functions from X to $\mathbb{R}$.

Theorem

$X = A \sqcup B$ if and only if $R_X \simeq R_A \times R_B$.

Proof.

$(\Rightarrow)$: Suppose that $X = A \sqcup B$. Let f be the function that is 1 on A and 0 on B. Then $R_X \simeq \langle f \rangle \times \langle 1 - f \rangle$.
$(\Leftarrow)$: Suppose that $R_X \simeq R_A \times R_B$. Let f be the function that corresponds to $(1, 0)$ under this isomorphism. Then $A = f^{-1}(1)$ and $B = f^{-1}(0)$. □

Connectedness

If X is a topological space, we say that a subset of X is *disconnected* if it can be written as the union of two disjoint nonempty open sets A, B. In which case we may write $X = A \sqcup B$. The subset is said to be connected otherwise.
Let R_X denote the ring of continuous functions from X to $\mathbb{R}$.

Theorem

$X = A \sqcup B$ *if and only if* $R_X \simeq R_A \times R_B$.

Proof.

$(\Rightarrow)$: Suppose that $X = A \sqcup B$. Let f be the function that is 1 on A and 0 on B. Then $R_X \simeq \langle f \rangle \times \langle 1 - f \rangle$.
$(\Leftarrow)$: Suppose that $R_X \simeq R_A \times R_B$. Let f be the function that corresponds to $(1, 0)$ under this isomorphism. Then $A = f^{-1}(1)$ and $B = f^{-1}(0)$. □

Irreducibility

If X is a topological space, we say that a subset of X is *reducible* if it can be written as the union of two distinct nonempty proper closed sets C, D. It is said to be irreducible otherwise. Let R_X denote the ring of continuous functions from X to $\mathbb{R}$.

Theorem

If R_X has zero divisors, then X is reducible.

Proof.

Suppose that $fg = 0$ with $f, g \neq 0$. Then $X = f^{-1}(0) \cup g^{-1}(0)$. Both sets are closed, and they are proper as neither f nor g is 0 by assumption. They are distinct, as otherwise $X = f^{-1}(0) \cup g^{-1}(0) = f^{-1}(0)$, which forces $f = 0$. □

Irreducibility

If X is a topological space, we say that a subset of X is *reducible* if it can be written as the union of two distinct nonempty proper closed sets C, D. It is said to be irreducible otherwise. Let R_X denote the ring of continuous functions from X to $\mathbb{R}$.

Theorem

If R_X has zero divisors, then X is reducible.

Proof.

Suppose that $fg = 0$ with $f, g \neq 0$. Then $X = f^{-1}(0) \cup g^{-1}(0)$. Both sets are closed, and they are proper as neither f nor g is 0 by assumption. They are distinct, as otherwise $X = f^{-1}(0) \cup g^{-1}(0) = f^{-1}(0)$, which forces $f = 0$. □

Irreducibility

If X is a topological space, we say that a subset of X is *reducible* if it can be written as the union of two distinct nonempty proper closed sets C, D. It is said to be irreducible otherwise. Let R_X denote the ring of continuous functions from X to $\mathbb{R}$.

Theorem

If R_X has zero divisors, then X is reducible.

Proof.

Suppose that $fg = 0$ with $f, g \neq 0$. Then $X = f^{-1}(0) \cup g^{-1}(0)$. Both sets are closed, and they are proper as neither f nor g is 0 by assumption. They are distinct, as otherwise $X = f^{-1}(0) \cup g^{-1}(0) = f^{-1}(0)$, which forces $f = 0$. □

Closed Sets

Let X be a compact Hausdorff space. For a closed set $C \subseteq X$ and some subset $S \subseteq R_X$, define

$$V(C) := \{f \in R_X : f|_C = 0\}, \qquad Z(S) := \{x \in X : f(x) = 0 \text{ for all } f \in S\}.$$

Theorem

If C is closed, then $Z(V(C)) = C$.

Proof.

Every $f \in V(C)$ vanishes on C, so $C \subseteq Z(V(C))$.
Conversely, if $x \notin C$, by Urysohn's lemma there exists f with $f|_C = 0$ and $f(x) = 1$. □

Closed Sets

Let X be a compact Hausdorff space. For a closed set $C \subseteq X$ and some subset $S \subseteq R_X$, define

$$V(C) := \{f \in R_X : f|_C = 0\}, \qquad Z(S) := \{x \in X : f(x) = 0 \text{ for all } f \in S\}.$$

Theorem

If C is closed, then $Z(V(C)) = C$.

Proof.

Every $f \in V(C)$ vanishes on C, so $C \subseteq Z(V(C))$.
Conversely, if $x \notin C$, by Urysohn's lemma there exists f with $f|_C = 0$ and $f(x) = 1$. □

Closed Sets

Let X be a compact Hausdorff space. For a closed set $C \subseteq X$ and some subset $S \subseteq R_X$, define

$$V(C) := \{f \in R_X : f|_C = 0\}, \qquad Z(S) := \{x \in X : f(x) = 0 \text{ for all } f \in S\}.$$

Theorem

If C is closed, then $Z(V(C)) = C$.

Proof.

Every $f \in V(C)$ vanishes on C, so $C \subseteq Z(V(C))$.
Conversely, if $x \notin C$, by Urysohn's lemma there exists f with $f|_C = 0$ and $f(x) = 1$. □

Ideals

Let X be a compact Hausdorff space. For a closed set $C \subseteq X$ and some subset $S \subseteq R_X$, define

$$V(C) := \{f \in R_X : f|_C = 0\}, \qquad Z(S) := \{x \in X : f(x) = 0 \text{ for all } f \in S\}.$$

Theorem

$Z(S) = Z(\langle S \rangle)$ for all S.

Proof.

$Z(\langle S \rangle) \subseteq Z(S)$ follows immediately from the definition, since $S \subseteq \langle S \rangle$.
Conversely, suppose that f, g vanish on $Z(S)$. Then $(f+g)|_{Z(S)} = 0 + 0 = 0$.
For any $h \in R_X$, $(h \cdot f)|_{Z(S)} = h|_{Z(S)} \cdot 0 = 0$. Hence every element of $\langle S \rangle$ vanishes on $Z(S)$, i.e. $Z(S) \subseteq Z(\langle S \rangle)$. □

Ideals

Let X be a compact Hausdorff space. For a closed set $C \subseteq X$ and some subset $S \subseteq R_X$, define

$$V(C) := \{f \in R_X : f|_C = 0\}, \qquad Z(S) := \{x \in X : f(x) = 0 \text{ for all } f \in S\}.$$

Theorem

$Z(S) = Z(\langle S \rangle)$ for all S.

Proof.

$Z(\langle S \rangle) \subseteq Z(S)$ follows immediately from the definition, since $S \subseteq \langle S \rangle$.
Conversely, suppose that f, g vanish on $Z(S)$. Then $(f+g)|_{Z(S)} = 0 + 0 = 0$.
For any $h \in R_X$, $(h \cdot f)|_{Z(S)} = h|_{Z(S)} \cdot 0 = 0$. Hence every element of $\langle S \rangle$ vanishes on $Z(S)$, i.e. $Z(S) \subseteq Z(\langle S \rangle)$. □

Ideals

Let X be a compact Hausdorff space. For a closed set $C \subseteq X$ and some subset $S \subseteq R_X$, define

$$V(C) := \{f \in R_X : f|_C = 0\}, \qquad Z(S) := \{x \in X : f(x) = 0 \text{ for all } f \in S\}.$$

Theorem

$Z(S) = Z(\langle S \rangle)$ *for all* S.

Proof.

$Z(\langle S \rangle) \subseteq Z(S)$ follows immediately from the definition, since $S \subseteq \langle S \rangle$.
Conversely, suppose that f, g vanish on $Z(S)$. Then $(f+g)|_{Z(S)} = 0 + 0 = 0$.
For any $h \in R_X$, $(h \cdot f)|_{Z(S)} = h|_{Z(S)} \cdot 0 = 0$. Hence every element of $\langle S \rangle$ vanishes on $Z(S)$, i.e. $Z(S) \subseteq Z(\langle S \rangle)$. □

Closed set - ideal correspondence

Theorem

$C \mapsto V(C)$ defines an injective, inclusion-reversing map from closed sets to ideals.

Hence proving facts about the ideals of R will prove facts about the closed sets of X. This map is almost never surjective, however. For instance, when $X = [0, 1]$, the principal ideals generated by x and x^2 are distinct (as $x/x^2 = 1/x$ is not continuous at 0) but share the same zero loci, namely $\{0\}$.

- For those in the know, this is exactly the issue that Hilbert's Nullstellensatz fixes. That theorem is the algebraic analogue of the $C([0, 1])$ example we saw earlier, and how you prove that maximal ideals correspond to (closed) points in your topological space. This is the "correct" geometric interpretation of the Nullstellensatz.

Closed set - ideal correspondence

Theorem

$C \mapsto V(C)$ defines an injective, inclusion-reversing map from closed sets to ideals.

Hence proving facts about the ideals of R will prove facts about the closed sets of X. This map is almost never surjective, however. For instance, when $X = [0, 1]$, the principal ideals generated by x and x^2 are distinct (as $x/x^2 = 1/x$ is not continuous at 0) but share the same zero loci, namely $\{0\}$.

- For those in the know, this is exactly the issue that Hilbert's Nullstellensatz fixes. That theorem is the algebraic analogue of the $C([0, 1])$ example we saw earlier, and how you prove that maximal ideals correspond to (closed) points in your topological space. This is the "correct" geometric interpretation of the Nullstellensatz.

The Pièce de Résistance

Introduction

We have now seen that various topological properties can be detected by ring-theoretic properties of some corresponding ring of functions on that space. But that itself is not that impressive - why not just look at the original space itself? Why bother with rings of functions at all?

As mentioned before, in algebraic geometry it turns out that we are only interested in polynomial (or rational) functions, as those are the only functions that can be well described by abstract algebra.

Hence, for this example, we will take $R = \mathbb{R}[x, y]$, considered as a ring of polynomial functions on the vector space $\mathbb{R}^2$ in the natural way.
Let our topological space X be the graph of the parabola $y = x^2$. By our previous discussion on quotients, we have that $R_X \simeq \mathbb{R}[x, y]/\langle y - x^2 \rangle$. This is the space of all unique real-valued polynomial functions on X.

Introduction

We have now seen that various topological properties can be detected by ring-theoretic properties of some corresponding ring of functions on that space. But that itself is not that impressive - why not just look at the original space itself? Why bother with rings of functions at all?

As mentioned before, in algebraic geometry it turns out that we are only interested in polynomial (or rational) functions, as those are the only functions that can be well described by abstract algebra.

Hence, for this example, we will take $R = \mathbb{R}[x, y]$, considered as a ring of polynomial functions on the vector space $\mathbb{R}^2$ in the natural way.
Let our topological space X be the graph of the parabola $y = x^2$. By our previous discussion on quotients, we have that $R_X \simeq \mathbb{R}[x, y]/\langle y - x^2 \rangle$. This is the space of all unique real-valued polynomial functions on X.


Introduction

We have now seen that various topological properties can be detected by ring-theoretic properties of some corresponding ring of functions on that space. But that itself is not that impressive - why not just look at the original space itself? Why bother with rings of functions at all?

As mentioned before, in algebraic geometry it turns out that we are only interested in polynomial (or rational) functions, as those are the only functions that can be well described by abstract algebra.


Hence, for this example, we will take $R = \mathbb{R}[x, y]$, considered as a ring of polynomial functions on the vector space $\mathbb{R}^2$ in the natural way.
Let our topological space X be the graph of the parabola $y = x^2$. By our previous discussion on quotients, we have that $R_X \simeq \mathbb{R}[x, y]/\langle y - x^2 \rangle$. This is the space of all unique real-valued polynomial functions on X.

A simple question



How many times does the curve intersect the x-axis?

A simple question



How many times does the curve intersect the x-axis?

There is one. This isn't a trick question.

Or is it?



The problem

- The graph has two horizontal intercepts for every positive y-value. In other words, the "limit" of a line with two points on it is a line with a single point on it. That can't be right, 2 shouldn't suddenly become 1.
- For those that know some linear algebra, you might recognize that what is happening is that the geometric multiplicity of the zero of the graph at $(0, 0)$ is equal to 1, but the algebraic multiplicity of the zero of the graph is equal to 2.
- In other words, even though the equation $x^2 = 0$ is saying that there should be two solutions at this point, there is physically not enough room in the graph for those two solutions to have separate points.

The problem

- The graph has two horizontal intercepts for every positive y-value. In other words, the "limit" of a line with two points on it is a line with a single point on it. That can't be right, 2 shouldn't suddenly become 1.
- For those that know some linear algebra, you might recognize that what is happening is that the geometric multiplicity of the zero of the graph at $(0, 0)$ is equal to 1, but the algebraic multiplicity of the zero of the graph is equal to 2.
- In other words, even though the equation $x^2 = 0$ is saying that there should be two solutions at this point, there is physically not enough room in the graph for those two solutions to have separate points.

The problem

- The graph has two horizontal intercepts for every positive y-value. In other words, the "limit" of a line with two points on it is a line with a single point on it. That can't be right, 2 shouldn't suddenly become 1.
- For those that know some linear algebra, you might recognize that what is happening is that the geometric multiplicity of the zero of the graph at $(0, 0)$ is equal to 1, but the algebraic multiplicity of the zero of the graph is equal to 2.
- In other words, even though the equation $x^2 = 0$ is saying that there should be two solutions at this point, there is physically not enough room in the graph for those two solutions to have separate points.

The fix


- Taking the horizontal intercept of the graph with the line $y = c$ is the same as restricting X with the subspace $X \cap \{y = c\}$.
- By the discussion we had before, for the ring of polynomial functions this corresponds to the quotient map $(\mathbb{R}[x, y]/\langle y - x^2 \rangle)/\langle y - c \rangle$.
- When $c > 0$, the ring is isomorphic to $\mathbb{R} \oplus \mathbb{R}$ which has two nonzero proper ideals, which tells you that there are exactly two closed sets in the intersection as we expect.

The fix

- Taking the horizontal intercept of the graph with the line $y = c$ is the same as restricting X with the subspace $X \cap \{y = c\}$.
- By the discussion we had before, for the ring of polynomial functions this corresponds to the quotient map $(\mathbb{R}[x, y]/\langle y - x^2 \rangle)/\langle y - c \rangle$.
- When $c > 0$, the ring is isomorphic to $\mathbb{R} \oplus \mathbb{R}$ which has two nonzero proper ideals, which tells you that there are exactly two closed sets in the intersection as we expect.

The fix

- Taking the horizontal intercept of the graph with the line $y = c$ is the same as restricting X with the subspace $X \cap \{y = c\}$.
- By the discussion we had before, for the ring of polynomial functions this corresponds to the quotient map $(\mathbb{R}[x, y]/\langle y - x^2 \rangle)/\langle y - c \rangle$.
- When $c > 0$, the ring is isomorphic to $\mathbb{R} \oplus \mathbb{R}$ which has two nonzero proper ideals, which tells you that there are exactly two closed sets in the intersection as we expect.



The fix (cont.)

What happens now when $c = 0$?

- By the third isomorphism theorem, the ring of polynomial functions is now isomorphic to $\mathbb{R}[x]/\langle x^2 \rangle$.
- This has one single nonzero proper ideal, which tells us there is one point of intersection.
- But $\mathbb{R}[x]/\langle x^2 \rangle$ is a dimension *two* vector space over the real numbers, so the ring has *remembered the fact that this point is supposed to have a "weight" of two.*
- In other words, the ring of functions has managed to keep track of information about the space X that X itself is unable to remember. It somehow knows more about X than X knows about itself!

The fix (cont.)

What happens now when $c = 0$?

- By the third isomorphism theorem, the ring of polynomial functions is now isomorphic to $\mathbb{R}[x]/\langle x^2 \rangle$.
- This has one single nonzero proper ideal, which tells us there is one point of intersection.
- But $\mathbb{R}[x]/\langle x^2 \rangle$ is a dimension *two* vector space over the real numbers, so the ring has *remembered the fact that this point is supposed to have a "weight" of two.*
- In other words, the ring of functions has managed to keep track of information about the space X that X itself is unable to remember. It somehow knows more about X than X knows about itself!

The fix (cont.)


What happens now when $c = 0$?

- By the third isomorphism theorem, the ring of polynomial functions is now isomorphic to $\mathbb{R}[x]/\langle x^2 \rangle$.
- This has one single nonzero proper ideal, which tells us there is one point of intersection.
- But $\mathbb{R}[x]/\langle x^2 \rangle$ is a dimension *two* vector space over the real numbers, so the ring has *remembered the fact that this point is supposed to have a "weight" of two*.
- In other words, the ring of functions has managed to keep track of information about the space X that X itself is unable to remember. It somehow knows more about X than X knows about itself!

The fix (cont.)

What happens now when $c = 0$?

- By the third isomorphism theorem, the ring of polynomial functions is now isomorphic to $\mathbb{R}[x]/\langle x^2 \rangle$.
- This has one single nonzero proper ideal, which tells us there is one point of intersection.
- But $\mathbb{R}[x]/\langle x^2 \rangle$ is a dimension *two* vector space over the real numbers, so the ring has *remembered the fact that this point is supposed to have a "weight" of two*.
- In other words, the ring of functions has managed to keep track of information about the space X that X itself is unable to remember. It somehow knows more about X than X knows about itself!



But what does this mean?

As a real vector space, the ring $\mathbb{R}[x]/\langle x^2 \rangle$ has a basis $\{1, \overline{x}\}$. Is there a way to figure out what this means?

- Recall that when $c > 0$, we have that the ring of polynomials is isomorphic to $\mathbb{R}[x]/\langle x - \sqrt{c} \rangle \oplus \mathbb{R}[x]/\langle x + \sqrt{c} \rangle$.
- This tells us that if you give me some unknown polynomial f; then knowing the values $f(\sqrt{c})$ and $f(-\sqrt{c})$ is exactly all the information you need to figure out what f is on the intersection.
- Following this interpretation, this is telling you that even at the single point $(0, 0)$, you still need two pieces of information to be able to figure out what the unknown polynomial was on the intersection:
  - Its actual value at the origin (i.e. $f(0)$); and
  - the *derivative* of the polynomial (i.e. $f'(0)$).

But what does this mean?

As a real vector space, the ring $\mathbb{R}[x]/\langle x^2 \rangle$ has a basis $\{1, \overline{x}\}$. Is there a way to figure out what this means?

- Recall that when $c > 0$, we have that the ring of polynomials is isomorphic to $\mathbb{R}[x]/\langle x - \sqrt{c} \rangle \oplus \mathbb{R}[x]/\langle x + \sqrt{c} \rangle$.
- This tells us that if you give me some unknown polynomial f; then knowing the values $f(\sqrt{c})$ and $f(-\sqrt{c})$ is exactly all the information you need to figure out what f is on the intersection.
- Following this interpretation, this is telling you that even at the single point $(0, 0)$, you still need two pieces of information to be able to figure out what the unknown polynomial was on the intersection:
  - Its actual value at the origin (i.e. $f(0)$); and
  - the *derivative* of the polynomial (i.e. $f'(0)$).

But what does this mean?

As a real vector space, the ring $\mathbb{R}[x]/\langle x^2 \rangle$ has a basis $\{1, \overline{x}\}$. Is there a way to figure out what this means?

- Recall that when $c > 0$, we have that the ring of polynomials is isomorphic to $\mathbb{R}[x]/\langle x - \sqrt{c} \rangle \oplus \mathbb{R}[x]/\langle x + \sqrt{c} \rangle$.
- This tells us that if you give me some unknown polynomial f; then knowing the values $f(\sqrt{c})$ and $f(-\sqrt{c})$ is exactly all the information you need to figure out what f is on the intersection.
- Following this interpretation, this is telling you that even at the single point $(0, 0)$, you still need two pieces of information to be able to figure out what the unknown polynomial was on the intersection:
  - Its actual value at the origin (i.e. $f(0)$); and
  - the *derivative* of the polynomial (i.e. $f'(0)$).

But wait, there's more!

What happens when $c < 0$?

- The ring of polynomial functions is now isomorphic to $\mathbb{R}[x]/\langle x^2 - c \rangle$, which is isomorphic to $\mathbb{C}$.
- $\mathbb{C}$ has no nonzero proper ideals, so this tells us that there are zero points of intersection.
- But $\mathbb{C}$ still has real dimension 2, so it is saying that there *still* should be two solutions!
- The ring of polynomials has discovered the fact that our original graph was missing **the entire imaginary axis**, and X is missing some solutions.

But wait, there's more!

What happens when $c < 0$?

- The ring of polynomial functions is now isomorphic to $\mathbb{R}[x]/\langle x^2 - c \rangle$, which is isomorphic to $\mathbb{C}$.
- $\mathbb{C}$ has no nonzero proper ideals, so this tells us that there are zero points of intersection.
- But $\mathbb{C}$ still has real dimension 2, so it is saying that there *still* should be two solutions!
- The ring of polynomials has discovered the fact that our original graph was missing **the entire imaginary axis**, and X is missing some solutions.

But wait, there's more!


What happens when $c < 0$?

- The ring of polynomial functions is now isomorphic to $\mathbb{R}[x]/\langle x^2 - c \rangle$, which is isomorphic to $\mathbb{C}$.
- $\mathbb{C}$ has no nonzero proper ideals, so this tells us that there are zero points of intersection.
- But $\mathbb{C}$ still has real dimension 2, so it is saying that there *still* should be two solutions!
- The ring of polynomials has discovered the fact that our original graph was missing **the entire imaginary axis**, and X is missing some solutions.

But wait, there's more!

What happens when $c < 0$?

- The ring of polynomial functions is now isomorphic to $\mathbb{R}[x]/\langle x^2 - c \rangle$, which is isomorphic to $\mathbb{C}$.
- $\mathbb{C}$ has no nonzero proper ideals, so this tells us that there are zero points of intersection.
- But $\mathbb{C}$ still has real dimension 2, so it is saying that there *still* should be two solutions!
- The ring of polynomials has discovered the fact that our original graph was missing **the entire imaginary axis**, and X is missing some solutions.



What is Algebraic Geometry?

- Algebraic geometry is the study of the *zero loci of systems of polynomials* using algebraic structures, mainly groups, rings, fields and modules.
- Modern algebraic geometry is motivated by the philosophy that *every* ring R should be the ring of functions on some topological space (namely Spec R, if you know what that is).
  - The space Spec R is known as a *scheme*, and was one of the twelve "great ideas" that Grothendieck contributed to the world of mathematics.

2.3 PERIODS

I. Prehistory (-400-1630)

II. Exploration (1630-1795)

III. The golden age of projective geometry (1795-1850)

IV. Riemann and birational geometry (1850-1866)

V. Development and chaos (1866-1920)

a) Algebraic approach
b) Brill-Noether
c) Integrals on higher dimensional varieties
d) The Italian school

VI. New structures in algebraic geometry (1920-1950)

a) Kählerian manifolds and return to Riemann
b) Abstract algebraic geometry

VII. Sheaves and schemes (1950-)

a) The Riemann Roch theorem and sheaf cohomology
b) Serre [...] varieties
c) Schemes and topologies

[1] Image source: https://arxiv.org/pdf/1803.00163

What is Algebraic Geometry?

- Algebraic geometry is the study of the *zero loci of systems of polynomials* using algebraic structures, mainly groups, rings, fields and modules.
- Modern algebraic geometry is motivated by the philosophy that *every* ring R should be the ring of functions on some topological space (namely $\operatorname{Spec} R$, if you know what that is).
  - The space $\operatorname{Spec} R$ is known as a *scheme*, and was one of the twelve "great ideas" that Grothendieck contributed to the world of mathematics.

2.3 PERIODS

I. Prehistory (-400-1630)

II. Exploration (1630-1795)

III. The golden age of projective geometry (1795-1850)

IV. Riemann and birational geometry (1850-1866)

V. Development and chaos (1866-1920)

a) Algebraic approach
b) Brill-Noether
c) Integrals on higher dimensional varieties
d) The Italian school

VI. New structures in algebraic geometry (1920-1950)

a) Kählerian manifolds and return to Riemann
b) Abstract algebraic geometry

VII. Sheaves and schemes (1950-)

a) The Riemann Roch theorem and sheaf cohomology
b) Serre [...] varieties
c) Schemes and topologies

[1]Image source: `https://arxiv.org/pdf/1803.00163`

Any Questions?




Figure 7. An algebraic family of schemes.