

New Track Fitting

Hsin-Yi Chou^a, Yuan-Hann Chang^a, Yi Yang^b, Han-Sheng Li^b, Sadakazu Haino^c

^a*National Central University, Chung-Li, Tao Yuan 32054, Taiwan*

^b*National Cheng Kung University, Tainan 70101, Taiwan*

^c*Institute of Physics, Academia Sinica, Nankang, Taipei 11529, Taiwan*

Abstract

This template helps you to create a properly formatted L^AT_EX manuscript.

Keywords: AMS-02, Track Fitting

1. Equation of Motion

Notation.

<i>step length</i> : S	$[cm]$
<i>path length</i> : s	$[cm]$
<i>position</i> : \vec{p}	$[cm]$
<i>unit direction</i> : \hat{u}	$[1]$
<i>orthogonal direction</i> : \hat{n}_τ and \hat{n}_ρ	$[1]$
<i>eta</i> : $\tilde{\eta} = \tilde{q}/(\gamma\beta)$	$[1]$
<i>rigidity</i> : $R = (m/q) \cdot \gamma\beta$	$[GV]$
<i>mass</i> : m	$[GeV]$
<i>charge</i> : q	$[e]$
<i>charge sign</i> : $\tilde{q} = q/ q $	$[1]$
<i>electron mass</i> : $m_e = 0.510999$	$[MeV]$
<i>electron charge</i> : $q_e = 1$	$[e]$
<i>velocity</i> : β	$[1]$
<i>Lorentz factor</i> : $\gamma = 1/\sqrt{1 - \beta^2}$	$[1]$
<i>magnetic field</i> : \vec{B}	$[kGauss]$
<i>lambda</i> : $\lambda = 2.99792458 \cdot 10^{-4}$	$[GV \cdot kGauss^{-1} \cdot cm^{-1}]$
<i>radiation length</i> : X_0	$[g \cdot cm^{-2}]$
<i>mass density</i> : D	$[g \cdot cm^{-3}]$
<i>cross length</i> : $L = D/X_0$	$[cm^{-1}]$
<i>passage length</i> : l	$[cm]$
<i>Rydberg constant</i> : $C_{\text{Rydberg}} = 0.0136$	$[GeV]$
<i>K</i> : $K = 0.307075$	$[MeV \cdot mol^{-1} \cdot cm^2]$
<i>atomic mass</i> : A	$[g \cdot mol^{-1}]$
<i>atomic number</i> : Z	$[1]$
<i>density effect</i> : $\delta(\gamma\beta)$	$[MeV]$
<i>mean excitation energy</i> : I	$[eV]$
<i>Landau factor</i> : $j = 0.200$	$[1]$

Lorentz Force.

$$\begin{aligned}
\frac{d\hat{u}}{ds} &= \Lambda \cdot (\hat{u} \times \vec{B}) \cdot \tilde{\eta} \\
\Lambda &= \lambda \cdot |q/m| \\
(\hat{u} \times \vec{B}) &= \begin{pmatrix} \hat{u}_y \vec{B}_z - \hat{u}_z \vec{B}_y \\ \hat{u}_z \vec{B}_x - \hat{u}_x \vec{B}_z \\ \hat{u}_x \vec{B}_y - \hat{u}_y \vec{B}_x \end{pmatrix}
\end{aligned} \tag{2}$$

Multiple Scattering.

$$\begin{aligned}
\frac{d\hat{u}}{ds} &= (\omega_\tau \cdot \hat{n}_\tau + \omega_\rho \cdot \hat{n}_\rho) \cdot \Omega_\theta \\
\hat{n}_\tau &= (\vec{B} - (\hat{u} \cdot \vec{B}) \cdot \hat{u}) / |\hat{u} \times \vec{B}| \\
\hat{n}_\rho &= (\hat{u} \times \vec{B}) / |\hat{u} \times \vec{B}| \\
\Omega_\theta &= C_{\text{Rydberg}} \cdot |q/m| \cdot (|\tilde{\eta}| \sqrt{\tilde{\eta}^2 + 1}) \cdot \sqrt{l \cdot L} \cdot (1 + 0.038 \cdot \ln(l \cdot L)) \cdot (1/S) \\
\omega_\tau, \omega_\rho &\sim \text{Normal}(0, 1) \\
f(\omega) &= \frac{1}{\sqrt{2\pi}} \cdot \exp\left(-\frac{\omega^2}{2}\right)
\end{aligned} \tag{3}$$

Ionisation Energy Loss.

$$\begin{aligned}
\frac{d\tilde{\eta}}{ds} &= \tilde{\eta} \cdot (\omega_I \cdot \Omega_\delta + \Omega_\Delta) \\
\Omega_I &= (K/2) \cdot (D \cdot l \cdot Z/A) \cdot (q^2) \cdot (\tilde{\eta}^2 + 1) \\
\Omega_\delta &= \Omega_I \cdot (1/m) \cdot (\tilde{\eta}^2 \sqrt{\tilde{\eta}^{-2} + 1}) \cdot (1/S) \\
\Omega_\Delta &= \Omega_\delta \cdot \left[\ln\left(\frac{2m_e \cdot \tilde{\eta}^{-2}}{I}\right) + \ln\left(\frac{\Omega_I}{I}\right) + j - \beta^2 - \delta(\gamma\beta) \right] \\
\omega_I &\sim \text{Landau}(0, 1) \text{ require } [\omega_I \geq (-\Omega_\Delta/\Omega_\delta)] \\
f(\omega_I) &\simeq \frac{1}{\sqrt{2\pi}} \cdot \exp\left(-\frac{1}{2}(\omega_I + \exp(-\omega_I))\right)
\end{aligned} \tag{4}$$

Bremsstrahlung Energy Loss.

$$\begin{aligned}
\frac{d\tilde{\eta}}{ds} &= \tilde{\eta} \cdot (\omega_B \cdot \Omega_B) \\
\Omega_B &= \left(\frac{q}{q_e} \cdot \frac{m_e}{m} \right)^2 \cdot (\sqrt{\tilde{\eta}^{-2} + 1} - 1) \cdot \left(\frac{l \cdot L}{\ln 2} \right) \cdot (\tilde{\eta}^2 \sqrt{\tilde{\eta}^{-2} + 1}) \cdot (1/S) \\
\omega_B &\sim \Gamma(t, t) \text{ require } [0 \leq \omega_B < \inf] \\
f(\omega_B) &= \left(\frac{t^t}{\Gamma(t)} \right) \cdot (\omega_B^{t-1} \cdot \exp(-\omega_B \cdot t)) \text{ where } t = \left(\frac{l \cdot L}{\ln 2} \right)
\end{aligned} \tag{5}$$

Summary.

$$\begin{aligned}
\frac{d\vec{p}}{ds} &= \hat{u} \\
\frac{d\hat{u}}{ds} &= \Lambda \cdot (\hat{u} \times \vec{B}) \cdot \tilde{\eta} + (\omega_\tau \cdot \hat{n}_\tau + \omega_\rho \cdot \hat{n}_\rho) \cdot \Omega_\theta \\
\frac{d\tilde{\eta}}{ds} &= \tilde{\eta} \cdot [(\omega_I \cdot \Omega_\delta + \Omega_\Delta) + (\omega_B \cdot \Omega_B)]
\end{aligned} \tag{6}$$

2. Propagation

Status.

$$\begin{aligned}
\xi_{\vec{p}}^T &= \begin{pmatrix} x & y & z \end{pmatrix} \\
\xi_{\hat{u}}^T &= \begin{pmatrix} \hat{u}_x & \hat{u}_y & \hat{u}_z \end{pmatrix} \\
\xi_{\tilde{\eta}}^T &= \begin{pmatrix} \tilde{\eta} \end{pmatrix} \\
\xi_{\omega}^T &= \begin{pmatrix} \omega_\tau & \omega_\rho & \omega_I & \omega_B \end{pmatrix} \\
\xi^T &= \begin{pmatrix} \xi_{\vec{p}}^T & \xi_{\hat{u}}^T & \xi_{\tilde{\eta}}^T & \xi_{\omega}^T \end{pmatrix} \\
\xi_G^T &= \begin{pmatrix} \xi_{\vec{p}}^T & \xi_{\hat{u}}^T & \xi_{\tilde{\eta}}^T \end{pmatrix} \\
\xi_L^T &= \begin{pmatrix} \xi_{\omega}^T \end{pmatrix}
\end{aligned} \tag{7}$$

Differential Status.

$$\begin{aligned}
\zeta_{\vec{p}} = \frac{d\xi_{\vec{p}}}{ds} &= \hat{u} \\
\zeta_{\hat{u}} = \frac{d\xi_{\hat{u}}}{ds} &= \Lambda \cdot (\hat{u} \times \vec{B}) \cdot \tilde{\eta} + (\omega_\tau \cdot \hat{n}_\tau + \omega_\rho \cdot \hat{n}_\rho) \cdot \Omega_\theta \\
\zeta_{\tilde{\eta}} = \frac{d\xi_{\tilde{\eta}}}{ds} &= \tilde{\eta} \cdot [(\omega_I \cdot \Omega_\delta + \Omega_\Delta) + (\omega_B \cdot \Omega_B)]
\end{aligned} \tag{8}$$

Transfer Status A.

$$\begin{aligned}
\kappa_{\vec{p}} = \frac{d\zeta_{\vec{p}}}{d\xi} &= \begin{pmatrix} \frac{d\zeta_{\vec{p}}}{d\xi_{\vec{p}}} & \frac{d\zeta_{\vec{p}}}{d\xi_{\hat{u}}} & \frac{d\zeta_{\vec{p}}}{d\xi_{\tilde{\eta}}} & \frac{d\zeta_{\vec{p}}}{d\xi_{\omega}} \end{pmatrix} \\
\kappa_{\hat{u}} = \frac{d\zeta_{\hat{u}}}{d\xi} &= \begin{pmatrix} \frac{d\zeta_{\hat{u}}}{d\xi_{\vec{p}}} & \frac{d\zeta_{\hat{u}}}{d\xi_{\hat{u}}} & \frac{d\zeta_{\hat{u}}}{d\xi_{\tilde{\eta}}} & \frac{d\zeta_{\hat{u}}}{d\xi_{\omega}} \end{pmatrix} \\
\kappa_{\tilde{\eta}} = \frac{d\zeta_{\tilde{\eta}}}{d\xi} &= \begin{pmatrix} \frac{d\zeta_{\tilde{\eta}}}{d\xi_{\vec{p}}} & \frac{d\zeta_{\tilde{\eta}}}{d\xi_{\hat{u}}} & \frac{d\zeta_{\tilde{\eta}}}{d\xi_{\tilde{\eta}}} & \frac{d\zeta_{\tilde{\eta}}}{d\xi_{\omega}} \end{pmatrix} \\
\kappa_{\vec{p}} = \frac{d\zeta_{\vec{p}}}{d\xi} &= \begin{pmatrix} 0 & \frac{d\zeta_{\vec{p}}}{d\xi_{\hat{u}}} & 0 & 0 \end{pmatrix} \\
\kappa_{\hat{u}} = \frac{d\zeta_{\hat{u}}}{d\xi} &= \begin{pmatrix} 0 & \frac{d\zeta_{\hat{u}}}{d\xi_{\hat{u}}} & \frac{d\zeta_{\hat{u}}}{d\xi_{\tilde{\eta}}} & \frac{d\zeta_{\hat{u}}}{d\xi_{\omega}} \end{pmatrix} \\
\kappa_{\tilde{\eta}} = \frac{d\zeta_{\tilde{\eta}}}{d\xi} &= \begin{pmatrix} 0 & 0 & \frac{d\zeta_{\tilde{\eta}}}{d\xi_{\tilde{\eta}}} & \frac{d\zeta_{\tilde{\eta}}}{d\xi_{\omega}} \end{pmatrix}
\end{aligned} \tag{9}$$

Transfer Status B.

$$\begin{aligned}
\frac{d\zeta_{\vec{p}_i}}{d\xi_{\hat{u}_j}} &= \delta_{ij} \\
\frac{d\zeta_{\hat{u}_i}}{d\xi_{\hat{u}_j}} &= \Lambda \cdot \frac{\partial(\hat{u} \times \vec{B})_i}{\partial \hat{u}_j} \cdot \tilde{\eta} \\
\frac{d\zeta_{\hat{u}_i}}{d\xi_{\tilde{\eta}}} &= \Lambda \cdot (\hat{u} \times \vec{B})_i \\
\frac{d\zeta_{\hat{u}_i}}{d\xi_{\omega_\tau}} &= (\Omega_\theta \cdot \hat{n}_\tau)_i \\
\frac{d\zeta_{\hat{u}_i}}{d\xi_{\omega_\rho}} &= (\Omega_\theta \cdot \hat{n}_\rho)_i \\
\frac{d\zeta_{\tilde{\eta}}}{d\xi_{\tilde{\eta}}} &= (\omega_I \cdot \Omega_\delta + \Omega_\Delta) + (\omega_B \cdot \Omega_B) \\
\frac{d\zeta_{\tilde{\eta}}}{d\xi_{\omega_I}} &= \tilde{\eta} \cdot \Omega_\delta \\
\frac{d\zeta_{\tilde{\eta}}}{d\xi_{\omega_B}} &= \tilde{\eta} \cdot \Omega_B \\
\frac{\partial(\hat{u} \times \vec{B})}{\partial \hat{u}} &= \begin{pmatrix} 0 & \vec{B}_z & -\vec{B}_y \\ -\vec{B}_z & 0 & \vec{B}_x \\ \vec{B}_y & -\vec{B}_x & 0 \end{pmatrix}
\end{aligned} \tag{10}$$

Euler Method.

$$\begin{aligned}
\xi^0 &= \xi(0) \\
\xi^f &= \xi(S) \\
\xi_G^f &= \xi_G^0 + \begin{pmatrix} S \cdot \zeta_p^0 + \frac{S^2}{2} \cdot \zeta_u^0 \\ S \cdot \zeta_u^0 \\ S \cdot \zeta_{\tilde{\eta}}^0 \end{pmatrix} \\
J^f &= \begin{pmatrix} J_{GG}^f & J_{GL}^f \end{pmatrix} \\
J_{GG}^f &= I + \begin{pmatrix} S \cdot \kappa_{\tilde{p}G}^0 + \frac{S^2}{2} \cdot \kappa_{uG}^0 \\ S \cdot \kappa_{uG}^0 \\ S \cdot \kappa_{\tilde{\eta}G}^0 \end{pmatrix} \\
J_{GL}^f &= \begin{pmatrix} \frac{S^2}{2} \cdot \kappa_{uL}^0 \\ S \cdot \kappa_{uL}^0 \\ S \cdot \kappa_{\tilde{\eta}L}^0 \end{pmatrix}
\end{aligned} \tag{11}$$

Euler-Heun Method.

$$\begin{aligned}
\xi^0 &= \xi(0) \\
\xi^1 &= \xi(S) \\
\xi_G^1 &= \xi_G^0 + \begin{pmatrix} S \cdot \zeta_p^0 + \frac{S^2}{2} \cdot \zeta_u^0 \\ S \cdot \zeta_u^0 \\ S \cdot \zeta_{\eta}^0 \end{pmatrix} \\
J^1 &= \begin{pmatrix} J_{GG}^1 & J_{GL}^1 \end{pmatrix} \\
J_{GG}^1 &= I + \begin{pmatrix} S \cdot \kappa_{pG}^0 + \frac{S^2}{2} \cdot \kappa_{uG}^0 \\ S \cdot \kappa_{uG}^0 \\ S \cdot \kappa_{\eta G}^0 \end{pmatrix} \\
J_{GL}^1 &= \begin{pmatrix} \frac{S^2}{2} \cdot \kappa_{uL}^0 \\ S \cdot \kappa_{uL}^0 \\ S \cdot \kappa_{\eta L}^0 \end{pmatrix} \\
\xi^f &= \xi(S) \\
\xi_G^f &= \xi_G^0 + \begin{pmatrix} S \cdot \zeta_p^0 + \frac{S^2}{6} \cdot (2\zeta_u^0 + \zeta_u^1) \\ \frac{S}{2} \cdot (\zeta_u^0 + \zeta_u^1) \\ \frac{S}{2} \cdot (\zeta_{\eta}^0 + \zeta_{\eta}^1) \end{pmatrix} \\
J^f &= \begin{pmatrix} J_{GG}^f & J_{GL}^f \end{pmatrix} \\
J_{GG}^f &= I + \begin{pmatrix} S \cdot \kappa_{pG}^0 + \frac{S^2}{6} \cdot (2\kappa_{uG}^0 + \kappa_{uG}^1 J_{GG}^1) \\ \frac{S}{2} \cdot (\kappa_{uG}^0 + \kappa_{uG}^1 J_{GG}^1) \\ \frac{S}{2} \cdot (\kappa_{\eta G}^0 + \kappa_{\eta G}^1 J_{GG}^1) \end{pmatrix} \\
J_{GL}^f &= \begin{pmatrix} \frac{S^2}{6} \cdot (2\kappa_{uL}^0 + (\kappa_{uG}^1 J_{GL}^1 + \kappa_{uL}^1)) \\ \frac{S}{2} \cdot (\kappa_{uL}^0 + (\kappa_{uG}^1 J_{GL}^1 + \kappa_{uL}^1)) \\ \frac{S}{2} \cdot (\kappa_{\eta L}^0 + (\kappa_{\eta G}^1 J_{GL}^1 + \kappa_{\eta L}^1)) \end{pmatrix}
\end{aligned} \tag{12}$$

Runge-Kutta-Nystrom Method A.

$$\begin{aligned}
\xi^0 &= \xi(0) \\
\xi^1 &= \xi\left(\frac{S}{2}\right) \\
\xi_G^1 &= \xi_G^0 + \begin{pmatrix} \frac{S}{2} \cdot \zeta_p^0 + \frac{S^2}{8} \cdot \zeta_u^0 \\ \frac{S}{2} \cdot \zeta_u^0 \\ \frac{S}{2} \cdot \zeta_{\tilde{\eta}}^0 \end{pmatrix} \\
J^1 &= \begin{pmatrix} J_{GG}^1 & J_{GL}^1 \end{pmatrix} \\
J_{GG}^1 &= I + \begin{pmatrix} \frac{S}{2} \cdot \kappa_{pG}^0 + \frac{S^2}{8} \cdot \kappa_{uG}^0 \\ \frac{S}{2} \cdot \kappa_{uG}^0 \\ \frac{S}{2} \cdot \kappa_{\tilde{\eta}G}^0 \end{pmatrix} \\
J_{GL}^1 &= \begin{pmatrix} \frac{S^2}{8} \cdot \kappa_{uL}^0 \\ \frac{S}{2} \cdot \kappa_{uL}^0 \\ \frac{S}{2} \cdot \kappa_{\tilde{\eta}L}^0 \end{pmatrix} \\
\xi^2 &= \xi\left(\frac{S}{2}\right) \\
\xi_G^2 &= \xi_G^1 + \begin{pmatrix} \frac{S}{2} \cdot \zeta_p^0 + \frac{S^2}{8} \cdot \zeta_u^0 \\ \frac{S}{2} \cdot \zeta_u^1 \\ \frac{S}{2} \cdot \zeta_{\tilde{\eta}}^1 \end{pmatrix} \\
J^2 &= \begin{pmatrix} J_{GG}^2 & J_{GL}^2 \end{pmatrix} \\
J_{GG}^2 &= I + \begin{pmatrix} \frac{S}{2} \cdot \kappa_{pG}^0 + \frac{S^2}{8} \cdot \kappa_{uG}^0 \\ \frac{S}{2} \cdot \kappa_{uG}^1 J_{GG}^1 \\ \frac{S}{2} \cdot \kappa_{\tilde{\eta}G}^1 J_{GG}^1 \end{pmatrix} \\
J_{GL}^2 &= \begin{pmatrix} \frac{S^2}{8} \cdot \kappa_{uL}^0 \\ \frac{S}{2} \cdot (\kappa_{uG}^1 J_{GL}^1 + \kappa_{uL}^1) \\ \frac{S}{2} \cdot (\kappa_{\tilde{\eta}G}^1 J_{GL}^1 + \kappa_{\tilde{\eta}L}^1) \end{pmatrix}
\end{aligned} \tag{13}$$

Runge-Kutta-Nystrom Method B.

$$\begin{aligned}
\xi^3 &= \xi(S) \\
\xi_G^3 &= \xi_G^0 + \begin{pmatrix} S \cdot \zeta_p^0 + \frac{S^2}{2} \cdot \zeta_u^2 \\ S \cdot \zeta_u^2 \\ S \cdot \zeta_{\tilde{\eta}}^2 \end{pmatrix} \\
J^3 &= \begin{pmatrix} J_{GG}^3 & J_{GL}^3 \end{pmatrix} \\
J_{GG}^3 &= I + \begin{pmatrix} S \cdot \kappa_{pG}^0 + \frac{S^2}{2} \cdot \kappa_{uG}^2 J_{GG}^2 \\ S \cdot \kappa_{uG}^2 J_{GG}^2 \\ S \cdot \kappa_{\tilde{\eta}G}^2 J_{GG}^2 \end{pmatrix} \\
J_{GL}^3 &= \begin{pmatrix} \frac{S^2}{2} \cdot (\kappa_{uG}^2 J_{GL}^2 + \kappa_{uL}^2) \\ S \cdot (\kappa_{uG}^2 J_{GL}^2 + \kappa_{uL}^2) \\ S \cdot (\kappa_{\tilde{\eta}G}^2 J_{GL}^2 + \kappa_{\tilde{\eta}L}^2) \end{pmatrix} \\
\xi^f &= \xi(S) \\
\xi_G^f &= \xi_G^0 + \begin{pmatrix} S \cdot \zeta_p^0 + \frac{S^2}{6} \cdot (\zeta_u^0 + \zeta_u^1 + \zeta_u^2) \\ \frac{S}{6} \cdot (\zeta_u^0 + 2\zeta_u^1 + 2\zeta_u^2 + \zeta_u^3) \\ \frac{S}{6} \cdot (\zeta_{\tilde{\eta}}^0 + 2\zeta_{\tilde{\eta}}^1 + 2\zeta_{\tilde{\eta}}^2 + \zeta_{\tilde{\eta}}^3) \end{pmatrix} \\
J^f &= \begin{pmatrix} J_{GG}^f & J_{GL}^f \end{pmatrix} \\
J_{GG}^f &= I + \begin{pmatrix} S \cdot \kappa_{pG}^0 + \frac{S^2}{6} \cdot (\kappa_{uG}^0 + \kappa_{uG}^1 J_{GG}^1 + \kappa_{uG}^2 J_{GG}^2) \\ \frac{S}{6} \cdot (\kappa_{uG}^0 + 2\kappa_{uG}^1 J_{GG}^1 + 2\kappa_{uG}^2 J_{GG}^2 + \kappa_{uG}^3 J_{GG}^3) \\ \frac{S}{6} \cdot (\kappa_{\tilde{\eta}G}^0 + 2\kappa_{\tilde{\eta}G}^1 J_{GG}^1 + 2\kappa_{\tilde{\eta}G}^2 J_{GG}^2 + \kappa_{\tilde{\eta}G}^3 J_{GG}^3) \end{pmatrix} \\
J_{GL}^f &= \begin{pmatrix} \frac{S^2}{6} \cdot (\kappa_{uL}^0 + (\kappa_{uG}^1 J_{GL}^1 + \kappa_{uL}^1) + (\kappa_{uG}^2 J_{GL}^2 + \kappa_{uL}^2)) \\ \frac{S}{6} \cdot (\kappa_{uL}^0 + 2(\kappa_{uG}^1 J_{GL}^1 + \kappa_{uL}^1) + 2(\kappa_{uG}^2 J_{GL}^2 + \kappa_{uL}^2) + (\kappa_{uG}^3 J_{GL}^3 + \kappa_{uL}^3)) \\ \frac{S}{6} \cdot (\kappa_{\tilde{\eta}L}^0 + 2(\kappa_{\tilde{\eta}G}^1 J_{GL}^1 + \kappa_{\tilde{\eta}L}^1) + 2(\kappa_{\tilde{\eta}G}^2 J_{GL}^2 + \kappa_{\tilde{\eta}L}^2) + (\kappa_{\tilde{\eta}G}^3 J_{GL}^3 + \kappa_{\tilde{\eta}L}^3)) \end{pmatrix} \\
\end{aligned} \tag{14}$$

3. Track Fitting

Negative-Log-Likelihood.

$$\mathcal{L} = -\ln(\text{all probability density function multiply}) + \text{Constant} \tag{15}$$

$$\begin{aligned}
\mathcal{L} = & \left(\frac{1}{2}\right) \cdot \sum \left((\xi_{\vec{p}} - m_{\vec{p}})^T \sigma_{\vec{p}}^{-1} (\xi_{\vec{p}} - m_{\vec{p}}) \right) \\
& + \left(\frac{1}{2}\right) \cdot \sum \omega_{\tau}^2 \\
& + \left(\frac{1}{2}\right) \cdot \sum \omega_{\rho}^2 \\
& + \left(\frac{1}{2}\right) \cdot \sum (\omega_I + \exp(-\omega_I)) \\
& + \sum (\omega_B t + (1-t) \cdot \ln(\omega_B))
\end{aligned} \tag{16}$$

Negative-Log-Likelihood.

$$\begin{aligned}
H_{\xi} = \frac{\partial \mathcal{L}}{\partial \xi^T} = & \sum \left(\left(\frac{d\vec{p}_{\xi}}{d\xi} \right)^T \sigma_m^{-1} (\vec{p}_{\xi} - \vec{p}_m) \right) + \\
& \sum \left(\left(\frac{\partial \omega_L}{\partial \xi} \right)^T \cdot \omega_L \right) + \\
& \sum \left(\left(\frac{\partial \omega_{\theta}}{\partial \xi} \right)^T \cdot \omega_{\theta} \right) + \\
& \sum \left(\left(\frac{\partial \omega_I}{\partial \xi} \right)^T \cdot \left(\frac{1}{2} \right) (1 - \exp(-\omega_I)) \right) + \\
& \sum \left(\left(\frac{\partial \omega_B}{\partial \xi} \right)^T \cdot (t + (1-t) \cdot \omega_B^{-1}) \right)
\end{aligned} \tag{17}$$

Negative-Log-Likelihood.

$$\begin{aligned}
C_{\xi}^{-1} = \frac{\partial \mathcal{L}}{\partial \xi^T \partial \xi} = & \sum \left(\left(\frac{d\vec{p}_{\xi}}{d\xi} \right)^T \sigma_m^{-1} \left(\frac{d\vec{p}_{\xi}}{d\xi} \right) \right) + \\
& \sum \left(\left(\frac{\partial \omega_L}{\partial \xi} \right)^T \cdot \left(\frac{\partial \omega_L}{\partial \xi} \right) \right) + \\
& \sum \left(\left(\frac{\partial \omega_{\theta}}{\partial \xi} \right)^T \cdot \left(\frac{\partial \omega_{\theta}}{\partial \xi} \right) \right) + \\
& \sum \left(\left(\frac{\partial \omega_I}{\partial \xi} \right)^T \cdot \left(\frac{1}{2} \right) \exp(-\omega_I) \cdot \left(\frac{\partial \omega_I}{\partial \xi} \right) \right) + \\
& \sum \left(\left(\frac{\partial \omega_B}{\partial \xi} \right)^T \cdot ((t-1) \cdot \omega_B^{-2}) \cdot \left(\frac{\partial \omega_B}{\partial \xi} \right) \right)
\end{aligned} \tag{18}$$

Minimization. The Gradient Descent Method

$$\Delta\xi = c \cdot H_\xi \text{ where } c \text{ is constant} \quad (19)$$

The Gauss-Newton Method

$$\Delta\xi = C_\xi \cdot H_\xi \quad (20)$$

The Levenberg-Marquardt Method

$$\Delta\xi = (C_\xi + c \cdot \text{diag}(C_\xi)) \cdot H_\xi \text{ where } c \text{ is constant} \quad (21)$$

4. The Elsevier article class

5 *Installation.* If the document class *elsarticle* is not available on your computer, you can download and install the system package *texlive-publishers* (Linux) or install the L^AT_EX package *elsarticle* using the package manager of your T_EX installation, which is typically T_EX Live or MikT_EX.

Usage. Once the package is properly installed, you can use the document class
10 *elsarticle* to create a manuscript. Please make sure that your manuscript follows the guidelines in the Guide for Authors of the relevant journal. It is not necessary to typeset your manuscript in exactly the same way as an article, unless you are submitting to a camera-ready copy (CRC) journal.

Functionality. The Elsevier article class is based on the standard article class
15 and supports almost all of the functionality of that class. In addition, it features commands and options to format the

- document style
- baselineskip
- front matter
- 20 • keywords and MSC codes
- theorems, definitions and proofs

- lables of enumerations
- citation style and labeling.

5. Front matter

25 The author names and affiliations could be formatted in two ways:

- (1) Group the authors per affiliation.
- (2) Use footnotes to indicate the affiliations.

See the front matter of this document for examples. You are recommended to conform your choice to the journal you are submitting to.

30 6. Bibliography styles

There are various bibliography styles available. You can select the style of your choice in the preamble of this document. These styles are Elsevier styles based on standard styles like Harvard and Vancouver. Please use BibTeX to generate your bibliography and include DOIs whenever available.

35 Here are two sample references: [1, 2]. Here are two sample references: [3].

Propagation. It uses RK4 method.

References

- [1] R. Feynman, F. Vernon Jr., The theory of a general quantum system interacting with a linear dissipative system, *Annals of Physics* 24 (1963) 118–173.
40 doi:10.1016/0003-4916(63)90068-X.
- [2] P. Dirac, The lorentz transformation and absolute time, *Physica* 19 (1–12) (1953) 888–896. doi:10.1016/S0031-8914(53)80099-6.
- [3] A. Strandlie, R. Frühwirth, Track and vertex reconstruction: From classical to adaptive methods, *Reviews of Modern Physics* 82 (2) (2010) 1419–1458.
45 doi:10.1103/RevModPhys.82.1419.
URL <http://link.aps.org/doi/10.1103/RevModPhys.82.1419>