

Liver imaging report — FDG PET-CT of 23 September 2026 and 19 June 2026

Anastasia Zherlitsyna, born 25.04.1976 — metastatic GIST with liver metastases

Auto-generated summary, compiled 27 September 2026 from the patient's records
(docs_eng archive)

Contents

1 Purpose and sources	2
2 Formulated diagnosis	2
3 Current situation (as of 27 September 2026)	3
4 History	4
5 The 23.09.2026 PET-CT (KMKOC)	5
5.1 Independent segmentation of the 23.09.2026 study	5
6 The 19.06.2026 PET-CT: three readings	11
7 Independent image analysis of the 19.06.2026 study	12
7.1 Data and method	12
7.2 Findings	12
7.3 Second FDG-active site: the wall of the right-lobe mass	14
7.4 3D reconstruction	16
8 Evolution of the liver disease 2023–2026	18
9 Gastric-wall GIST, uterus and right ovarian endometrioma 2022–2026	19
9.1 Gastric-wall mass (primary GIST site)	19
9.2 Uterus with leiomyomas and right adnexal mass (whole pelvic complex traced)	22
10 Feasibility of liver-directed treatment	24
10.1 Opinion of the interventional radiologist (Dr. Oleksandr Korshak, Shalimov National Scientific Centre of Surgery and Transplantology, Kyiv; 10–11.09.2026, messenger consultation)	24
10.2 Guideline basis for local treatment of focal progression	25
10.3 Evidence summary (OpenEvidence review, 05.09.2026)	25
10.4 Proposed systemic drugs and coverage of the three KIT clones	25
11 Questions for the multidisciplinary tumour board	26
12 Appendix A — PET-CT report, KMKOC, 23.09.2026 (English translation)	29
13 Appendix B — Expert opinion on the FDG PET-CT, 08.09.2026 (English translation)	31
14 Appendix C — LISOD retrospective radiology report, 08.07.2026 (English translation)	33
15 Appendix D — PET-CT report, Feofaniya, 19.06.2026 (English translation)	34
16 Appendix E — Slice-by-slice verification of the lesion outline	36

1 Purpose and sources

This document is an auto-generated summary, compiled and formatted by software from the patient's records and from image analysis by the family; it has not been reviewed by a physician. It is not a radiology report or a medical opinion, and it does not replace the original documents, which are linked throughout. Please verify any figure used for clinical decisions against the source document or fresh imaging.

This report gathers, in one place, everything that the available imaging says about the liver disease of the patient as of September 2026, so that an interventional radiologist, hepatobiliary surgeon or oncologist can judge the feasibility of liver-directed treatment without opening the full archive. It contains: (1) the formulated diagnosis and current clinical situation; (2) the reading of the latest FDG PET-CT of 23.09.2026 and an independent AI-assisted segmentation of that study (volumes, June → September comparison, 3D reconstruction); (3) the three professional readings of the 19.06.2026 FDG PET-CT; (4) an independent quantitative image analysis of the June study with raw and labelled CT slices and a 3D reconstruction, including the second FDG-active site in the right-lobe mass; (5) the evolution of the liver lesions across the five PET-CTs of 2023–2026; (6) the opinion of the thermal-ablation specialist consulted on 10.09.2026; (7) a summary of the treatment-option evidence reviewed with OpenEvidence on 05.09.2026.

The image analysis was produced by the patient's family with open-source software on the original DICOM data and is **not** a radiologist's report. It is included to make the anatomy explicit; measurements should be confirmed on fresh imaging before any procedure.

Sources are held on the family's Google Drive (`docs_eng/`): the Ukrainian originals, English translations, DICOM discs (`anastasia_disks/`) and the overview document `Overview.docx`.

2 Formulated diagnosis

ICD-10: C16.9 (malignant neoplasm of stomach, unspecified); ICD-O morphology 8936/3, gastrointestinal stromal tumour.

Diagnosis: Gastrointestinal stromal tumour (GIST) of the stomach, KIT exon 11 deletion (Q556_V559>H), with metastases to the liver (bilobar, dominant right-lobe mass) and peritoneum, stage IV. Initially (08/2022) diagnosed and treated as leiomyosarcoma of the uterine body T2NxM1; diagnosis revised 04–06/2023 on histology (DOG-1, CD117 positive; Israeli pathology revision of all samples). Concomitant: uterine leiomyomas; right ovarian endometrioma; type 2 diabetes; iron-deficiency anaemia.

- **08/2022** — liver-metastasis biopsy read as metastatic uterine leiomyosarcoma; bilobar liver masses, paragastric mass 51 × 46 mm, pelvic masses.
- **09/2022 – 01/2023** — status after 4 cycles gemcitabine + docetaxel + bevacizumab: progression in the liver (MRI 10/2022, CT 12/2022). Oncomine (12/2022): KIT exon 11 deletion (VAF 42.5 %), subclonal KIF5B–MET fusion, TMB low, MSS; germline panel negative.
- **05–12/01/2023** — pazopanib 800 mg, stopped on day 8: posterior reversible encephalopathy syndrome with left occipital haemorrhage, hospitalised 13–25/01/2023, resolved; CT 09/02/2023 with 12 % regression.
- **01/2023** — repeat liver biopsy: reported as metastasis of uterine leiomyosarcoma.
- **02–03/2023** — status after 3 cycles trabectedin: progression on PET-CT 11/04/2023.
- **04–06/2023** — EUS-FNA of the paragastric mass and liver core biopsy: GIST of the stomach, GIST metastasis in the liver (CSD, 21–26/04/2023); Israeli revision 04/06/2023 confirms GIST in all samples.
- **03/05/2023** — imatinib 400 mg/day started.
- **08/2023** — marked metabolic response (PET-CT).
- **03/2024** — stable.
- **05/2025** — stable, “PET negative”, cystic residues.
- **06–09/2026** — PET-CT 19/06/2026: return of enhancing, FDG-active mural components in the known liver lesions after 3 years of imatinib (started 03.05.2023) (LISOD re-read 08/07/2026; Israeli expert reading 08/09/2026: metabolic progression, left-lobe lesion 42 × 36 mm, right-lobe mass ~15 × 10 cm, no bone lesions).
- **08/2026** — FoundationOne Liquid CDx (ctDNA fraction 32 %): KIT exon 11 Q556_V559>H (17.0 %) plus secondary KIT exon 13 V654A (8.8 %) and exon 18 A829P (4.0 %) — polyclonal imatinib resistance.
- **08/09/2026** — recommendation: sunitinib 50 mg 4/2 (regorafenib 160 mg d1–21 as an option); not yet started; imatinib continues.

- **10/09/2026** — thermal ablation of the left-lobe lesion judged contraindicated by Dr. O. Korshak (interventional radiology, Shalimov Centre, Kyiv): contact with the left portal branch and bile duct; other local options under discussion.
- **17/09/2026** — Dr. E. Tahover (GIST oncologist, Tel Aviv Sourasky): imatinib has failed; start a second-line TKI now (sunitinib or ripretinib; regorafenib an alternative); liver-directed interventional treatment first.
- **18/09/2026** — Dr. D. Fedorov (liver surgeon, Kyiv): two-stage liver resection feasible and potentially radical (right hemihepatectomy first, then the segment II/III focus after regeneration); workup after fresh PET-CT.
- **23/09/2026** — PET-CT (KMKOC): increased FDG activity in multiple liver foci vs 19.06.2026 (SUVmax 4.9) — progression; left-lobe metabolic area 35 × 54 mm, confluent; new active focus 15 × 24 mm in segment S8; gastric and pelvic lesions stable. Second-line drug not started as of 27.09.2026.

3 Current situation (as of 27 September 2026)

Extent of the current progression. This is not a single small nodule, and it grew between June and September:

- **PET-CT 23.09.2026 (KMKOC):** FDG activity increased in multiple liver foci compared with 19.06.2026, SUVmax up to 4.9 — progression of the underlying disease.
- **Left lobe (segments II/III):** the active area is now confluent, up to 35 × 54 mm (June CT lesion 46 × 48 × 42 mm, 27 mL; expert reading 42 × 36 mm). On the June contrast CT it touches the left portal branch and the left hepatic duct.
- **Right lobe:** a new active focus up to 15 × 24 mm in segment S8, inside the ametabolic area of the dominant mass (now 98 × 148 mm). In June the right-lobe activity was a plaque on the wall of that mass.
- **Own volumetry (section 5):** FDG-active tissue in the left lobe **19 → 36 mL** (2.2 → 3.4 × liver background); new 4 mL focus in S8; active wall of the right-lobe mass 6 → 10 mL.
- **ctDNA tumour fraction 32 %** (FoundationOne Liquid CDx, 20.08.2026) is high and points to a substantial amount of active tumour, carrying two resistance clones (KIT exon 13 V654A and exon 18 A829P).

Patient. Anastasia Zherlitsyna, born 25.04.1976 (50), Kyiv, Ukraine.

Diagnosis. Metastatic gastric GIST with bilobar liver metastases; treated as uterine leiomyosarcoma T2NxM1 from 28.09.2022 until the April–May 2023 biopsies established GIST (DOG-1 and CD117 positive; Israeli revision 04.06.2023 confirmed GIST in all three samples). Uterine leiomyomas and a right ovarian endometrioma are benign bystanders.

Molecular status (most important). Driver KIT exon 11 deletion Q556_V559>H (Oncomine, Dec 2022, VAF 42.5 %). FoundationOne Liquid CDx of 20.08.2026, ctDNA tumour fraction 32 %: exon 11 driver at VAF 17.0 % plus two new KIT resistance mutations — exon 13 V654A (8.8 %) and exon 18 A829P (4.0 %), in different kinase domains: polyclonal imatinib resistance. The report flags imatinib, sunitinib and avapritinib as resistant and lists regorafenib as NCCN category 1.

Sites of disease (PET-CT 23.09.2026, KMKOC). Multiple ametabolic liver lesions with metabolic activity growing along the contour and inside some of them, SUVmax up to 4.9; dominant right-lobe mass 98 × 148 mm; left-lobe active area up to 35 × 54 mm, confluent; active focus up to 15 × 24 mm in segment S8; calcified FDG-negative gastric lesion 43 × 32 mm, unchanged; uterine lesions stable; new oesophageal uptake (SUVmax 2.0–3.2, possibly inflammatory — endoscopy advised); no bone lesions.

Sites of disease in June (PET-CT 19.06.2026, expert reading 08.09.2026). Dominant right-lobe liver mass about 15 × 10 cm, cystic/necrotic residue, with a 5.7 mL FDG-active plaque on its anterolateral wall plus a 0.3 mL spot on the medial wall (SUVmax 2.34, own image analysis — see below; the reading describes this as “increased peripheral uptake”); left-lobe segment II/III lesion 42 × 36 mm (own CT measurement 46 × 48 × 42 mm, 27 mL), metabolically silent in 2024–2025 and FDG-active in June 2026 — the current progression, in contact with the left portal branch; calcified, near-inactive gastric-wall residue 37 × 31 mm; no bone lesions.

Treatment. Imatinib 400 mg/day since 03.05.2023, continuing. On 30.07.2026 and 08.09.2026 the LISOD oncologist recommended sunitinib 50 mg, 28 days on / 14 off (regorafenib 160 mg days 1–21 q28 as an alternative) — sunitinib has NOT been started as of 27.09.2026 (no second-line drug started); on 17.09.2026 Dr. Tahover urged starting a second-line TKI without further delay.

Safety context. PRES on pazopanib in January 2023 with left occipital haemorrhage and multifocal vasoconstriction, largely resolved by June 2023 — potent VEGFR inhibitors carry extra risk. Baseline echocardiogram 30.07.2026 normal (EF 67 %). Comorbidities on record: type 2 diabetes, iron-deficiency anaemia.

Open clinical question. Local treatment of the FDG-active left-lobe lesion: on 10.09.2026 Dr. Oleksandr Korshak (interventional radiology, Shalimov Centre of Surgery and Transplantology, Kyiv) judged thermal ablation (RFA/MWA) contraindicated because the lesion contacts the left portal branch and the accompanying left bile duct; TARE, TACE/DEB-TACE, SBRT, IRE, resection of segments II–III and non-thermal histotripsy (HistoSonics, newly CE-approved; opinion to be requested from the manufacturer, as Dr. Korshak suggested) are under discussion, with imatinib continued — an approach NCCN and ESMO allow to be considered for limited progression on imatinib (guideline basis below). The PET-CT of 23.09.2026 adds an active focus in segment S8 of the right lobe, which any local plan must cover as well; S8 lies within the part removed by a right hemihepatectomy, the first stage of the two-stage resection Dr. Fedorov proposed on 18.09.2026.

4 History

Period	Systemic treatment	Examination / event	Finding
Aug 2022	—	Liver-metastasis biopsy 29.08.2022; tumour board 31.08.2022	Uterine leiomyosarcoma T2NxM1 (revised to GIST in May 2023)
Sep 2022 – Jan 2023	Gemcitabine + Docetaxel + Avastin, 4 cycles	MRI 14.10.2022, CT 01.12.2022, CT 04.01.2023	Progression in the liver
Dec 2022 – Jan 2023	(same)	Oncomine on the 2022 biopsy; BRCA panel ; BAM files	KIT exon 11 p.Gln556_Val559delinsHis (VAF 42.5 %); subclonal KIF5B-MET fusion; TMB low, MSS; germline negative
5–12 Jan 2023	Pazopanib 800 mg (stopped day 8)	Brain MRI 13.01 and 20.01.2023	PRES with left occipital haemorrhage
Feb – Mar 2023	Trabectedin, 3 cycles	PET-CT 11.04.2023; biopsies 21–25.04.2023	Progression; GIST established
May – Aug 2023	Imatinib 400 mg/day (start 03.05.2023)	PET-CT 17.08.2023	Pronounced response, loss of activity and solid rim components
Mar 2024	Imatinib 400 mg/day (cont.)	PET-CT 14.03.2024, Israeli revision 25.03.2024	Stable, minimal uptake
May 2025	Imatinib 400 mg/day (cont.)	PET-CT 13.05.2025, LISOD read 19.05.2025	PET negative; new inactive 11 mm mural nodule
19 Jun 2026	Imatinib 400 mg/day (cont.)	PET-CT 19.06.2026	First read as stable
8 Jul 2026	Imatinib 400 mg/day (cont.)	LISOD retrospective read	New FDG-active mural components — progression
30 Jul 2026	Imatinib 400 mg/day (cont.); sunitinib 50 mg 4/2 prescribed, not started	LISOD oncologist; echocardiogram	Progression documented; EF 67 %
20 Aug 2026	Imatinib 400 mg/day (cont.)	FoundationOne Liquid CDx , blood	ctDNA 32 %; KIT exon 11 (17 %) + exon 13 V654A (8.8 %) + exon 18 A829P (4.0 %); BRCA2, CDKN2A, MTOR; TMB 3, MSS

Period	Systemic treatment	Examination / event	Finding
8 Sep 2026	Imatinib 400 mg/day (cont.)	Expert PET-CT reading, Israel	Metabolic progression, especially the left-lobe lesion
8 Sep 2026	Imatinib 400 mg/day (cont.)	LISOD oncologist (Ukrainian original)	Sunitinib 50 mg 4/2 recommended; regorafenib 160 mg d1–21 q28 as an option; neither started as of 11.09.2026
17 Sep 2026	Imatinib 400 mg/day (cont.)	Oncology consultation, Dr. E. Tahover (Tel Aviv Sourasky)	Start second-line TKI now (sunitinib or ripretinib; regorafenib alternative); interventional radiology first for the liver
18 Sep 2026	Imatinib 400 mg/day (cont.)	Surgical consultation, Dr. D. Fedorov	Two-stage liver resection feasible, potentially radical; fresh PET-CT, then workup
23 Sep 2026	Imatinib 400 mg/day (cont.); second line not started	PET-CT, KMKOC (translation, Ukrainian original)	Increased FDG activity in multiple liver foci — progression; new S8 focus; gastric and pelvic lesions stable

5 The 23.09.2026 PET-CT (KMKOC)

Low-dose **non-contrast** CT and FDG PET, GE Discovery STE16, 260 MBq (DICOM header: 230 MBq), 80 min uptake; compared with 19.06.2026. [English translation](#) · [Ukrainian original](#) · [DICOM disc](#). The full report is Appendix A.

Radiologist's conclusion: increased metabolic activity in multiple liver foci compared with 19.06.2026 — progression of the underlying disease. Largest right-lobe lesion 98 × 148 mm; left-lobe metabolic area up to 35 × 54 mm (confluent); focus up to 15 × 24 mm inside the ametabolic area in segment S8; SUVmax up to 4.9 (blood pool 1.6). Gastric lesion on the lesser curvature 43 × 32 mm, calcified, FDG-negative, stable. Anterior uterine lesion 67 × 49 mm (SUVmax 1.8) and ametabolic node 37 × 47 mm, stable. Oesophageal uptake SUVmax 2.0–3.2, possibly inflammatory — EGD recommended. No bone lesions.

5.1 Independent segmentation of the 23.09.2026 study

Method. The radiologist's report and automatic organ maps (TotalSegmentator) served only as starting points. Every structure was outlined slice by slice by AI segmentation agents and checked by independent adversarial reviewer agents, which re-rendered every slice and tried to refute each outline. Outlines were corrected until they passed review; all passed on the second review. Liver background SUV 1.97 ± 0.63 ; FDG-active threshold **3.24** (liver mean + 2 SD); blood pool 1.58. **This is an AI-assisted analysis, not a radiology report.** Without contrast, vessels, bile ducts and small or isodense lesions cannot be assessed.

Structure	23.09.2026 (own measurement)	19.06.2026	Radiologist 23.09.2026
FDG-active tissue, left lobe (II/III)	36.3 mL ; 60 × 29 mm axial, 52 mm craniocaudal; SUVmax 6.66, 3.4 × liver	19 mL; SUVmax 2.27, 2.2 × liver	35 × 54 mm, confluent
Left-lobe lesion, CT outline	35 ± 4 mL; 45 × 30 mm axial, 56 mm craniocaudal	27 mL; 46 × 48 × 42 mm	—

Structure	23.09.2026 (own measurement)	19.06.2026	Radiologist 23.09.2026
FDG-active focus, segment S8 (new)	4.1 mL ; 23 × 13 mm, 29 mm craniocaudal; SUVmax 6.75, 3.4 × liver	not active	15 × 24 mm
FDG-active wall of the right-lobe mass	9.7 mL in 9 foci; anterolateral plaque 6.8 mL (3.1 × liver); medial spot next to the IVC 0.9 mL	5.7 mL plaque (2.6 × liver) + 0.3 mL spot	“multiple foci”
Dominant right-lobe mass	1200 mL; 146 × 116 mm axial, 180 mm craniocaudal (incl. the 74 mL S8 lobule)	~1217 mL	98 × 148 mm
Other liver lesions	none found (lesions < ~8 mm cannot be excluded)	—	—
Gastric-wall lesion	36 mL; 42 × 31 mm; calcified, FDG-cold	43 × 33 × 44 mm	43 × 32 mm
Anterior uterine lesion	104 mL; 66 × 51 mm; ≈ 1.0 × liver	—	67 × 49 mm
Upper-margin pelvic node	23 mL; 43 × 37 × 29 mm; not active	—	37 × 47 mm

Our voxel SUVmax (6.66–6.75) exceeds the report’s 4.9. The dose difference in the DICOM header (230 vs 260 MBq) explains about 13 %; the rest is voxel maximum vs reader ROI. **Ratios to liver background** are the comparable measure between studies: left lobe 2.2 → 3.4 × liver, mass wall 2.6 → 3.1 × liver. The right-lobe mass short axis differs from the report because of how it is measured; our narrowest slices also give 98 mm.




Figure 1: 3D close-up of the left lobe, 23.09.2026: CT lesion outline (cyan) and the FDG-active tissue (yellow), which extends beyond the lesion on its medial and posterior side; green = right-lobe mass.




Figure 2: Left-lobe FDG-active tissue on twelve axial levels: FDG-active tissue yellow, CT lesion outline cyan, liver green; patient right on the image left, anterior up.



Figure 3: Anterior 3D view of the liver, 23.09.2026: right-lobe mass (green, image left = patient right), left-lobe lesion (cyan) and all FDG-active tissue (yellow) — the S8 focus at the top of the mass, the anterolateral wall plaque and small wall foci.




Figure 4: 3D close-up of the segment S8 focus (yellow) inside the upper lobule of the right-lobe mass.




Figure 5: Segment S8 focus, axial slices.




Figure 6: Right-lobe mass outline with the FDG-active wall tissue, axial slices.




Figure 7: Gastric-wall lesion (orange), calcified and FDG-cold. The SUVmax in each panel title is the hottest voxel anywhere in the panel, not in the lesion.




Figure 8: Pelvis: anterior uterine lesion (magenta), upper-margin node (cyan), uterus/pelvic complex (green). The right ovarian endometrioma cannot be separated on this non-contrast study; the panel-title SUVmax is bladder urine.

6 The 19.06.2026 PET-CT: three readings

	Feofaniya, 19.06.2026 (primary read)	LISOD, 08.07.2026 (retrospective read, Dr. M. Novikov)	Israel, 08.09.2026 (expert opinion, Dr. V. Kulikova)
Documents	translation (Ukrainian original)	translation (Ukrainian original)	translation (Russian original)
Comparison	prior studies	13.05.2025 and 14.03.2024	13.05.2025

	Feofaniya, 19.06.2026 (primary read)	LISOD, 08.07.2026 (retrospective read, Dr. M. Novikov)	Israel, 08.09.2026 (expert opinion, Dr. V. Kulikova)
Right-lobe mass	known cystic residue	new enhancing, moderately FDG-active mural components; liver enlarged to 18.7 cm	about 15 × 10 cm (was 11–12.5 cm), peripheral uptake somewhat more pronounced
Left-lobe lesion	—	mural components with uptake (SUVmax about 3.2 reported)	about 42 × 36 mm, size unchanged, uptake intensity increased
Gastric-wall mass	unchanged	unchanged, calcifications	about 37 × 31 mm, minimal uptake, unchanged
Portal vein / ducts	—	—	portal vein patent; intrahepatic ducts not dilated
Bones	—	31 mm lucent lesion, right femoral head (to characterise)	no pathological foci or destruction in the scanned area
Conclusion	initially read as stable	return of viable mural tumour components — progression	metabolic progression of the liver lesions, especially the left-lobe lesion; enlargement of the right-lobe mass with somewhat increased activity

Full texts of the LISOD and Israeli readings are reproduced in the appendices.

7 Independent image analysis of the 19.06.2026 study

7.1 Data and method

The study contains a low-dose CT, an arterial-phase and a **venous-phase contrast CT (series 9, 2 mm, 412 slices)** and the whole-body FDG PET (SUVbw, 2.67 × 2.67 × 5 mm). The venous-phase CT and the PET were converted from DICOM with the correct patient geometry and fused in world coordinates. Organ, vessel and tumour masks were produced with TotalSegmentator on the corrected volume and reviewed slice by slice; the lesion outline is a density threshold (hypodense relative to enhanced liver) seeded at the FDG peak, and the FDG-active volume is the connected region above 50 % of the peak-to-background contrast. Display convention: radiological (anterior up, patient left on image right).

Note on an earlier analysis. A first automated analysis of this study (September 2026) used NIfTI volumes whose x and y axes were swapped relative to their header; laterality derived from those files was wrong and the PET was misregistered by about 7.6 mm per axis. That analysis was discarded; everything below comes from the corrected frame and agrees with the 08.09.2026 expert reading.

7.2 Findings

Finding	Measurement (corrected frame, RAS mm)
Left-lobe lesion, CT	segment II (83 %) / III (17 %); hypodense, mean 44 HU vs liver 98 HU; 46 × 48 × 42 mm , 27 mL; centre (−54, 173, −1201)
FDG-active tumour	SUVmax 2.27 (liver background 1.04); 19 mL above the 1.65 threshold; about half inside the hypodense core, the rest isodense enhancing tissue on the medial (hilar) side and inferior pole

Finding	Measurement (corrected frame, RAS mm)
Left portal branch	in contact with the lesion (0.8 mm) and with the FDG-active part (0 mm); contact point (−35, 164, −1200)
Distances to other structures	IVC 31 mm; stomach 19 mm; heart 28 mm; the lesion reaches the liver surface (depth 0–31 mm)
Bile ducts	not resolvable on CT (not dilated); the left hepatic duct runs in the same Glissonean sheath as the left portal branch, so it lies within a few millimetres of the lesion

SUV values differ between readers and reconstructions (LISOD reported SUVmax about 3.2 for the same lesion); the relative pattern is the finding, not the absolute value. Sub-2 mm distances measured on a 2 mm CT with AI vessel masks mean “in contact”.




Figure 9: Left-lobe lesion, 19.06.2026. Top row: clean venous-phase CT (three axial levels and one coronal). Bottom row: the same slices with FDG uptake (SUV > 1.3), CT lesion outline (cyan), FDG-active tumour (yellow), portal vein (blue) and other liver vessels (green). The enhancing vessel touching the medial edge of the lesion is the left portal branch.




Figure 10: Nine consecutive axial slices (3 mm apart) through the lesion, same overlays as Figure 1.

7.3 Second FDG-active site: the wall of the right-lobe mass

The right-lobe mass is not uniformly inactive. Its wall was searched for tissue above liver background + 2 SD (threshold SUV 1.59, liver background 0.90) in a band spanning 15 mm inside and 8 mm outside the mass outline, then compared against blood-pool activity (IVC 0.94, portal vein 1.09) to exclude vascular spill-over.

Finding	Measurement
Main component	5.7 mL plaque on the anterolateral wall , 27 × 24 × 54 mm, SUVmax 2.34 (2.6 × liver background), soft-tissue density (37 HU vs 15 HU cyst fluid)
Second component	0.3 mL spot on the medial wall , near the portal vein, SUVmax 1.83
Rest of the wall / cyst	below threshold at every other point checked

This is what the professional readings describe as “increased peripheral uptake” (Feofaniya, 19.06.2026) and enhancing mural components (LISOD retrospective read). Its intensity (SUVmax 2.34) is essentially the same as the left-lobe focus (2.27) — the right-lobe mass carries an active site in addition to the left-lobe lesion, not instead of it.




Figure 11: FDG-active wall of the right-lobe mass: twelve axial levels through the 5.7 mL component (cyan outline, $SUV \geq 1.59$), mass outline in green, anterior up, patient right on the image left.




Figure 12: Coronal and axial maximum-intensity projections of FDG uptake in and around the liver: green = right-lobe mass outline, cyan = its FDG-active wall, yellow = the left-lobe lesion outline (axial panel).

7.4 3D reconstruction

This scene now includes both FDG-active sites — the left-lobe focus and the newly traced active wall of the right-lobe mass — matching the interactive 3D model on the case website.




Figure 13: 3D scene, anterior view: liver ghosted, right-lobe mass translucent green, its FDG-active wall opaque yellow, portal vein and branches blue, left-lobe CT lesion translucent cyan, its FDG-active tumour opaque yellow, dotted orange = possible course of the left bile duct (anatomical assumption offset antero-superior to the portal branch; not visible on CT), red = contact point.



Figure 14: Close-up on the right-lobe mass and its FDG-active wall (yellow), with the left-lobe lesion visible behind.




Figure 15: Close-up, left anterior oblique: the FDG-active tissue wraps the terminal left portal branch.




Figure 16: Inferior view of the same scene.

An interactive version of this model — with layers that can be switched on and off — is on the case website ([viewer_all.html](#)); a static glTF export is in the Drive folder docs_eng/2026_new/Left_lobe_focus_2026-09-10/.

8 Evolution of the liver disease 2023–2026




Figure 17: Dominant right-lobe mass through its largest cross-section on all six PET-CTs (24 cm field). Top: CT. Bottom: CT with FDG uptake above $1.3 \times$ liver background and the mass outline (automated until June 2026; reviewed agent segmentation on 23.09.2026).




Figure 18: Left-lobe lesion region on the last four PET-CTs (16 cm field). Top: CT. Bottom: CT with FDG uptake above 1.3 × liver background and the lesion outline.

Study	Left-lobe lesion, CT (mm)	Volume (mL)	SUVmax	Uptake / liver background
14.03.2024	59 × 49 × 40	26.9	PET not SUV-calibrated	1.29
13.05.2025	46 × 45 × 32	13.2	1.74 (liver 1.46)	1.19
19.06.2026	46 × 48 × 42	27.4	2.20 (liver 1.16)	1.90
23.09.2026	41 × 46 × 56	35.1	6.13 (liver 2.04)	3.01

The lesion is a pre-existing hypodense metastasis (present at the same site in March 2024 and May 2025 with uptake at or below liver background) that became FDG-active in June 2026 — the pattern of an emerging imatinib-resistant clone inside a treated lesion — and was more active and larger on 23.09.2026 (the 23.09.2026 values come from the reviewed agent segmentation; that CT is non-contrast). Outlines were checked slice by slice on each date. The 2023–2024 PET series were not exported in SUV units, so uptake is given relative to the liver background for all dates.

9 Gastric-wall GIST, uterus and right ovarian endometrioma 2022–2026

On each of the eleven studies (five CTs of 2022–2023 from the DICOM discs and six PET-CTs) the gastric-wall mass and the pelvic mass complex were traced slice by slice from a hand-placed seed (soft-tissue density window, air and bone excluded, growth limited to tissue continuous with the seed) and every traced slice was checked visually. The 23.09.2026 outlines come from the reviewed agent segmentation of that study (low-dose non-contrast CT), so its volumes are not strictly comparable with the earlier traces. The earlier PET-CTs were acquired with intravenous contrast; the 2023–2024 PET series were not exported in SUV units, so uptake is given relative to the liver background throughout.

9.1 Gastric-wall mass (primary GIST site)

PET-CT	Max in-plane diameter (mm)	Volume (mL)	Mean HU	Max uptake / liver	Reported
23 Aug 2022 (CT)	55	53	63	CT only	51 × 46 mm
1 Dec 2022 (CT)	53	65	65	CT only	56 × 46 mm (German re-read 5.2 × 6.1 cm)
4 Jan 2023 (CT)	52	70	58	CT only	65 × 50 mm
9 Feb 2023 (CT)	50	52	51	CT only	55 × 40 mm
11 Apr 2023	54	36	75	5.5 × (not SUV-calibrated)	58 mm, SUVmax 8.2, active
28 Jun 2023 (CT)	54	47	43	CT only	47 × 35 mm, low density
17 Aug 2023	48	46	41	1.8 × (not SUV-calibrated)	44 mm, almost fluid, inactive
14 Mar 2024	49	35	45	1.3 × (not SUV-calibrated)	40 mm, inactive
13 May 2025	43	23	40	1.8 ×	35 mm, calcifications
19 Jun 2026	41	24	38	1.1 ×	43 × 33 × 44 mm (Feofaniya); 37 × 31 mm, calcified, unchanged (Israel)
23 Sep 2026	41	36	51	1.2 ×	43 × 32 mm, calcified, FDG-negative, stable (KMKOC; non-contrast study)

The gastric-wall tumour collapsed on imatinib: from an enhancing, intensely active mass in April 2023 (about 5.5 × liver) to a larger but low-density, almost fluid, near-inactive residue by August 2023 (cystic change, as the report describes), then a plateau at about 4 cm through 2024–2026 with progressive calcification and uptake at or near liver background — in agreement with the reports. In April 2023 the mass was traced from its FDG-avid volume (restricted to soft-tissue density) because its enhancement matched the adjacent tissue on CT. On 23.09.2026 it is unchanged in linear size (41 mm) and FDG-cold; its 36 mL against 24 mL in June reflects the method (the September outline includes the calcified lateral clumps the June trace left out), not growth.




Figure 19: Gastric-wall GIST through its largest cross-section on all eleven studies 2022–2026 (12 cm field); amber titles = contrast CT, cyan titles = PET-CT with FDG above 1.3 x liver background. Cyan = traced outline.




Figure 20: 3D reconstruction of the traced gastric-wall mass on the eleven dates at one scale (bar = 2 cm), anterior view.




Figure 21: The same masses with ghost organs from the automatic segmentation of each study, clipped to a box around the mass: stomach (lilac), pancreas (sand), spleen (rose), liver edge (pale), left kidney (grey); bar = 5 cm.

9.2 Uterus with leiomyomas and right adnexal mass (whole pelvic complex traced)

PET-CT	Max in-plane diameter (mm)	Volume (mL)	Mean HU	Max uptake / liver	Reported
23 Aug 2022 (CT)	67	175	59	CT only	fundal mass 93 × 63 mm; right adnexal cyst 67 × 57 mm
1 Dec 2022 (CT)	82	167	53	CT only	fundal mass 87 × 64 mm
4 Jan 2023 (CT)	61	145	44	CT only	fundal mass 75 × 55 mm, uterus 45 × 42 mm; right ovarian formation 75 × 45 mm
9 Feb 2023 (CT)	66	164	44	CT only	fundal mass 75 × 55 mm
11 Apr 2023	96	178	62	1.8 × (not SUV-calibrated)	inactive uterine mass 72 mm; right ovarian mass ~6 cm
28 Jun 2023 (CT)	90	228	83	CT only	pedunculated myoma 90 × 70 mm; right ovarian formation 65 × 42 mm
17 Aug 2023	108	189	59	1.5 × (not SUV-calibrated)	96 mm, inactive
14 Mar 2024	98	148	65	1.3 × (not SUV-calibrated)	78 mm; Israeli revision 6.5 cm and 4.8 × 5.1 cm
13 May 2025	108	134	55	1.6 ×	84 mm

PET-CT	Max in-plane diameter (mm)	Volume (mL)	Mean HU	Max uptake / liver	Reported
19 Jun 2026	92	153	51	2.4 ×	68 × 55 × 48 mm + 46 × 37 mm, stable anterior-wall lesion 67 × 49 mm, SUVmax 1.8, and ametabolic node 37 × 47 mm, stable (KMKOC; non-contrast study; uptake maximum is bladder-urine spill-over)
23 Sep 2026	113	248	31	2.2 ×	

The traced object is the whole pelvic complex (uterine body, its fibroids and the right ovarian endometrioma, traced as one object because fibroid, myometrium and the non-enhancing cyst cannot be separated reliably on every study), which is why it is larger than the single-mass diameters in the reports. Over four years it stays between 130 and 190 mL with no consistent trend and low uptake (1.3–1.8 × liver; the 2.4 × in June 2026 comes from a bowel loop touching the outline) — the behaviour of leiomyoma, not of the GIST metastases. Our numbers exceed the single-mass diameters in the reports because the whole complex is traced; on some dates the outline may include an adjacent bowel loop. The 248 mL on 23.09.2026 is mostly a difference of method, not growth: the September outline reaches lower, to the bladder dome, and includes the anterior lesion that the June trace cut off at its lower edge. Over the same levels the difference is between –4 % and +41 %, within the 134–228 mL spread of the earlier traces, and the radiologist reports both pelvic lesions as stable.




Figure 22: The pelvic complex through its largest cross-section on all eleven studies (19 cm field); amber titles = contrast CT, cyan titles = PET-CT with FDG above 1.3 × liver background. Cyan = traced outline.




Figure 23: 3D reconstruction of the traced pelvic complex on the eleven dates at one scale (bar = 2 cm), anterior view.




Figure 24: The same masses with ghost organs: bladder (yellow), hip bones, sacrum and femoral heads (grey), colon (sand), clipped to a box around the mass; bar = 5 cm.

10 Feasibility of liver-directed treatment

10.1 Opinion of the interventional radiologist (Dr. Oleksandr Korshak, Shalimov National Scientific Centre of Surgery and Transplantology, Kyiv; 10–11.09.2026, messenger consultation)

Thermal ablation (RFA, MWA) is contraindicated for this lesion because two contraindications coincide: proximity to the left branch of the portal vein and, above all, direct adjacency to the left bile duct. Y-90 radioembolisation (TARE) costs from about EUR 40,000 and is not performed in Ukraine (nor in Czechia or Slovakia); whether transarterial chemoembolisation (DEB-TACE/TACE) or surgical resection is possible was still under discussion. On 11.09.2026 the same specialist was asked about focused-ultrasound histotripsy (HistoSonics, newly approved in Europe): a new and very expensive method without results in large numbers of GIST patients or for lesions of this size, results from other liver tumours cannot be transferred; he suggested asking HistoSonics directly.

Consistency with the imaging: the image analysis supports this opinion. The lesion abuts the left portal

branch and the metabolically active tumour is exactly on that side, so an ablation margin would have to include the branch and the accompanying duct.

10.2 Guideline basis for local treatment of focal progression

NCCN (von Mehren et al., JNCCN 2022;20(11):1204–1214; NCCN Guidelines for Patients, GIST 2023) and ESMO–EURACAN–GENTURIS (Casali et al., Ann Oncol 2022;33:20–33) both recommend, for limited progression on imatinib, continuing the TKI and treating the progressing lesion locally (resection if feasible, ablation, embolization or chemoembolization), reserving a change of systemic drug for generalised progression. In the Dutch GIST Registry (Brink et al., Eur J Surg Oncol 2023;49:106942) 123 of 457 metastatic patients had elective surgery or ablation; local treatment was associated with better overall survival (HR 0.56), with the best results when disease was confined to the liver (HR 0.27) and worse results when it was progressing on systemic treatment (HR 3.9). Ablation series for focal progression on imatinib (Dileo 2004, Hasegawa 2007, Avritscher and Gupta 2009, summarised at gistsupport.org) report complete ablation without complications and progression-free intervals of 8–16 months, with the caveat that RFA applies to lesions “not close to blood vessels or ducts” — the reason it was judged contraindicated here.

10.3 Evidence summary (OpenEvidence review, 05.09.2026)

- **Pattern.** Imatinib-refractory, liver-limited oligoprogression (“nodule within a mass”) driven by polyclonal secondary KIT resistance, while the exon-11 bulk disease still responds to imatinib. A combined local-plus-systemic strategy is favoured over a plain TKI switch, and imatinib should continue through any local therapy (2-year PFS after RFA 75 % with continued TKI vs 29 % without).
- **Local options for a viable nodule inside a cystic mass.** Arterially directed therapy (DEB-TACE/TACE, Y-90 TARE; NCCN 2A) deposits its effect only in perfused tissue; microwave ablation (after cyst aspiration) or cryoablation for a few well-defined nodules; **SBRT** for a perivascular or central nodule unsafe for a probe (prospective GIST series: 80 % stable target lesions, median 16 months); **IRE** as the vessel- and duct-sparing alternative for a nodule abutting major vessels or central ducts (GIST data anecdotal); RFA is the least suitable in fluid-containing targets.
- **Timing.** Ablation planning needs multiphase CT or MRI no older than about 30 days; the June scan is stale and the 32 % ctDNA fraction argues for re-imaging now and definitive local therapy within about a month, provided the disease is still oligoprogressive.
- **Systemic.** Sunitinib covers V654A but not A829P; regorafenib and ripretinib cover A829P and regorafenib remains clinically reasonable against V654A (VOYAGER ctDNA data); sorafenib has the broadest in-vitro coverage but only retrospective GIST evidence. The PRES history raises the stakes of potent VEGFR inhibitors: reduced starting dose and strict blood-pressure monitoring. Clinical-trial referral is recommended.
- **Extrahepatic disease.** The femoral-head lesion flagged by LISOD should be characterised (MRI), because extrahepatic disease reduces the value of arterial therapies; the Israeli reading found no bone lesions.

10.4 Proposed systemic drugs and coverage of the three KIT clones

Merged from the [FoundationOne report](#) (therapy list, NCCN categories, resistance flags) and the [OpenEvidence review](#) (subgroup data from INTRIGUE and VOYAGER, toxicity). Median PFS values are trial subgroup figures.

Drug / approach	FoundationOne	V654A (exon 13)	A829P (exon 18)	VEGFR / PRES risk	Status for her
Imatinib 400 mg (current)	resistance flagged	resistant	resistant	none	continuing; keep through any local therapy
Sunitinib 50 mg 4/2	resistance flagged (A829P)	strong (mPFS 15.0 mo)	resistant (2.3 mo)	high (grade ≥3 HTN 22.6 %, PRES warning)	prescribed 30.07 and 08.09.2026, not started

Drug / approach	FoundationOne	V654A (exon 13)	A829P (exon 18)	VEGFR / PRES risk	Status for her
Regorafenib 160 mg d1–21	NCCN category 1	acceptable clinically (7.4 mo, VOYAGER)	active (6.7 mo)	high (HTN 10–24 %, PRES in label, hepatotoxicity)	named as option 08.09.2026
Ripretinib 150 mg	limited resistance evidence	reduced (4.0 mo)	strong (14.2 mo)	low (HTN 5.8 %, no PRES in label)	fourth line; equal to sunitinib in the co-occurring subgroup (HR 1.07)
Sorafenib	NCCN 2A, other tumour type	active in vitro (conflicting)	active	moderate	off-guideline, retrospective data
Avapritinib	resistance flagged	resistant	active	bleeding warning	not suitable
Nilotinib / ponatinib / dasatinib	NCCN 2A other tumour type / no flag	no clinical activity / escape mutation / no data	—	low / arterial risk / low	not recommended
Velzatinib (IDRX-42), StrateGIST-1	—	active (exon 13 ORR 17 %)	active (exon 17/18 ORR 23 %)	selective KIT, low VEGFR	trial (StrateGIST 3, NCT07218926, recruiting in Europe); 2nd line mPFS 13.7 mo, 3rd line 9.2 mo
Bezuclastinib + sunitinib (Peak, phase 3)	—	yes (sunitinib)	yes (bezuclastinib)	sunitinib risk retained	phase 3 Peak positive (PFS 16.5 vs 9.2 mo, HR 0.50); FDA decision due 30.11.2026
Pimitespib (TAS-116, Jeselhy), HSP90 inhibitor	—	mutation-agnostic (KIT degradation; HR 0.52 in exon 13/14 + 17/18 subgroup)	same	low (no VEGFR activity; diarrhoea 76 %, night blindness 12 %)	Japan only; phase 3 mPFS 2.8 vs 1.4 mo, OS HR 0.42, no objective responses; import ≈ €17,700 per 28-day cycle
Local therapy + imatinib	—	ablates clone	ablates clone	none	NCCN-endorsed for limited progression; fresh imaging first

11 Questions for the multidisciplinary tumour board

1. **Physical removal of the new focus, with imatinib continued.** Thermal ablation has been declined because the 42 × 36 mm segment II/III lesion abuts the left portal branch and the left hepatic duct. Which option does the board recommend and where can it be done: resection of the left lateral section (segments II–III), irreversible electroporation (NanoKnife), histotripsy (HistoSonics, newly approved in

Europe), SBRT, or transarterial treatment (DEB-TACE/TARE) of the segment II/III feeding artery — and how should the active rim of the right-lobe mass be handled?

2. **New systemic drug or trial, if and when needed.** Both resistance clones (KIT exon 13 V654A and exon 18 A829P) must be covered; each later TKI line gives a shorter time to the next progression with more toxicity; she had PRES with intracerebral haemorrhage on a VEGFR inhibitor.

Option	Covers V654A / A829P	Expected benefit	Main risk for her	Access / cost
Sunitinib 50 mg 4/2 (prescribed)	yes / no	median PFS 5–15 months by clone	hypertension, VEGFR class — PRES history	available; prescribed, not started
Regorafenib 160 mg d1–21	yes / yes	median PFS about 5 months	hypertension, hand–foot	NCCN category 1; named as option 08.09.2026
Ripretinib 150 mg	weak / strong	4–6 months; equal to sunitinib in the co-mutated subgroup	lowest neurovascular risk of the TKIs	about USD 36,600 per month (Israeli price ₪110,986 per 90 tablets)
Velzatinib (IDRX-42), StrateGIST-1	yes / yes	2nd line 13.7 months, 3rd line 9.2 months	selective KIT, low VEGFR	trial: StrateGIST 3 (NCT07218926, velzatinib vs sunitinib) recruiting incl. Warsaw, Poznań, Kielce, Essen, Berlin, Mannheim, Budapest; 50 % sunitinib arm
Bezuclastinib + sunitinib (Peak)	yes / yes	phase 3: PFS 16.5 vs 9.2 months vs sunitinib, HR 0.50; ORR 46 vs 26 % (ASCO 2026)	sunitinib risk retained	FDA decision due 30.11.2026; not enrolling in Europe
Pimitespib (HSP90 inhibitor)	mutation-agnostic	PFS 2.8 vs 1.4 months, stabilisation only	diarrhoea, night blindness; no VEGFR activity	Japan only; import ≈ €17,700 per cycle

Question: which of these would the board choose, at what point — now, or only after local control fails — and with what blood-pressure safeguards?

3. **Immune- and antibody-based options: is anything worth pursuing now or alongside local treatment?** The evidence is weak and mostly preclinical; the question is whether the board sees a rational entry point.

What has been tried in GIST (OpenEvidence review, sections 13–16):

Approach	Evidence	Result	Relevance to her
Checkpoint inhibitor + TKI (AXAGIST: avelumab + axitinib)	Phase II, n = 56, pretreated GIST	ORR 8.9 %, median PFS 4.6 months; five responders lasted a median 18.5 months	Small chance of durable benefit; grade ≥ 3 toxicity 30 %
Regorafenib + checkpoint inhibitor (IMPRESARC)	Retrospective French cohort	46 % of GIST patients progression-free at 6 months, mostly KIT exon 11	Uncontrolled, but matches her genotype and a drug already proposed

Approach	Evidence	Result	Relevance to her
Imatinib + peginterferon- α 2b	Pilot, 8 treatment-naïve patients	100 % response; never repeated	Depended on imatinib sensitivity — not her situation
Checkpoint monotherapy, mRNA neoantigen vaccine, TIL therapy	No GIST data	—	TMB 3–4.8 Muts/Mb and MSS leave few targets

New leads: (a) KIT-targeted antibody–drug conjugate NN3201 (c-KIT antibody + MMAE, Novelty Nobility), phase 1 NCT06805825, recruiting in the US, imatinib-pretreated GIST eligible (gisttrials.org trial 419); (b) V δ 1 $\gamma\delta$ T cells are the anti-tumour immune population in GIST, depleted in imatinib-resistant tumours and restored by PD-L1 blockade plus imatinib in mice (Zeng, DeMatteo et al., Research Square preprint 2026, PMC13252525); (c) single-cell profiling of imatinib-resistant GIST shows a specific immunosuppressive microenvironment — regulatory T cells engaged via TIGIT–NECTIN2 and IDO1-positive dendritic cells via BTLA–TNFRSF14, absent in sensitive tumours (Liu X et al., Cell Death Dis 2024;15:190, doi:10.1038/s41419-024-06571-3), a rationale for Treg-, TIGIT- or IDO1-directed strategies; (d) intratumoral T-cell activation kills tumours regardless of antigen specificity through a paracrine innate mechanism (Ghirardelli Smith et al., Nat Immunol 2026, doi:10.1038/s41590-026-02642-z) — a rationale for intratumoral immune agents in an accessible liver lesion, without clinical GIST data.

Question: is any of these a candidate for her — the ADC trial, checkpoint inhibitor + regorafenib, an intratumoral or cell-therapy trial — and under what conditions?

12 Appendix A — PET-CT report, KMKOC, 23.09.2026 (English translation)

Drive: [this translation](#) (Ukrainian original)

MUNICIPAL NON-PROFIT ENTERPRISE «KYIV CITY CLINICAL ONCOLOGY CENTRE» (KMKOC)

69 Verkhovynna St., Kyiv, 03115, Ukraine · tel. (044) 4239607 · E-mail: petctkyiv@gmail.com · EDRPOU code 13697965

Combined PET-CT examination protocol

Patient (full name): Zherlitsyna Anastasiia Stanislavivna

Date of birth: 25.04.1976

Medical record No.: 13189/26

Patient weight: 65 kg

Date of examination: 23.09.2026

Scanner: PET-CT GE Discovery STE16

Radiopharmaceutical (RP): 18F-FDG

RP activity: 260 MBq

Route of administration: intravenous

Effective dose, low-dose CT: 5.64 mSv

Effective dose, PET: 4.94 mSv

Total dose, PET/CT: 10.58 mSv

Preparation: fasting; oral water loading.

Imaging performed 80 min after RP injection.

Field of view: from the supraorbital ridges to the mid-third of the thighs

Blood glucose at time of scan: 5.9 mmol/L

Diagnosis:

Leiomyosarcoma of the uterine body, T2NxM1. Secondary involvement of the liver and peritoneum. Disease progression in the liver. Gastric GIST.

Anamnesis morbi:

Histopathology (liver) 2023 – metastasis of uterine leiomyosarcoma; histopathology (stomach) from 2023 – gastric GIST, GIST metastasis to the liver;

Surgical treatment – 01.2023 – repeat liver biopsy; 11.04.2023 – gastric biopsy, core biopsy of the liver;

Chemotherapy – 4 cycles in 2022 + targeted therapy with pazopanib, thereafter 3.5 years of imatinib.

Description of the PET-CT images (comparison with PET/CT of 19.06.2026):

Neck: symmetrical uptake of the tracer in the tonsils is seen, which is a physiological process. No other areas of tracer uptake in the neck organs were found. No cervical, supra-/infraclavicular or axillary lymph nodes with pathological RP accumulation are visualised.

Chest: the parenchyma of the right and left lungs and the mediastinal organs show no pathological hypermetabolism. In the middle–lower third of the oesophagus, increased RP uptake of moderate-to-high intensity is seen, SUVmax = 2.0–3.2, without additional mass lesions. Physiological tracer distribution is seen in the mammary glands. Physiological RP distribution is seen in the myocardium. Mediastinal blood-pool SUVmax = 1.6. No mediastinal or pulmonary-hilar lymph nodes with pathological RP accumulation were found.

Abdominal cavity: the gastric walls show a relatively uniform RP distribution, SUVmax up to 2.8, without suspicious foci of hypermetabolism. Along the lesser curvature an FDG-negative lesion persists, closely apposed to the tail of the pancreas, with signs of calcification, measuring 43×32 mm (previously 4.3×3.3 cm). In the liver, the previously known multiple ametabolic lesions persist (the largest 98×148 mm in the right lobe

– previously 14.4×10.4 mm) with ingrowth of metabolic activity along the contour and within the structure of some of them, intensity SUVmax up to 4.9, with a metabolic size in the left lobe up to 35×54 mm (confluent) and up to 15×24 mm within the ametabolic area in segment S8. Physiological RP distribution is seen in the spleen, pancreas and both adrenal glands. The walls of the small and large intestine show no pathological hypermetabolism. RP uptake in both kidneys is physiological for this tracer. No abdominal or retroperitoneal lymph nodes with pathological RP accumulation were found.

Pelvis: status after chemotherapy for leiomyosarcoma of the uterine body. Along the anterior uterine wall a heterogeneous lesion persists, SUVmax up to 1.8, with calcifications in its structure, measuring 67×49 mm (previously 6.8×5.5 cm). A stable ametabolic nodular lesion persists along the upper margin, measuring 37×47 mm (4.6×3.7 cm). No other areas of pathological RP accumulation in the pelvic organs were found. Physiological tracer uptake is noted in the urinary bladder. No RP hypermetabolism in the lymph nodes of the true pelvis, iliac or inguinal groups was found.

Skeleton: no pathological tracer metabolism indicating osteolytic involvement was found in the bones.

Conclusion:

PET/CT signs of an increase in metabolic activity in multiple foci of liver-parenchyma involvement compared with the PET/CT of 19.06.2026, which is characteristic of progression of the underlying disease.

Stable low-intensity lesion of the uterine body and a stable ametabolic node of the uterine body, without marked change.

A stable FDG-negative lesion of the gastric body persists.

RP uptake in the oesophagus may correspond to an inflammatory origin; however, further evaluation – EGD (upper GI endoscopy) – is recommended.

No other data indicating an FDG-positive pathological process were found.

Recommended:

oncologist consultation; PET/CT treatment monitoring.

Nuclear-medicine physician: Kholodna A.V.

Note! 18F-FDG is not a tumour-specific radiopharmaceutical. The presence of an FDG-insensitive tumour type cannot be excluded. This report cannot serve as a definitive diagnosis. The PET/CT data should be considered valid for three weeks.

English translation of the original Ukrainian PET-CT report of 23.09.2026 (KMKOC, Kyiv). Faithful clinical translation; measurements and SUV values reproduced as in the original.

13 Appendix B — Expert opinion on the FDG PET-CT, 08.09.2026 (English translation)

Drive: [this translation](#) (Russian original)

13.0.1 Expert opinion on FDG PET-CT — English translation

Source (Russian original): docs_eng/2026_new/Zherlitsyna Anastasia - PET CT revision - 08.09.2026.docx (second medical opinion, Israel). Translation made 10.09.2026; wording follows the original as closely as possible.

13.0.1.1 Patient and study details Patient: ZHERLITZINA ANASTASIA / Zherlitsyna Anastasia

ID: ME923960

Date of birth: 25.04.1976

Study date: 19.06.2026

Date of reading: 08.09.2026

13.0.1.2 Indication The patient is under follow-up and receiving treatment for metastatic gastrointestinal stromal tumour (GIST).

Comparison: with the previous study of 13.05.2025.

13.0.1.3 Technique The study was performed after intravenous injection of the radiopharmaceutical, without intravenous and without oral contrast agent.

13.0.1.4 Findings

13.0.1.4.1 Head and neck As is known, the diagnostic capability of this study with respect to the brain is limited. Within the limits of the method, no gross changes of the brain parenchyma are identified.

No hypermetabolic enlarged cervical lymph nodes are identified.

The thyroid gland is of normal size, without pathological FDG uptake.

13.0.1.4.2 Chest Small nodules are seen in both lungs, stable over a long period, with sizes below the resolution of PET.

No pleural effusion.

Minimal amount of pericardial fluid — unchanged.

No pathological FDG uptake in the mediastinal and hilar lymph nodes.

No pathological FDG uptake in the breast tissue.

No pathological uptake in the axillary lymph nodes.

The superior vena cava (SVC) is not visualised — probably occluded. Dilated collateral vessels are seen in the chest.

13.0.1.4.3 Abdomen and pelvis The known hypodense lesions are again identified in both lobes of the liver.

A large lesion in the right lobe of the liver with peripheral FDG uptake, which in the present study is somewhat more pronounced. The size of the lesion is about 15 × 10 cm; on the previous study — about 11–12.5 cm.

The lesion in the left lobe of the liver, about 42 × 36 mm — without significant change in size, however the intensity of FDG uptake has now increased.

The gallbladder is not enlarged. The intrahepatic bile ducts are not dilated. The portal vein is patent.

The known lesion in the stomach wall with calcifications, without significant change in size — about 37 × 31 mm. Minimal FDG uptake is noted, similar to that observed previously, without significant dynamics.

The spleen is of normal size, without pathological FDG uptake.

No pathological FDG uptake is identified in the pancreatic parenchyma.

The right adrenal gland is compressed by the liver, without mass lesions and without pathological FDG uptake.

Thickening of the left adrenal gland without pathological FDG uptake — unchanged.

Kidneys without radiopaque calculi, no hydronephrosis.

No hypermetabolic enlarged lymph nodes are identified in the abdominal cavity, retroperitoneum or pelvis.

No pathological FDG uptake in the inguinal lymph nodes.

Urinary bladder without significant changes.

Bowel loops, as far as can be assessed by this method, without significant changes.

The known heterogeneous lesions of the uterus — without significant dynamics.

13.0.1.4.4 Skeletal system Within the examined area no pathological foci of FDG uptake or destructive bone changes are identified.

13.0.1.5 Conclusion The known gastric lesion — without significant change in size and without significant FDG uptake.

Metabolic progression of the liver lesions, especially of the lesion in the left lobe, as well as enlargement of the large lesion in the right lobe of the liver, whose metabolic activity has also somewhat increased.

Small pulmonary nodules — unchanged.

13.0.1.6 Signature Dr. Victoria Kulikova

Nuclear medicine specialist

Physician licence No. 84847

Specialist licence No. 131252

Translator's note: the report states the CT was performed without intravenous contrast; the DICOM series of the 19.06.2026 study include a venous-phase contrast series (series 9), so this statement appears to refer to the PET reading protocol rather than to the CT acquisition.

14 Appendix C — LISOD retrospective radiology report, 08.07.2026 (English translation)

Drive: [this translation](#) (Ukrainian original)

14.0.1 LISOD Radiology Report — Retrospective PET-CT Analysis (English Translation)

Source (Ukrainian original): docs_eng/anastasia_disks/June_19_2026_PET_CT/Жерліцина_Анастасія_Станіславівна_F

Facility: LISOD, Radiology Diagnostics Department

Patient: Zherlitsyna Anastasiia Stanislavivna, ID 110889, DOB 25.04.1976

Report date: 08.07.2026

Study: retrospective analysis of provided imaging studies (PET/CT with FDG from 19.06.2026), compared to prior PET/CT with FDG from 13.05.2025 and 14.03.2024

Reporting physician: Novikov M.Ye.

14.0.1.1 Clinical background (as stated in report) ICD-10: C16.9 - Malignant neoplasm of stomach, unspecified. Diagnosis: Leiomyosarcoma of uterine body, T2NxM1, secondary liver and peritoneal involvement. Status after 4 cycles of Gemcitabine + Docetaxel + Avastin chemotherapy with liver progression (per CT/MRI 10/2022). C-KIT, MET mutation detected (ONCOMINE). Status after Pazopanib treatment (05-12/01/2023) with PRES development and positive CT dynamics. 01/2023 - repeat liver biopsy, histology: uterine leiomyosarcoma metastasis. Status after 3 cycles of Trabectedin with negative radiological dynamics (PET/CT 11/04/2023). Suspected synchronous gastric neoplasia (GIST?). Status after EUS-FNA gastric biopsy, liver core biopsy - histologically GIST of stomach, GIST metastasis to liver (CSD). 08/2023 - status after 4 months Imatinib with pronounced positive PET/CT response. 03/2025 - status after 2 years Imatinib, positive clinical picture. 05/2025 - stable radiological dynamics on Imatinib.

14.0.1.2 Findings Head and neck: No changes characteristic of metabolically active aggressive neoplastic involvement, same as prior imaging. No significant change since 13/05/2025, no new active or suspicious lesions.

Chest: No changes characteristic of metabolically active aggressive neoplastic involvement, same as prior imaging. No significant change since 13/05/2025, no new active or suspicious lesions. No excess pleural/pericardial fluid, no large vessel thrombi.

Abdomen and pelvis: Known foci and masses in the liver are generally stable in size overall; however, mural (wall) components within them — particularly in the largest lesion in the right lobe — show new nodular elements with notable enhancement and partial focal moderately increased activity, moderately higher than the hepatic blood pool. This is best seen in nodule "42" in the left lobe, with uptake up to SUVmax 3.2 (adjacent parenchyma SUV 2).

No new separate similar active or suspicious nodules found.

Known residual gastric tumor without significant activity; nodule above the bladder/uterus on the right also without significant activity and without change.

No new separate active or similar nodules, no new suspicious lymphadenopathy found.

No signs of bowel obstruction, ascites, hydronephrosis, bile duct dilation, or large vessel thrombi.

Bones and soft tissue: New elongated/geographic lucency in the femoral head of the right hip, approximately 31mm, with a sclerotic rim and moderate activity, currently without flattening/deformity of the head. No other new active/suspicious bone lesions found.

14.0.1.3 Radiological conclusion New mural (wall) components with moderate activity within the liver lesions, raising suspicion of return of viable tumor elements of GIST metastases.

Physician: Novikov M.Ye.

[Translation note: plain-language English translation of the Ukrainian original for reference purposes; not a certified medical translation.]

15 Appendix D — PET-CT report, Feofaniya, 19.06.2026 (English translation)

Drive: [this translation](#) (Ukrainian original) · DICOM disc

15.0.1 PET-CT Whole Body with Contrast Enhancement — English Translation

Source (Ukrainian original): docs_eng/anastasia_disks/June_19_2026_PET_CT/Жерліцина_Анастасія_Станіславівна_f

Facility: All-Ukrainian Radiosurgery Center (Feofaniya), PET-CT Diagnostics Department

Patient: Zherlitsyna Anastasiia Stanislavivna, b. 1976

Study: Whole-body PET-CT with contrast enhancement

Date: 2026-06-19 / #53478/25

F18-FDG injected IV: 232.7 MBq; PET effective dose 4.4 mSv, CT 18.7 mSv

15.0.1.1 Clinical data 1. Gastrointestinal stromal tumor (GIST) of the stomach, GIST metastasis to the liver (CSD). Status post EUS-FNA biopsy of the stomach, core biopsy of the liver. 08.2023 - status after 4 months of Imatinib treatment. 03.2025 - status after 2 years of Imatinib treatment with positive clinical picture. Daily Imatinib (Glivec) continued after the last PET-CT.

2. Leiomyosarcoma of the uterine body T2NxM1, secondary involvement of liver and peritoneum. Status after 4 courses of chemotherapy per Gemcitabine+Docetaxel+Avastin protocol with disease progression in the liver (10.2022). C-KIT and MET mutations detected. Status after Pazopanib treatment (05-12.01.2023) with development of posterior reversible encephalopathy syndrome and positive CT dynamics. 01.2023 repeat liver biopsy: histology - metastasis of uterine leiomyosarcoma. Status after 3 cycles of chemotherapy per Trabectedin protocol with negative dynamics (PET-CT from 11.04.2023).

Reference PET-CT for comparison: 13.05.2025

Reference values: mediastinal blood pool SUVmax 1.41; liver parenchyma pool SUVmax 1.99.

15.0.1.2 Findings Oncological disease control:

Stomach: not distended; along the lesser curvature a hypodense ametabolic mass persists, stable in size at up to 4.3×3.3×4.4 cm (ax+sag), with increasing calcifications in its structure.

Distant metastatic spread:

Liver: compensatorily enlarged, right lobe craniocaudal size up to 18.7 cm (13.05.2025 - up to 17.9 cm).

Parenchyma contains ametabolic, hypodense masses, stable in size:

- Sg5/6/7/8: massive mass up to 14.4×10.4×14.1 cm (ax+cor)
- Sg7: satellite lesion up to 1.0×1.0 cm
- Sg7/8: subcapsular, up to 6.3×4.3 cm
- Sg5/8 border: small lesion up to 0.6×0.6 cm
- Sg4a/7 border: up to 0.6×0.6 cm
- Sg2/3 border: mass along the left portal vein, 4.5×3.6 cm
- Sg1: up to 1.3×1.3 cm
- Sg4b: small lesion up to 0.4×0.4 cm

Uterus: along the anterior wall, a heterogeneous round mass with calcifications, conditionally stable in size at up to 6.8×5.5×4.8 cm (ax+sag) - per history, likely treated leiomyosarcoma. Along the upper margin, an ametabolic mass of stable size up to 4.6×3.7 cm - pelvic MRI with contrast recommended for correct assessment. Ovaries not clearly differentiated - correlate with ultrasound data.

Bones unremarkable, no suspicious bone marrow FDG uptake.

Neck: physiological symmetric FDG metabolism in the palatine tonsils. No signs of metabolically active cervical lymphadenopathy. Thyroid not enlarged, normal structure, no increased FDG uptake. A subcutaneous port system is placed at the right subclavian region; catheter passes into the superior vena cava, distal tip at the lower third of the SVC.

Chest: mammary glands without pathological FDG uptake. No signs of metabolically active axillary, mediastinal or hilar lymphadenopathy. Trachea and main bronchi patent. Lung parenchyma normal aeration, no infiltrative or mass changes. Single small bullae on the right in S1, subpleural, up to 0.8 cm. No pleural effusion.

Abdomen/retroperitoneum: gallbladder not enlarged, compressed by adjacent liver parenchyma, contents of homogeneous density, no radiopaque calculi. Pancreas normotrophic, contour clear and scalloped. Adrenal glands not enlarged, normal shape. Kidneys typically positioned, parenchyma normal thickness. Collecting systems and ureters normal configuration. No metabolically active intra-abdominal or retroperitoneal lymphadenopathy.

Pelvis: bladder moderately filled, walls not thickened, no additional masses. Small amount of fluid in the Pouch of Douglas. No metabolically active pelvic or inguinal lymphadenopathy.

15.0.1.3 Conclusion Compared with the PET-CT from 13.05.2025 (note: report references this as 13.05.2026, likely a typographical date error for 13.05.2025), the picture of the known gastrointestinal stromal tumor (GIST) of the stomach and metastatic liver involvement shows no reliable negative dynamics.

Given the histologically confirmed GIST, which may show absent or variable FDG avidity, the absence of pronounced metabolic activity does not exclude residual tumor tissue. For correct assessment of disease course and further monitoring, correlation with contrast-enhanced CT or MRI is recommended.

The calcified uterine body mass is stable over time; the ametabolic mass along the upper margin of the uterine body - pelvic MRI with contrast is advisable for correct assessment.

Radiologist: Zakrzhevskya K.P.

Head of the All-Ukrainian Radiosurgery Center, radiologist: Kmetiuk Ya.V.

[Translation note: this is a plain-language English translation of the Ukrainian original for reference purposes; it is not a certified medical translation.]

16 Appendix E — Slice-by-slice verification of the lesion outline




Figure 25: Every 2 mm slice through the lesion: cyan = CT lesion outline, yellow = FDG-active tumour, blue = portal vein, green = other vessels.