

PROTOCOL COMPUTER TOMOGRAPHY DESCRIPTION

Patient's name: Zherlitsyna Anastasia Stanislavivna,
46 years old

Type of research: CT

Phone number: 0685975246

Use of contrast agent: +C

Patient's ID: 7140773

Document No. 28689

Outpatient card No. 28689 dated 12.12.2022

Department: Polyclinic department

from **01/04/2023 15:40:00**

Radiation load: 2737 Total DLP mGy*cm

Doctor: Shevchenko Volodymyr Serhiiovych

Protocol:

The middle structures are not displaced. The subarachnoid spaces are expanded. Post-contrast scans did not reveal pathological foci in the substance of the brain and cerebellum. The ventricles of the brain are not expanded. The craniovertebral region is unremarkable. There are no pathological changes in the pituitary gland. Orbits without features. The paranasal sinuses are pneumatized. Focal infiltrative changes in the lungs were not detected, multiple small thin-walled ones were present throughout. Mediastinal lymph nodes are not enlarged. The trachea and main bronchi are permeable. Pulmonary embolism was not detected. No free fluid was detected in the pleural cavities and the pericardial cavity. No CT-visible pathology was detected in the mammary glands. Supra-, subclavian, axillary lymph nodes are not enlarged. The craniocaudal size of the liver is 242 mm, bilobular single massive heterogeneous foci with central zones of reduced density, size S1 15 mm, S2/3 85x65 mm, S8 subcapsular 100x70 mm, S5,6,7 200x130x180 mm almost completely represented by the central cystic component, subcapsular in S8 25x13 mm. Intrahepatic bile ducts are minimally dilated in the left lobe at S7.8 to 2-4 mm. Portal vein without thrombosis. Gallbladder: does not contain radiopaque calculi, common bile duct is not dilated. The pancreas is displaced to the left by an enlarged liver, the dimensions are typical, the parenchyma is without focal changes, the duct is not dilated. Kidneys, pelvicalyceal system, ureters without features, the right kidney is displaced by an enlarged liver to the level of the wing of the ilium. The concentration and excretory functions are not disturbed. Adrenal glands without features. The spleen is not enlarged, the structure is homogeneous. Lymph nodes of the abdominal cavity, retroperitoneal space, pelvic localization, inguinal are not enlarged. Ascites was not detected. Stomach, intestinal loops without CT-visible pathology. Interloop in the left mesogastric implant 65x50 mm of soft tissue density, closely adjacent to the walls of the body of the stomach in the lower third and the capsule of the tail of the pancreas.

The body of the uterus is 45x42 mm, the myometrium is heterogeneous, in the area of the bottom, a heterogeneous formation with tuberos contours up to 75x55 mm protrudes beyond the contours. The cervix without CT-visible pathology. Ovaries: in the projection of the right, a three-dimensional formation with clear, even contours, homogeneous, with a density of up to +45 H units, dimensions of 75x45 mm, the left one is structurally unchanged, with dimensions of up to 30x15 mm. Bladder: the contents are homogeneous, the wall is not thickened. There are no destructive changes in the bone window, no degenerative changes, no visible areas of lytic destruction within the limits of the method.

Area research:

The brain, organs of the chest, abdominal cavity and pelvis.

Conclusion:

Leyomiosarcoma of the uterus, condition after 4 cycles of chemotherapy. CT signs of secondary damage to the liver parenchyma, interloop implant of the left mesogastric, bulky formation in the projection of the right ovary (implant in the Douglas space?), minimal signs of portal hypertension. From December 1, 2022, a CT picture without visible dynamics.

Recommendations:

PROVIDE PREVIOUS CT (CD) DISKS AT EACH SUBSEQUENT EXAMINATION!!!

Radiologist

/ Shevchenko Volodymyr Serhiiovych /