

Helical Polymerization as Dissipative Geometry:

A Unified Framework for Chromosome Architecture, Mutational Selection,

and the Wet Rock Origin of Life

Version 7 — May 2026

Hernán Diego Herrero

Iniciativa de Investigación Hijos del Sol y la Basura · Córdoba, Argentina

Submitted to: *Journal of Theoretical Biology* · May 2026

Abstract

Background: The Helical Polymerization model proposes that biological macromolecules arise as dissipative geometric attractors in polar aqueous media under sustained directional electromagnetic energy input. Version 7 integrates three converging lines of evidence: a formal variational derivation of the helix as the energy-minimizing geometry under phase-coherence constraints; the first empirical test of a central model prediction against 20,052 independent human mutations; and a computational phase map that, combined with reaction thermodynamics, predicts the precise physical conditions for helical polymerization. The intersection of these three lines generates a novel prediction about the origin of life: the optimal prebiotic environment is not a hydrothermal vent, not a primordial ocean, but a thin film of water on a mineral surface under direct solar UV — *a wet rock in sunlight*.

Methods: Antinode positions for TP53 ($L = 1,179$ codons) were computed for the first 12 normal modes of an open-ended chain and tested against four IARC TP53 Database datasets: somatic mutations ($n = 15,878$), pathogenic germline variants ($n = 4,174$), germline prevalence estimates ($n = 113$), and germline reference entries ($n = 433$). For each dataset, mean distance to the nearest antinode was compared against 10,000 permutation null distributions. Monte Carlo simulations ($N = 10,000$ chains, 800 Langevin steps) mapped helical formation probability P_h across the $J \times kT$ parameter space under directional UV versus isotropic thermal fields.

Results: Somatic TP53 hotspots concentrate at predicted antinode positions ($Z = -8.59$; $p < 0.0001$). Pathogenic germline variants show a stronger signal ($Z = -14.92$; $p < 0.0001$). Non-selected germline datasets show no enrichment ($p > 0.60$), producing the double dissociation specifically predicted by the structural selection hypothesis. The phase map reveals that directional UV favors helical formation only in the low-temperature, moderate-coupling region. The exothermicity of polymerization creates negative thermal feedback, generating discrete pulses of helical formation — a dissipative oscillator whose frequency is determined by the thermal capacity of the interfacial water film and the diurnal UV cycle.

Conclusions: The geometry of the linear chromosome constrains the spatial distribution of functionally disruptive mutations across evolutionary timescales. The prebiotic environment that emerges from the phase map is the thin aqueous film on mineral surfaces under direct UV radiation. Life did not originate in the depths of the ocean. It originated on a wet rock in the sun.

Keywords: *helical polymerization · dissipative attractor · antinode hypothesis · TP53 · structural selection · wet rock origin · standing waves · phase map · prebiotic chemistry · geometric biology*

EPISTEMOLOGICAL NOTE

This framework is explicitly phenomenological. It does not claim that chromosomes are literal electromagnetic antennae in the engineering sense. The claim is that certain geometric stability classes — dissipative attractors, resonant configurations, standing-wave antinodes — recur across physical systems at different scales, and that biological macromolecules are examples of this general class. The framework is organized in four layers of decreasing empirical support, with falsification criteria for each consolidated in Section 9.

The TP53 analysis (Layer 2) is the first completed empirical test. It passed. All other tests remain to be performed.

1. INTRODUCTION

Three facts about biological macromolecules resist easy explanation. First: the helix appears as the stable geometry across wildly different molecular systems — DNA, RNA, proteins, collagen — at different scales, across all domains of life. The chemical explanations for each system are specific and well-established; what remains unexplained is why the solution is always helical. Second: somatic mutation hotspots in tumor suppressor genes are not randomly distributed, and their spatial concentration is not fully explained by intrinsic sequence mutability. Third: the origin of life lacks a satisfying physical account of how polymers of sufficient length and regularity could form spontaneously under prebiotic conditions.

The Helical Polymerization framework proposes a common physical explanation for all three. The helix is the dissipative attractor of polar polymer systems under directional energy input. The antinode structure of linear chromosomes constrains the positions of maximum mechanical stress and therefore of maximum selective enrichment under functional pressure. And the conditions required for the whip-crack transition that produces stable helical oligomers are precisely those of a thin aqueous film on mineral surfaces under solar UV.

1.1 What Version 7 establishes

Version 7 formalizes the helix as the solution to a variational problem — not an intuitive analogy but a mathematical necessity under phase-coherence constraints. It presents the TP53 antinode analysis as the first quantitative test of a model prediction against independent data, with a statistically unambiguous result and an internal control. And it derives from the intersection of the computational phase map and reaction thermodynamics a specific and novel prediction about the prebiotic

environment that was not assumed and was not sought. It emerged from the data.

2. LAYER 1: THE HELIX AS DISSIPATIVE ATTRACTOR

2.1 The variational derivation

Consider a polar polymer chain growing in an aqueous medium under sustained directional electromagnetic energy input. Three constraints apply simultaneously: continuity (each monomer is bonded to the previous one), phase coherence (the energy dissipation must be spatially coherent along the chain), and thermodynamic stability in the dissipative steady state. Phase coherence is decisive: the condition that successive monomers maintain a constant phase relationship with the incident field is precisely the geometric definition of a helix.

Minimizing dissipated energy per unit chain length subject to these constraints yields:

$$E[\theta(s)] = \int_0^L [\frac{1}{2} \kappa (\theta')^2 - J \cos(\theta - \theta_0) - \eta \cdot \theta' \cdot E_{UV}(s)] ds$$

where $\theta(s)$ is the torsion angle at arc-length s , κ is torsional stiffness, J is the dipolar coupling constant, θ_0 is the optimal helicoidal angle ($\approx 32.7^\circ$), η is the UV coupling coefficient, and $E_{UV}(s)$ is the incident field. The stationary solution at $\theta(s) = \theta_0$ — the helical configuration — is a stable attractor: perturbations decay exponentially with rate J/γ .

2.2 The whip-crack bifurcation

The attractor framework predicts a saddle-node bifurcation at a critical chain length λ_c . Below λ_c , dissolution dominates. Above λ_c , the coherent field of the existing chain stabilizes incoming monomers — autocatalysis. The whip-crack is the inevitable consequence of crossing λ_c under any conditions within Region III of the phase map:

$$dL/dt = k_{on}(L) \cdot [M] - k_{off}(L) \text{ where } k_{on}(L) \propto \exp(J \cdot L / (kT \cdot \lambda_c)) \text{ for } L > \lambda_c$$

2.3 The phase map

The parameter space is partitioned into four regions by the bifurcation manifold $P_h(J, kT) = 0.5$: Region I (dissolution), Region II (sub-threshold growth), Region III (whip-crack autocatalysis, requires directional UV), Region IV (chaotic fragmentation).

3. LAYER 1: THE CIRCULAR/LINEAR DISTINCTION AS TOPOLOGICAL CONSEQUENCE

3.1 Antinode positions of open and closed chains

For an open-ended chain of length L , the free-end boundary conditions require maximum amplitude at both termini. The antinode positions of mode n are:

$$x_k = (2k-1) \cdot L / (2n), k = 1, 2, \dots, n$$

A closed chain has no fixed antinode positions — the energy distribution of any mode is uniform by rotational symmetry. This is a mathematical consequence of boundary conditions, not a biological

observation.

3.2 Physical origin of the distinction

Isotropic thermal field → circular topology: In a hydrothermal vent, energy arrives from all directions. No elongation direction is privileged. The polymer closes on itself — circular chromosome, no fixed antinodes, no meiotic crossover positions. This is the archaeal solution.

Anisotropic UV field → linear topology: At a mineral-water surface under solar UV, energy arrives perpendicular to the interface. The field creates a preferred elongation direction, producing a linear chain with free ends, fixed antinode positions, and structurally distinguished sites of maximum amplitude. This is the bacterial and eukaryotic solution.

3.3 Biological consequences

The topological distinction generates four consequences derivable without evolutionary narrative: geometric impossibility of meiosis in archaea (no fixed crossover positions), physical necessity of telomeres (obligatory end-antinodes require protection), concentration of functional hotspots at interior antinodes (tested in Section 4), and greater archaeal stability under isotropic perturbation but vulnerability to acute anisotropic stress.

4. LAYER 2: EMPIRICAL VALIDATION — TP53 ANTINODE ANALYSIS

4.1 Prediction P-A

The interior antinode positions of a linear chromosome should be enriched among variants preferentially selected for severe functional disruption, because these positions accumulate the greatest torsional stress. The double dissociation — selected datasets significant, non-selected not significant — is the specific falsifiable form of this prediction.

4.2 Results

Dataset	n	d_obs	d_null	Z	p-value	Result
Somatic (MutationView)	15,878	6.85	7.40 ± 0.77	−8.59	< 0.0001	✓ CONFIRMED
Pathogenic germline	4,174	5.56	7.40 ± 0.38	−14.92	< 0.0001	✓ CONFIRMED
Germline prevalence	113	11.12	7.40 ± 0.77	+4.82	1.000	✗ NOT SIG.
Germline reference	433	7.50	7.40 ± 0.38	+0.26	0.610	✗ NOT SIG.

Table 1. Antinode analysis across four TP53 datasets. d_obs = observed mean distance to nearest antinode (codons); d_null = null distribution mean ± SD (10,000 permutations). Green = functionally selected datasets; orange = non-selected controls. The double dissociation is the specific prediction of the structural selection hypothesis.

Somatic hotspots show mean distance 6.85 codons versus a null of 7.40 (Z = −8.59; p < 0.0001). The four highest-frequency hotspots — codons 248, 273, 175, 245 — fall within ±5 codons of predicted antinodes. 82% of top somatic hotspots (90th percentile) fall within the ±8 codon antinode window. Pathogenic germline variants show a stronger signal (Z = −14.92), consistent with more stringent

functional selection.



Figure 1. TP53 somatic mutation hotspots vs predicted antinode positions. Upper: distribution of 15,878 somatic mutations (blue) with antinode positions (gold vertical lines). Green shading marks hotspots within the ± 8 codon window. Lower panels: permutation test distributions — observed values (red lines) lie far outside the null distributions in both selected datasets.



Figure 2. Comparative analysis: somatic vs germline datasets. The double dissociation between functionally selected datasets (significant antinode enrichment, green) and non-selected datasets (no enrichment, orange) is the specific prediction of the structural selection hypothesis. The stronger germline signal ($Z = -14.92$) reflects more stringent functional selection.

4.3 Interpretation

The result supports the structural selection hypothesis in its pre-specified form. The double dissociation rules out the alternative that antinodes happen to overlap with intrinsically mutable sequences independently of selection. The result does not prove the literal electromagnetic mechanism proposed by the model; it proves that the geometric prediction is correct. The mechanism remains to be determined. These are separate claims and are evaluated separately.

5. LAYER 1: COMPUTATIONAL PHASE MAP — MONTE CARLO SIMULATIONS

Monte Carlo simulations ($N = 10,000$ chains, 800 Langevin steps) mapped $P_h(J, kT)$ under directional UV versus isotropic thermal fields. The phase map reveals that directional UV favors helical formation only in the low-temperature, moderate-coupling region ($kT < 0.15, J \in [0.2, 0.5]$). Outside this window, the isotropic field produces comparable or higher P_h — consistent with the model's prediction of two distinct physical origin regimes for circular and linear topologies.




Figure 3. Monte Carlo phase map. Upper left: P_h vs chain length N under nominal reactor parameters. Center: $P_h(J, kT)$ maps for anisotropic UV (green) and isotropic thermal (blue) conditions. Right center: differential map $\Delta P_h = P_h(\text{aniso}) - P_h(\text{iso})$ — green regions where UV directional field favors helix. The star marks the nominal Córdoba reactor conditions. Lower: temporal dynamics $P_h(t)$.

5.1 The thermal feedback discovery

The phase map reveals an unexpected constraint: helical formation under directional UV requires low temperature ($kT < 0.15$). But polymerization is exothermic — it releases heat. This creates a negative thermal feedback loop:

- Interface cools to $kT < 0.15$ through evaporative cooling and nocturnal radiation.
- UV activation initiates helical polymerization in the directional field.
- Exothermic reaction raises local temperature above the kT threshold.
- Polymerization slows as the system exits the Region III window.
- Interface cools again, restoring conditions for the next pulse.

This produces discrete pulses of helical formation — a dissipative oscillator. Each pulse generates a population of oligomers of characteristic length. Variation between pulses produces variation in products — the basis of selection without copying.

6. LAYER 2–3: THE WET ROCK ORIGIN OF LIFE

6.1 What the phase map predicts

The computational phase map defines the prebiotic environment satisfying Region III conditions with natural thermal regulation:

- **Low thermal mass:** thin water film (micrometers to millimeters) with low heat capacity and rapid thermal cycling — the natural regulator of the oscillator frequency.
- **Directional UV:** perpendicular solar radiation at the mineral surface — the anisotropic condition for linear topology.
- **pH gradient:** mineral alkalinity plus dissolved CO₂ acidity — exists at the mineral-water interface without engineering.
- **Precursor concentration:** evaporative accumulation of polar monomers in the film microtopography.
- **Diurnal oscillator:** the daily UV cycle as the clock for pulsatile polymerization and nocturnal cooling for system reset.

This is not a hydrothermal vent. It is not a primordial ocean. It is a wet rock in the sun.

6.2 The moss on a stone

The physical description above has a biological correlate that is not coincidental. Mosses grow in exactly these conditions: thin water films on mineral surfaces under direct solar radiation, tolerating complete desiccation and recovering with water. Cryptobiosis — the capacity to suspend all metabolic activity during desiccation and recover — is not an evolutionary adaptation to a hostile environment. It is the signature of an origin in a pulsatile environment where hydration and dehydration alternated on a diurnal cycle. The first biopolymers were not adapted to tolerate desiccation. They were produced by a mechanism that operated in those cycles.

6.3 Experimental predictions for the Córdoba reactor

- **Temperature dependence:** reducing reactor temperature from 25°C to 10°C should increase helical content measurable by circular dichroism spectroscopy.
- **UV directionality:** replacing collimated with diffuse UV (equal total intensity) should reduce helical content and eliminate the $P_h(\text{aniso}) > P_h(\text{iso})$ differential.
- **Pulsatile dynamics:** ThT fluorescence measured continuously should show discrete peaks rather than monotonic increase — each peak a polymerization pulse followed by thermal reset.

7. COMPUTATIONAL VISUALIZATION: 3D POLYMER DYNAMICS

A 3D molecular dynamics simulation of an 80-monomer chain under Langevin dynamics with explicit torsional potential, bending rigidity, and directional UV coupling shows convergence from a disordered initial state toward periodic oscillatory geometry. The X-Y projection shows emergence of helical oscillation along the elongation axis. The local torsional energy profile shows concentration at theoretically predicted antinode positions, and the chain termini show systematically

elevated energy consistent with obligatory end-antinodes in open-ended chains.




Figure 4. 3D polymer dynamics simulation. Left: final configuration of the 80-monomer chain after 15,000 simulation steps. Color encodes local torsional energy — yellow = high (antinode positions), purple = low. Red dots mark theoretically predicted antinode positions; white dots mark chain termini. Upper right: $P_h(t)$ showing convergence from disordered initial conditions. Lower right: local torsional energy profile with predicted antinodes marked.

The simulation reaches $P_h = 0.31$ — a partially ordered sub-threshold state consistent with the phase map at $J = 0.5$, $kT = 0.03$. Full convergence to $P_h > 0.5$ requires parameters in the green region of the differential phase map. These simulations will be extended using LAMMPS-based molecular dynamics with explicit solvent.

8. LAYER 3: BIOLOGICAL DERIVATIONS

8.1 Amitosis as bifurcation event

Cell division in the framework is a physical consequence of the same dynamics that produced the chromosome. As the helical chain grows beyond a critical length, accumulated torsional energy is released as two simultaneous propagating waves from the central compression node. The daughter segments are products of the same bifurcation with stochastic variation — heritable variation without a copying mechanism.

8.2 The circular chromosome prediction

The most directly testable prediction of the topological distinction: archaea, with circular chromosomes and no fixed antinode positions, should show no significant antinode enrichment of functionally selected mutations. If the TP53 double dissociation reflects the geometry of linear chromosomes specifically, then equivalent analysis in archaeal genes should produce a null result regardless of selection pressure. This is the next critical test of the framework.

9. FALSIFICATION CRITERIA

Layer	Prediction	Falsification criterion
1 — Math	Helix minimizes dissipated energy per unit length under identical boundary conditions	Non-helical geometry with trivial dissipation under identical boundary conditions
1 — Computation	$P_h(\text{aniso}) > P_h(\text{iso})$ in low-kT, moderate-J region	$P_h(\text{aniso}) \leq P_h(\text{iso})$ across the full intermediate parameter range
2 — TP53 ✓	Double dissociation: selected significant, non-selected not significant	Selected not significant, non-selected significant — CONFIRMED 0.05 — NOT OBSERVED
2 — Replication	Same double dissociation in BRCA1 (L=1,863) and BRCA2 (L=2,201)	Antinode enrichment absent in one of the replication tumor suppressors
3 — Archaea	Circular chromosomes show NO antinode enrichment of functionally selected mutations	Significant antinode enrichment in archaea comparable to linear-chromosomes
3 — Reactor	Pulsatile ThT; temperature dependence; UV directionality effect	No effect ThT; no temperature effect 10–25°C; no UV directionality effect
4 — Prebiotic	Helical oligomers on mineral surfaces under directional UV	No helical oligomers under any mineral-surface UV condition

Table 2. Falsification criteria by model layer. Green row = completed test with confirmed result.

10. DISCUSSION

10.1 The convergence of three independent lines

The mathematical derivation, the TP53 empirical analysis, and the computational phase map were developed independently and converged on the same framework. The TP53 data predates the model's formalization. The phase map was computed to characterize the simulator, not to predict the prebiotic environment. The wet rock prediction emerged from the data, not from assumption. This convergence is evidence, not circular reasoning.

10.2 Strengths and limitations

- Strongest:** The TP53 double dissociation. $Z = -14.92$ in 4,174 pathogenic germline variants with internal null control. Among the strongest statistical signals in computational genomics.
- Strong:** The variational derivation. Physically sound, requires only standard variational calculus and polymer physics.
- Moderate:** The thermal feedback mechanism and wet rock prediction. Physically coherent but require experimental validation at the Córdoba reactor.
- Weak:** Layer 4 extensions. Interesting and falsifiable; included as research program, not evidence.

10.3 Next steps

Three immediate priorities: (1) Replication of the TP53 analysis in BRCA1 and RB1 to establish the antinode enrichment as a general principle. (2) The three Córdoba reactor experiments — temperature dependence, UV directionality, and pulsatile ThT dynamics. (3) LAMMPS molecular dynamics on explicit mineral surfaces to test helical oligomer formation under predicted prebiotic conditions.

11. CONCLUSIONS

The helix is not a chemically specific solution to molecular structure. It is a physically general one — the dissipative attractor of polar polymer systems under sustained directional energy input. The mathematical derivation establishes this from first principles. The TP53 analysis demonstrates that its geometric consequences — the antinode structure of linear chromosomes — are observable in the distribution of 20,052 real mutations with high statistical significance and a matched internal control. The computational phase map shows that the conditions required for its spontaneous formation are precisely those of a wet mineral surface under solar UV.

The first polymers did not form in the darkness of a hydrothermal vent. They formed in pulses, on the surface of a rock, under the sun, in a thin film of water that cooled by night and was recharged by rain. The evidence that the distribution of cancer mutations in TP53 carries the geometric signature of this process — that the same standing-wave physics that governed the formation of the first polymers still constrains the spatial distribution of the most functionally disruptive mutations three billion years later — is either a remarkable coincidence or a fundamental principle.

The data suggest it is a principle.

DATA AND CODE AVAILABILITY

All mutation data: IARC TP53 Database release 21 (<https://tp53.isb-cgc.org>). Analysis scripts: `tp53_antinodos.py`, `tp53_germline.py`, `helix_montecarlo_local.py`, `v7_helix_3d.py`. Available from the corresponding author. Python 3.11, NumPy, SciPy, pandas, matplotlib.

REFERENCES

- Alexandrov, L. B., et al. (2013). Signatures of mutational processes in human cancer. *Nature*, 500, 415–421.
- Bouaoun, L., et al. (2016). TP53 variations in human cancers: new lessons from the IARC TP53 database. *Human Mutation*, 37, 865–876.
- Chialvo, D. R. (2010). Emergent complex neural dynamics. *Nature Physics*, 6, 744–750.
- Friston, K. (2010). The free-energy principle: A unified brain theory? *Nature Reviews Neuroscience*, 11, 127–138.
- Greenman, C., et al. (2007). Patterns of somatic mutation in human cancer genomes. *Nature*, 446, 153–158.

- Herrero, H. D. (2026a). Helical Polymerization as the Physical Basis of Life, v6. Preprint, *Origins of Life and Evolution of Biospheres*. Hijos del Sol y la Basura, Córdoba.
- Herrero, H. D. (2026b). Número de Reynolds Psicológico (Re_{ps}), v1.5. Hijos del Sol y la Basura, Córdoba.
- Herrero, H. D. (2026c). Sistemas Sociales Disipativos y Economía Disipativa. Preprint, Zenodo. Córdoba.
- IARC TP53 Database. Release 21. International Agency for Research on Cancer. <https://tp53.isb-cgc.org>. Accessed May 2026.
- Petitjean, A., et al. (2007). Impact of mutant p53 functional properties on TP53 mutation patterns and tumor phenotype. *Human Mutation*, 28, 622–629.
- Prigogine, I. (1984). *Order Out of Chaos*. Bantam Books, New York.
- Xie, L., et al. (2013). Sleep drives metabolite clearance from the adult brain. *Science*, 342, 373–377.

APPENDIX A: ANTINODE DERIVATION

For a chain of length L with free-end boundary conditions, the standing-wave equation with $\partial u / \partial x = 0$ at $x = 0$ and $x = L$ has spatial eigenfunctions $X_n(x) = \cos((2n-1)\pi x / 2L)$. The antinodes of mode n — positions where $|X_n(x)| = 1$ — are:

$$x_k = (2k-1) \cdot L / (2n), k = 1, 2, \dots, n$$

For TP53 ($L = 1,179$, $n = 1-12$): 63 unique antinode positions covering 10.7% of the gene within ± 8 codons. Significant somatic enrichment ($p < 0.001$) obtained for $n = 6$ through $n = 20$ in sensitivity analysis.

APPENDIX B: LANGEVIN SIMULATION

Each torsion angle θ_i is updated at each step:

$$\theta_i(t+dt) = \theta_i(t) + \gamma [-J \cdot \sin(\theta_i - \theta_{i-1}) + F_{UV}(\theta_i)] \cdot dt + \sqrt{(2kT \cdot dt)} \cdot \xi_i$$

$\gamma = 0.80$, $dt = 0.10$, $\theta_i = \pi / 5.5 \approx 32.7^\circ$. Anisotropic UV: $F_{UV} = I_{UV} \cdot \alpha \cdot \sin(\theta_i)$, $\alpha = 0.38$. Isotropic: $F_{UV} = I_{UV} \cdot \alpha \cdot \xi$ (white noise). Helical classification: $>55\%$ of angles within ± 0.6 rad of θ_i . $N = 10,000$ chains, 800 steps. All code available on request.