

A COINTUITIONISTIC ADJOINT LOGIC

HARLEY EADES III AND GIANLUIGI BELLIN

e-mail address: heades@augusta.edu

Computer Science, Augusta University, Augusta, GA

e-mail address: gianluigi.bellin@univr.it

Dipartimento di Informatica, Università di Verona, Strada Le Grazie, 37134 Verona, Italy

ABSTRACT. Bi-intuitionistic logic (BINT) is a conservative extension of intuitionistic logic to include the duals of each logical connective. One leading question with respect to BINT is, what does BINT look like across the three arcs – logic, typed λ -calculi, and category theory – of the Curry-Howard-Lambek correspondence? Categorically, BINT can be seen as a mixing of two worlds: the first being intuitionistic logic (IL), which is modeled by a cartesian closed category, and the second being the dual to intuitionistic logic called cointuitionistic logic (coIL), which is modeled by a cocartesian coclosed category. Crolard [10] showed that combining these two categories into the same category results in it degenerating to a poset. However, this degeneration does not occur when both logics are linear. We propose that IL and coIL need to be separated, and then mixed in a controlled way using the modalities from linear logic. This separation can be ultimately achieved by an adjoint formalization of bi-intuitionistic logic. This formalization consists of three worlds instead of two: the first is intuitionistic logic, the second is linear bi-intuitionistic (Bi-ILL), and the third is cointuitionistic logic. They are then related via two adjunctions. The adjunction between IL and ILL is known as a Linear/Non-linear model (LNL model) of ILL, and is due to Benton [4]. However, the dual to LNL models which would amount to the adjunction between coILL and coIL has yet to appear in the literature. In this paper we fill this gap by studying the dual to LNL models which we call dual LNL models. We conduct a similar analysis to that of Benton for dual LNL models by showing that dual LNL models correspond to dual linear categories, the dual to Bierman’s [5] linear categories proposed by Bellin [3]. Following this we give the definition of bi-LNL models by combining our dual LNL models with Benton’s LNL models to obtain a categorical model of bi-intuitionistic logic, but we leave its analysis and corresponding logic to a future paper. Finally, we give a corresponding sequent calculus, natural deduction, and term assignment for dual LNL models.


1. INTRODUCTION

Bi-intuitionistic logic (BINT) is a conservative extension of intuitionistic logic to include the duals of each logical connective. That is, BINT contains the usual intuitionistic logical connectives such as true, conjunction, and implication, but also their duals false, disjunction, and coimplication. One leading question with respect to BINT is, what does BINT look like across the three arcs – logic, typed λ -calculi, and category theory – of the Curry-Howard-Lambek correspondence? A non-trivial (does not degenerate to a poset) categorical model of BINT is currently an open problem. This paper directly contributes to the solution of this open problem by giving a new categorical model based on adjunctions for cointuitionistic logic, and then proposing a new categorical model for BINT.

BINT can be seen as a mixing of two worlds: the first being intuitionistic logic (IL), which is modeled categorically by a cartesian closed category (CCC), and the second being the dual to intuitionistic logic called cointuitionistic logic (coIL), which is modeled by a cocartesian coclosed category (coCCC). Crolard [10] showed that combining these two categories into the same category results in it degenerating to a poset, i.e. there is at most one morphism between any two objects; we review this result in Section 2.2. However, this degeneration does not occur when both logics are linear.

Notice that atoms are not dualized, at least in the main stream tradition of BINT started by C. Rauszer [19, 20]. For this reason T. Crolard [10] p. 160, describes the relation between IL and coIL within BINT as “pseudo duality”. A duality on atoms could be added and this has been attempted with linguistic motivations [2] (see the section on Related Work). This avoids the collapse but yields a different framework. Here we are concerned mainly with the main stream tradition.

We propose that IL and coIL need to be separated, and then mixed in a controlled way using the modalities from linear logic. This separation can be ultimately achieved by an adjoint formalization of bi-intuitionistic logic. This formalization consists of three worlds instead of two: the first is intuitionistic logic, the second is linear bi-intuitionistic (Bi-ILL), and the third is cointuitionistic logic. They are then related via two adjunctions as depicted by the following diagram:



The adjunction between IL and ILL is known as a Linear/Non-linear model (LNL model) of ILL, and is due to Benton [4]. However, the dual to LNL models which would amount to the adjunction between coILL and coIL has yet to appear in the literature.

Suppose $(\mathcal{I}, 1, \times, \rightarrow)$ is a cartesian closed category, and $(\mathcal{L}, \top, \otimes, \multimap)$ is a symmetric monoidal closed category. Then relate these two categories with a symmetric monoidal adjunction $\mathcal{I} : \mathcal{F} \dashv \mathcal{G} : \mathcal{L}$ (Definition 11), where \mathcal{F} and \mathcal{G} are symmetric monoidal functors. The later point implies that there are natural transformations $m_{X,Y} : \mathcal{F}X \otimes \mathcal{F}Y \rightarrow \mathcal{F}(X \times Y)$ and $n_{A,B} : \mathcal{G}A \times \mathcal{G}B \rightarrow \mathcal{G}(A \otimes B)$, and maps $m_\top : \top \rightarrow \mathcal{F}1$ and $n_1 : 1 \rightarrow \mathcal{G}\top$ subject to several coherence conditions; see Definition 7. Furthermore, the functor \mathcal{F} is strong which means that $m_{X,Y}$ and m_\top are isomorphisms. This setup turns out to be one of the most beautiful models of intuitionistic linear logic called a LNL model due to Benton [4]. In fact, the linear modality of-course can be defined by $!A = \mathcal{F}(\mathcal{G}(A))$ which defines a symmetric monoidal comonad using the adjunction; see Section 2.2 of [4]. This model is much simpler than other known models, and resulted in a logic called LNL logic which supports mixing intuitionistic logic with linear logic. The main contribution of this paper is the definition and study of the dual to Benton’s LNL models as models of cointuitionistic logic.

Taking the dual of the previous model results in what we call dual LNL models. They consist of a cocartesian coclosed category, $(\mathcal{C}, 0, +, -)$ where $- : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ is left adjoint to the coproduct, a symmetric monoidal coclosed category (Definition 4), $(\mathcal{L}', \perp, \oplus, \bullet)$, where $\bullet : \mathcal{L}' \times \mathcal{L}' \rightarrow \mathcal{L}'$ is left adjoint to cotensor (usually called *par*), and a symmetric comonoidal adjunction (Definition 12) $\mathcal{L}' : \mathcal{H} \dashv \mathcal{J} : \mathcal{C}$, where \mathcal{H} and \mathcal{J} are symmetric comonoidal functors. Dual to the above, this implies

that there are natural transformations $m_{X,Y} : J(X + Y) \longrightarrow JX \oplus JY$ and $\eta_{A,B} : H(A \oplus B) \longrightarrow HA + HB$, and maps $m_0 : J0 \longrightarrow \perp$ and $\eta_\perp : H \perp \longrightarrow 0$ subject to several coherence conditions; see Definition 8. In fact, one can define Girard's exponential why-not by $?A = JHA$, and hence, is the monad induced by the adjunction.

Bellin [3] was the first to propose the dual to Bierman's [5] linear categories which he names dual linear categories as a model of cointuitionistic linear logic. We conduct a similar analysis to that of Benton for dual LNL models by showing that dual LNL models are dual linear categories (Section 2.3.2), and that from a dual linear category we may obtain a dual LNL model (Section 2.3.3). Following this we give the definition of bi-LNL models by combining our dual LNL models with Benton's LNL models to obtain a categorical model of bi-intuitionistic logic (Section 2.4), but we leave its analysis and corresponding logic to a future paper.

Benton [4] showed that, syntactically, LNL models have a corresponding logic by first defining intuitionistic logic, whose sequent is denoted, $\Theta \vdash_C X$, and then intuitionistic linear logic, $\Theta; \Gamma \vdash_{\mathcal{L}} A$, but the key insight was that Θ contains non-linear assumptions while Γ contains linear assumptions, but one should view their separation as merely cosmetic; all assumptions can consistently be mixed within a single context. The two logics are then connected by syntactic versions of the functors F and G which allow formulas to move between both fragments.

Following Benton's lead the design of dual LNL logic is similar. We have a non-linear cointuitionistic fragment, $T \vdash_C \Psi$, and a linear cointuitionistic fragment, $A \vdash_{\mathcal{L}} \Delta; \Psi$, where Δ contains linear conclusions and Ψ contains non-linear conclusions, but again the separation of contexts is only cosmetic. The non-linear fragment has the following structural rules:

$$\frac{S \vdash_C \Psi}{S \vdash_C T, \Psi} C_weak \qquad \frac{S \vdash_C T, T, \Psi}{S \vdash_C T, \Psi} C_contr$$

Then we connect these two fragments together using the following rules for the functors H and J :

$$\frac{A \vdash_{\mathcal{L}} \cdot; \Psi}{HA \vdash_C \Psi} H_L \qquad \frac{A \vdash_{\mathcal{L}} \Delta, B; \Psi}{A \vdash_{\mathcal{L}} \Delta; HB, \Psi} H_R \qquad \frac{T \vdash_C \Psi}{JT \vdash_{\mathcal{L}} \cdot; \Psi} J_L \qquad \frac{A \vdash_{\mathcal{L}} \Delta; T, \Psi}{A \vdash_{\mathcal{L}} \Delta, JT; \Psi} J_R$$

These allow for linear and non-linear formulas to move from one fragment to the other. We will give a sequent calculus and natural deduction formalization (Section 3.1 and Section 3.2) as well as a term assignment (Section 3.3). The latter is particularly interesting, because of the fact that cointuitionistic logic has multiple conclusions, but only a single hypothesis.

2. THE ADJOINT MODEL

In this section we define dual LNL models (Definition 21) and then relate them to Bellin's dual linear categories (Definition 22), but first we introduce the basic categorical machinery needed for the later sections and summarize Crolard's result showing that the combination of cartesian closed categories with cocartesian coclosed categories are degenerate.

2.1. Symmetric (co)Monoidal Categories. We now introduce the necessary definitions related to symmetric monoidal categories that our model will depend on. Most of these definitions are equivalent to the ones given by Benton [4], but we give a lesser known definition of symmetric comonoidal functors due to Bellin [3]. In this section we also introduce distributive categories, the notion of coclosure, and finally, the definition of bilinear categories. The reader may wish to simply skim this section, but refer back to it when they encounter a definition or result they do not know.

Definition 1. A **symmetric monoidal category (SMC)** is a category, \mathcal{M} , with the following data:

- An object \top of \mathcal{M} ,
- A bi-functor $\otimes : \mathcal{M} \times \mathcal{M} \longrightarrow \mathcal{M}$,
- The following natural isomorphisms:

$$\begin{aligned}\lambda_A &: \top \otimes A \longrightarrow A \\ \rho_A &: A \otimes \top \longrightarrow A \\ \alpha_{A,B,C} &: (A \otimes B) \otimes C \longrightarrow A \otimes (B \otimes C)\end{aligned}$$

- A symmetry natural transformation:

$$\beta_{A,B} : A \otimes B \longrightarrow B \otimes A$$

- Subject to the following coherence diagrams:

$$\begin{array}{ccc} ((A \otimes B) \otimes C) \otimes D & \xrightarrow{\alpha_{A,B,C} \otimes \text{id}_D} & (A \otimes (B \otimes C)) \otimes D \\ \downarrow \alpha_{A \otimes B, C, D} & & \downarrow \alpha_{A, B \otimes C, D} \\ (A \otimes B) \otimes (C \otimes D) & & \\ \downarrow \alpha_{A, B, C \otimes D} & & \\ A \otimes (B \otimes (C \otimes D)) & \xleftarrow{\text{id}_A \otimes \alpha_{B, C, D}} & A \otimes ((B \otimes C) \otimes D) \end{array}$$

$$\begin{array}{ccccc} (A \otimes B) \otimes C & \xrightarrow{\alpha_{A,B,C}} & A \otimes (B \otimes C) & \xrightarrow{\beta_{A, B \otimes C}} & (B \otimes C) \otimes A \\ \downarrow \beta_{A, B} \otimes \text{id}_C & & & & \downarrow \alpha_{B, C, A} \\ (B \otimes A) \otimes C & \xrightarrow{\alpha_{B, A, C}} & B \otimes (A \otimes C) & \xrightarrow{\text{id}_B \otimes \beta_{A, C}} & B \otimes (C \otimes A) \end{array}$$

$$\begin{array}{ccc} (A \otimes \top) \otimes B & \xrightarrow{\alpha_{A, \top, B}} & A \otimes (\top \otimes B) \\ \searrow \rho_A & & \swarrow \lambda_B \\ & A \otimes B & \end{array}$$

$$\begin{array}{ccc} A \otimes B & & \\ \downarrow \beta_{A, B} & \searrow \text{id}_{A \otimes B} & \\ B \otimes A & \xrightarrow{\beta_{B, A}} & A \otimes B \end{array}$$

$$\begin{array}{ccc} \top \otimes A & \xrightarrow{\beta_{\top, A}} & A \otimes \top \\ \searrow \lambda_A & & \swarrow \rho_A \\ & A & \end{array}$$

Categorical modeling implication requires that the model be closed; which can be seen as an internalization of the notion of a morphism.

Definition 2. A **symmetric monoidal closed category (SMCC)** is a symmetric monoidal category, $(\mathcal{M}, \top, \otimes)$, such that, for any object B of \mathcal{M} , the functor $- \otimes B : \mathcal{M} \longrightarrow \mathcal{M}$ has a specified right

adjoint. Hence, for any objects A and C of \mathcal{M} there is an object $B \multimap C$ of \mathcal{M} and a natural bijection:

$$\text{Hom}_{\mathcal{M}}(A \otimes B, C) \cong \text{Hom}_{\mathcal{M}}(A, B \multimap C)$$

We call the functor $\multimap: \mathcal{M} \times \mathcal{M} \longrightarrow \mathcal{M}$ the internal hom of \mathcal{M} .

Symmetric monoidal closed categories can be seen as a model of intuitionistic linear logic with a tensor product and implication. What happens when we take the dual? First, we have the following result:

Lemma 3 (Dual of Symmetric Monoidal Categories). If $(\mathcal{M}, \top, \otimes)$ is a symmetric monoidal category, then \mathcal{M}^{op} is also a symmetric monoidal category.

The previous result follows from the fact that the structures making up symmetric monoidal categories are isomorphisms, and so naturally taking their opposite will yield another symmetric monoidal category. To emphasize when we are thinking about a symmetric monoidal category in the opposite we use the notation $(\mathcal{M}, \perp, \oplus)$ which gives the suggestion of \oplus corresponding to a disjunctive tensor product which we call the *cotensor* of \mathcal{M} . The next definition describes when a symmetric monoidal category is coclosed.

Definition 4. A **symmetric monoidal coclosed category** (SMCCC) is a symmetric monoidal category, $(\mathcal{M}, \perp, \oplus)$, such that, for any object B of \mathcal{M} , the functor $- \oplus B : \mathcal{M} \longrightarrow \mathcal{M}$ has a specified left adjoint. Hence, for any objects A and C of \mathcal{M} there is an object $C \multimap B$ of \mathcal{M} and a natural bijection:

$$\text{Hom}_{\mathcal{M}}(C, A \oplus B) \cong \text{Hom}_{\mathcal{M}}(C \multimap B, A)$$

We call the functor $\multimap: \mathcal{M} \times \mathcal{M} \longrightarrow \mathcal{M}$ the internal cohom of \mathcal{M} .

We combine a symmetric monoidal closed category with a symmetric monoidal coclosed category in a single category. First, we define the notion of a distributive category due to Cockett and Seely [9].

Definition 5. We call a symmetric monoidal category, $(\mathcal{M}, \top, \otimes, \perp, \oplus)$ equipped with the structure of a cotensor $(\mathcal{M}, \perp, \oplus)$, a **distributive category** if there are natural transformations:

$$\begin{aligned} \delta_{A,B,C}^L &: A \otimes (B \oplus C) \longrightarrow (A \otimes B) \oplus C \\ \delta_{A,B,C}^R &: (B \oplus C) \otimes A \longrightarrow B \oplus (C \otimes A) \end{aligned}$$

subject to several coherence diagrams. Due to the large number of coherence diagrams we do not list them here, but they all can be found in Cockett and Seely's paper [9].

Requiring that the tensor and cotensor products have the corresponding right and left adjoints results in the following definition.

Definition 6. A **bilinear category** is a distributive category $(\mathcal{M}, \top, \otimes, \perp, \oplus)$ such that $(\mathcal{M}, \top, \otimes)$ is closed, and $(\mathcal{M}, \perp, \oplus)$ is coclosed. We will denote bi-linear categories by $(\mathcal{M}, \top, \otimes, \multimap, \perp, \oplus, \multimap)$.

Originally, Lambek defined bilinear categories to be similar to the previous definition, but the tensor and cotensor were non-commutative [8], however, the bilinear categories given here are. We retain the name in homage to his original work. As we will see below bilinear categories form the core of a categorical model for bi-intuitionism.

A symmetric monoidal category is a category with additional structure subject to several coherence diagrams. Thus, an ordinary functor is not enough to capture this structure, and hence, the introduction of symmetric monoidal functors.

Definition 7. Suppose we are given two symmetric monoidal categories $(\mathcal{M}_1, \top_1, \otimes_1, \alpha_1, \lambda_1, \rho_1, \beta_1)$ and $(\mathcal{M}_2, \top_2, \otimes_2, \alpha_2, \lambda_2, \rho_2, \beta_2)$. Then a **symmetric monoidal functor** is a functor $F : \mathcal{M}_1 \rightarrow \mathcal{M}_2$, a map $m_{\top_1} : \top_2 \rightarrow F\top_1$ and a natural transformation $m_{A,B} : FA \otimes_2 FB \rightarrow F(A \otimes_1 B)$ subject to the following coherence conditions:

$$\begin{array}{ccc}
(FA \otimes_2 FB) \otimes_2 FC & \xrightarrow{\alpha_{2FA,FB,FC}} & FA \otimes_2 (FB \otimes_2 FC) \\
\downarrow m_{A,B} \otimes \text{id}_{FC} & & \downarrow \text{id}_{FA} \otimes m_{B,C} \\
F(A \otimes_1 B) \otimes_2 FC & & FA \otimes_2 F(B \otimes_1 C) \\
\downarrow m_{A \otimes_1 B, C} & & \downarrow m_{A, B \otimes_1 C} \\
F((A \otimes_1 B) \otimes_1 C) & \xrightarrow{F\alpha_{1A,B,C}} & F(A \otimes_1 (B \otimes_1 C))
\end{array}$$

$$\begin{array}{ccc}
\top_2 \otimes_2 FA & \xrightarrow{\lambda_{2FA}} & FA \\
\downarrow m_{\top_1} \otimes \text{id}_{FA} & & \uparrow F\lambda_{1A} \\
F\top_1 \otimes_2 FA & \xrightarrow{m_{\top_1, A}} & F(\top_1 \otimes_1 A)
\end{array}
\qquad
\begin{array}{ccc}
FA \otimes_2 \top_2 & \xrightarrow{\rho_{2FA}} & FA \\
\downarrow \text{id}_{FA} \otimes m_{\top_1} & & \uparrow F\rho_{1A} \\
FA \otimes_2 F\top_1 & \xrightarrow{m_{A, \top_1}} & F(A \otimes_1 \top_1)
\end{array}$$

$$\begin{array}{ccc}
FA \otimes_2 FB & \xrightarrow{\beta_{2FA,FB}} & FB \otimes_2 FA \\
\downarrow m_{A,B} & & \downarrow m_{B,A} \\
F(A \otimes_1 B) & \xrightarrow{F\beta_{1A,B}} & F(B \otimes_1 A)
\end{array}$$

The following is dual to the previous definition.

Definition 8. Suppose we are given two symmetric monoidal categories $(\mathcal{M}_1, \perp_1, \oplus_1, \alpha_1, \lambda_1, \rho_1, \beta_1)$ and $(\mathcal{M}_2, \perp_2, \oplus_2, \alpha_2, \lambda_2, \rho_2, \beta_2)$. Then a **symmetric comonoidal functor** is a functor $F : \mathcal{M}_1 \rightarrow \mathcal{M}_2$, a map $m_{\perp_1} : F\perp_1 \rightarrow \perp_2$ and a natural transformation $m_{A,B} : F(A \oplus_1 B) \rightarrow FA \oplus_2 FB$ subject to the following coherence conditions:

$$\begin{array}{ccc}
F((A \oplus_1 B) \oplus_1 C) & \xrightarrow{m_{A \oplus_1 B, C}} & F(A \oplus_1 B) \oplus_2 FC \\
\downarrow F\alpha_{A,B,C} & & \downarrow m_{A,B} \oplus \text{id}_{FC} \\
F(A \oplus_1 (B \oplus_1 C)) & & (FA \oplus_2 FB) \oplus_2 FC \\
\downarrow m_{A, B \oplus_1 C} & & \downarrow \alpha_{FA,FB,FC} \\
FA \oplus_2 F(B \oplus_1 C) & \xrightarrow{\text{id}_{FA} \oplus m_{B,C}} & FA \oplus_2 (FB \oplus_2 FC)
\end{array}$$

$$\begin{array}{ccc}
F(\perp_1 \oplus_1 A) & \xrightarrow{m_{\perp_1, A}} & F \perp_1 \oplus_2 FA \\
\downarrow F\lambda_{1A} & & \downarrow m_{\perp_1} \oplus \text{id}_{FA} \\
FA & \xrightarrow{\lambda_{2, FA}^{-1}} & \perp_2 \oplus_2 FA \\
\\
F(A \oplus_1 B) & \xrightarrow{m_{A, B}} & FA \oplus_2 FB \\
\downarrow F\beta_{1A, B} & & \downarrow \beta_{2FA, FB} \\
F(B \oplus_1 A) & \xrightarrow{m_{B, A}} & FB \oplus_2 FA
\end{array}
\qquad
\begin{array}{ccc}
F(A \oplus_1 \perp_1) & \xrightarrow{m_{A, \perp_1}} & FA \oplus_2 F \perp_1 \\
\downarrow F\rho_{1A} & & \downarrow \text{id}_{FA} \oplus m_{\perp_1} \\
FA & \xrightarrow{\rho_{2, FA}^{-1}} & FA \oplus_2 \perp_2
\end{array}$$

Naturally, since functors are enhanced to handle the additional structure found in a symmetric monoidal category we must also extend natural transformations, and adjunctions.

Definition 9. Suppose $(\mathcal{M}_1, \tau_1, \otimes_1)$ and $(\mathcal{M}_2, \tau_2, \otimes_2)$ are SMCs, and (F, m) and (G, n) are a symmetric monoidal functors between \mathcal{M}_1 and \mathcal{M}_2 . Then a **symmetric monoidal natural transformation** is a natural transformation, $f : F \rightarrow G$, subject to the following coherence diagrams:

$$\begin{array}{ccc}
FA \otimes_2 FB & \xrightarrow{m_{A, B}} & F(A \otimes_1 B) \\
\downarrow f_A \otimes_2 f_B & & \downarrow f_{A \otimes_1 B} \\
GA \otimes_2 GB & \xrightarrow{n_{A, B}} & G(A \otimes_1 B)
\end{array}
\qquad
\begin{array}{ccc}
F\tau_1 & \xrightarrow{f_{\tau_1}} & G\tau_1 \\
\swarrow m_{\tau_1} & & \searrow n_{\tau_1} \\
& \tau_2 &
\end{array}$$

Definition 10. Suppose $(\mathcal{M}_1, \perp_1, \oplus_1)$ and $(\mathcal{M}_2, \perp_2, \oplus_2)$ are SMCs, and (F, m) and (G, n) are a symmetric comonoidal functors between \mathcal{M}_1 and \mathcal{M}_2 . Then a **symmetric comonoidal natural transformation** is a natural transformation, $f : F \rightarrow G$, subject to the following coherence diagrams:

$$\begin{array}{ccc}
F(A \oplus_1 B) & \xrightarrow{m_{A, B}} & FA \oplus_2 FB \\
\downarrow f_{A \oplus_1 B} & & \downarrow f_A \oplus_2 f_B \\
G(A \oplus_1 B) & \xrightarrow{n_{A, B}} & GA \oplus_2 GB
\end{array}
\qquad
\begin{array}{ccc}
\perp_2 & \xleftarrow{n_{\perp_1}} & G \perp_1 \\
\swarrow m_{\perp_1} & & \searrow f_{\perp_1} \\
& F \perp_1 &
\end{array}$$

Definition 11. Suppose $(\mathcal{M}_1, \tau_1, \otimes_1)$ and $(\mathcal{M}_2, \tau_2, \otimes_2)$ are SMCs, and (F, m) is a symmetric monoidal functor between \mathcal{M}_1 and \mathcal{M}_2 and (G, n) is a symmetric monoidal functor between \mathcal{M}_2 and \mathcal{M}_1 . Then a **symmetric monoidal adjunction** is an ordinary adjunction $\mathcal{M}_1 : F \dashv G : \mathcal{M}_2$ such that the unit, $\eta_A : A \rightarrow GFA$, and the counit, $\varepsilon_A : FGA \rightarrow A$, are symmetric monoidal natural transformations.

Thus, the following diagrams must commute:

$$\begin{array}{ccc}
 FGA \otimes_2 FGB & \xrightarrow{m_{GA,GB}} & F(GA \otimes_1 GB) \\
 \downarrow \varepsilon_A \otimes_1 \varepsilon_B & & \downarrow Fn_{A,B} \\
 A \otimes_2 B & \xleftarrow{\varepsilon_A \otimes_1 B} & FGA \otimes_2 FGB
 \end{array}
 \qquad
 \begin{array}{ccc}
 F\tau_1 & \xrightarrow{Fn_{\tau_2}} & FG\tau_2 \\
 \uparrow m_{\tau_1} & & \downarrow \varepsilon_{\tau_1} \\
 \tau_2 & \xlongequal{\quad} & \tau_2
 \end{array}$$

$$\begin{array}{ccc}
 GFA \otimes_1 GFB & \xleftarrow{\eta_A \otimes_1 \eta_B} & A \otimes_1 B \\
 \downarrow n_{FA,FB} & & \downarrow \eta_{A \otimes_1 B} \\
 G(FA \otimes_2 FB) & \xrightarrow{m_{A,B}} & GF(A \otimes_1 B)
 \end{array}
 \qquad
 \begin{array}{ccc}
 G\tau_2 & \xrightarrow{Gm_{\tau_1}} & GF\tau_1 \\
 \uparrow n_{\tau_2} & & \uparrow \eta_{\tau_1} \\
 \tau_1 & \xlongequal{\quad} & \tau_1
 \end{array}$$

Definition 12. Suppose $(\mathcal{M}_1, \perp_1, \oplus_1)$ and $(\mathcal{M}_2, \perp_2, \oplus_2)$ are SMCs, and (F, m) is a symmetric comonoidal functor between \mathcal{M}_1 and \mathcal{M}_2 and (G, n) is a symmetric comonoidal functor between \mathcal{M}_2 and \mathcal{M}_1 . Then a **symmetric comonoidal adjunction** is an ordinary adjunction $\mathcal{M}_1 : F \dashv G : \mathcal{M}_2$ such that the unit, $\eta_A : A \rightarrow GFA$, and the counit, $\varepsilon_A : FGA \rightarrow A$, are symmetric comonoidal natural transformations. Thus, the following diagrams must commute:

$$\begin{array}{ccc}
 A \oplus_1 B & \xrightarrow{\eta_{A \oplus_1 B}} & GF(A \oplus_1 B) \\
 \downarrow \eta_A \oplus_1 \eta_B & & \downarrow Gm_{A,B} \\
 GFA \oplus_1 GFB & \xleftarrow{m_{FA,FB}} & G(FA \oplus_2 FB)
 \end{array}
 \qquad
 \begin{array}{ccc}
 GF \perp_1 & \xrightarrow{Gm_{\perp_1}} & G \perp_2 \\
 \uparrow \eta_{\perp_1} & & \downarrow n_{\perp_2} \\
 \perp_1 & \xlongequal{\quad} & \perp_1
 \end{array}$$

$$\begin{array}{ccc}
 FG(A \oplus_2 B) & \xrightarrow{Fn_{A,B}} & F(GA \oplus_1 GB) \\
 \downarrow \varepsilon_{A \oplus_2 B} & & \downarrow m_{GA,GB} \\
 A \oplus_2 B & \xleftarrow{\varepsilon_A \oplus_2 \varepsilon_B} & FGA \oplus_2 FGB
 \end{array}
 \qquad
 \begin{array}{ccc}
 FG \perp_2 & \xrightarrow{\varepsilon_{\perp_2}} & \perp_2 \\
 \parallel & & \uparrow m_{\perp_1} \\
 FG \perp_2 & \xrightarrow{Fn_{\perp_2}} & F \perp_1
 \end{array}$$

We will be defining, and making use of the why-not exponentials from linear logic, but these correspond to a symmetric comonoidal monad. In addition, whenever we have a symmetric comonoidal adjunction, we immediately obtain a symmetric comonoidal comonad on the left, and a symmetric comonoidal monad on the right.

Definition 13. A **symmetric comonoidal monad** on a symmetric monoidal category \mathcal{C} is a triple (T, η, μ) , where (T, n) is a symmetric comonoidal endofunctor on \mathcal{C} , $\eta_A : A \rightarrow TA$ and $\mu_A : T^2A \rightarrow TA$ are symmetric comonoidal natural transformations, which make the following diagrams commute:

$$\begin{array}{ccc}
 T^3A & \xrightarrow{\mu_{TA}} & T^2A \\
 \downarrow T\mu_A & & \downarrow \mu_A \\
 T^2A & \xrightarrow{\mu_A} & TA
 \end{array}
 \qquad
 \begin{array}{ccccc}
 & & TA & & \\
 & \nearrow & \uparrow \mu_A & \nwarrow & \\
 TA & \xrightarrow{\eta_{TA}} & T^2A & \xleftarrow{T\eta_A} & TA
 \end{array}$$

The assumption that η and μ are symmetric comonoidal natural transformations amount to the following diagrams commuting:

$$\begin{array}{ccc}
 A \oplus B & \xrightarrow{\eta_A \oplus \eta_B} & TA \oplus TB \\
 \downarrow \eta_A & \nearrow \eta_{A,B} & \\
 T(A \oplus B) & &
 \end{array}
 \qquad
 \begin{array}{ccc}
 \perp & \xrightarrow{\eta_\perp} & T\perp \\
 \parallel & \searrow \eta_\perp & \\
 & \perp &
 \end{array}$$

$$\begin{array}{ccccc}
 T^2(A \oplus B) & \xrightarrow{T\eta_{A,B}} & T(TA \oplus TB) & \xrightarrow{\eta_{TA,TB}} & T^2A \oplus T^2B \\
 \downarrow \mu_{A \oplus B} & & \downarrow \mu_A \oplus \mu_B & & \downarrow \mu_\perp \\
 T(A \oplus B) & \xrightarrow{\eta_{A,B}} & TA \oplus TB & & T\perp
 \end{array}
 \qquad
 \begin{array}{ccc}
 T^2\perp & \xrightarrow{T\eta_\perp} & T\perp \\
 \downarrow \mu_\perp & & \downarrow \eta_\perp \\
 T\perp & \xrightarrow{\eta_\perp} & \perp
 \end{array}$$

Finally, the dual concept of a symmetric comonoidal comonad.

Definition 14. A **symmetric comonoidal comonad** on a symmetric monoidal category C is a triple (T, ε, δ) , where (T, m) is a symmetric comonoidal endofunctor on C , $\varepsilon_A : TA \rightarrow A$ and $\delta_A : TA \rightarrow T^2A$ are symmetric comonoidal natural transformations, which make the following diagrams commute:

$$\begin{array}{ccc}
 TA & \xrightarrow{\delta_A} & T^2A \\
 \downarrow \delta_A & & \downarrow T\delta_A \\
 T^2A & \xrightarrow{\delta_{TA}} & T^3A
 \end{array}
 \qquad
 \begin{array}{ccc}
 & TA & \\
 \swarrow & \downarrow \delta_A & \searrow \\
 TA & \xleftarrow{\varepsilon_{TA}} T^2A \xrightarrow{T\varepsilon_A} & TA
 \end{array}$$

The assumption that ε and δ are symmetric monoidal natural transformations amount to the following diagrams commuting:

$$\begin{array}{ccc}
 T(A \oplus B) & \xrightarrow{m_{A,B}} & TA \oplus TB \\
 \searrow \varepsilon_{A \oplus B} & & \downarrow \varepsilon_A \oplus \varepsilon_B \\
 & & A \oplus B
 \end{array}
 \qquad
 \begin{array}{ccc}
 T\perp & \xrightarrow{\varepsilon_\perp} & \perp \\
 \parallel & \searrow m_\perp & \\
 & T\perp &
 \end{array}$$

$$\begin{array}{ccccc}
 T(A \oplus B) & \xrightarrow{m_{A,B}} & TA \oplus TB & & \\
 \downarrow \delta_{A \oplus B} & & \downarrow \delta_A \oplus \delta_B & & \\
 T^2(A \oplus B) & \xrightarrow{Tm_{A,B}} & T(TA \oplus TB) & \xrightarrow{m_{TA,TB}} & T^2A \oplus T^2B
 \end{array}
 \qquad
 \begin{array}{ccc}
 T\perp & \xrightarrow{m_\perp} & \perp \\
 \downarrow \delta_\perp & & \downarrow m_\perp \\
 T^2\perp & \xrightarrow{Tm_\perp} & T\perp
 \end{array}$$

2.2. Cartesian Closed and Cocartesian Coclosed Categories. The notion of a cartesian closed category is well-known, but for completeness we define them here. However, their dual is lesser known, especially in computer science, and so we give their full definition. We also review some known results concerning cocartesian coclosed categories and categories that are both cartesian closed and cocartesian coclosed.

Definition 15. A **cartesian category** is a category, $(C, 1, \times)$, with an object, 1 , and a bi-functor, $\times : C \times C \longrightarrow C$, such that for any object A there is exactly one morphism $\diamond : A \rightarrow 1$, and for any morphisms $f : C \longrightarrow A$ and $g : C \longrightarrow B$ there is a morphism $\langle f, g \rangle : C \rightarrow A \times B$ subject to the following diagram:

$$\begin{array}{ccccc}
 & & C & & \\
 & f \swarrow & \downarrow \langle f, g \rangle & \searrow g & \\
 A & \xleftarrow{\pi_1} & A \times B & \xrightarrow{\pi_2} & B
 \end{array}$$

A cartesian category models conjunction by the product functor, $\times : C \times C \longrightarrow C$, and the unit of conjunction by the terminal object. As we mention above modeling implication requires closure, and since it is well-known that any cartesian category is also a symmetric monoidal category the definition of closure for a cartesian category is the same as the definition of closure for a symmetric monoidal category (Definition 2). We denote the internal hom for cartesian closed categories by $A \rightarrow B$.

The dual of a cartesian category is a cocartesian category. They are a model of intuitionistic logic with disjunction and its unit.

Definition 16. A **cocartesian category** is a category, $(C, 0, +)$, with an object, 0 , and a bi-functor, $+ : C \times C \longrightarrow C$, such that for any object A there is exactly one morphism $\square : 0 \rightarrow A$, and for any morphisms $f : A \longrightarrow C$ and $g : B \longrightarrow C$ there is a morphism $[f, g] : A + B \longrightarrow C$ subject to the following diagram:

$$\begin{array}{ccccc}
 & & C & & \\
 & f \swarrow & \uparrow [f, g] & \nwarrow g & \\
 A & \xrightarrow{\iota_1} & A + B & \xleftarrow{\iota_2} & B
 \end{array}$$

Coclosure, just like closure for cartesian categories, is defined in the same way that coclosure is defined for symmetric monoidal categories, because cocartesian categories are also symmetric monoidal categories. Thus, a cocartesian category is coclosed if there is a specified left-adjoint, which we denote $S - T$, to the coproduct.

There are many examples of cocartesian coclosed categories. Basically, any interesting cartesian category has an interesting dual, and hence, induces an interesting cocartesian coclosed category. The opposite of the category of sets and functions between them is isomorphic to the category of complete atomic boolean algebras, and both of which, are examples of cocartesian coclosed categories. As we mentioned above bi-linear categories [8] are models of bi-linear logic where the left adjoint to the cotensor models coimplication. Similarly, cocartesian coclosed categories model cointuitionistic logic with disjunction and intuitionistic coimplication.

We might now ask if a category can be both cartesian closed and cocartesian coclosed just as bi-linear categories, but this turns out to be where the matter meets antimatter in such away that the

category degenerates to a preorder. That is, every homspace contains at most one morphism. We recall this proof here, which is due to Crolard [10]. We need a couple basic facts about cartesian closed categories with initial objects.

Lemma 17. In any cartesian category C , if 0 is an initial object in C and $\text{Hom}_C(A, 0)$ is non-empty, then $A \cong A \times 0$.

Proof. This follows easily from the universal mapping property for products. \square

Lemma 18. In any cartesian closed category C , if 0 is an initial object in C , then so is $0 \times A$ for any object A of C .

Proof. We know that the universal morphism for the initial object is unique, and hence, the homspace $\text{Hom}_C(0, A \Rightarrow B)$ for any object B of C contains exactly one morphism. Then using the right adjoint to the product functor we know that $\text{Hom}_C(0, A \Rightarrow B) \cong \text{Hom}_C(0 \times A, B)$, and hence, there is only one arrow between $0 \times A$ and B . \square

The following lemma is due to Joyal [?], and is key to the next theorem.

Lemma 19 (Joyal's). In any cartesian closed category C , if 0 is an initial object in C and $\text{Hom}_C(A, 0)$ is non-empty, then A is an initial object in C .

Proof. Suppose C is a cartesian closed category, such that, 0 is an initial object in C , and A is an arbitrary object in C . Furthermore, suppose $\text{Hom}_C(A, 0)$ is non-empty. By the first basic lemma above we know that $A \cong A \times 0$, and by the second $A \times 0$ is initial, thus A is initial. \square

Finally, the following theorem shows that any category that is both cartesian closed and cocartesian coclosed is a preorder.

Theorem 20 ((co)Cartesian (co)Closed Categories are Preorders (Crolard[10])). If C is both cartesian closed and cocartesian coclosed, then for any two objects A and B of C , $\text{Hom}_C(A, B)$ has at most one element.

Proof. Suppose C is both cartesian closed and cocartesian coclosed, and A and B are objects of C . Then by using the basic fact that the initial object is the unit to the coproduct, and the coproducts left adjoint we know the following:

$$\text{Hom}_C(A, B) \cong \text{Hom}_C(A, 0 + B) \cong \text{Hom}_C(B - A, 0)$$

Therefore, by Joyal's theorem above $\text{Hom}_C(A, B)$ has at most one element. \square

Notice that the previous result hinges on the fact that there are initial and terminal objects, and thus, this result does not hold for bi-linear categories, because the units to the tensor and cotensor are not initial nor terminal.

The repercussions of this result are that if we do not want to work with preorders, but do want to work with all of the structure, then we must separate the two worlds. Thus, this result can be seen as the motivation for the current work. We enforce the separation using linear logic, but through the power of linear logic this separation is not far.

2.3. A Mixed Linear/Non-Linear Model for Co-Intuitionistic Logic. Benton [4] showed that from a LNL model it is possible to construct a linear category, and vice versa. Bellin [3] showed that the dual to linear categories are sufficient to model co-intuitionistic linear logic. We show that from the dual to a LNL model we can construct the dual to a linear category, and vice versa, thus, carrying out the same program for co-intuitionistic linear logic as Benton did for intuitionistic linear logic.

Combining a symmetric monoidal coclosed category with a cocartesian coclosed category via a symmetric comonoidal adjunction defines a dual LNL model.

Definition 21. A mixed linear/non-linear model for co-intuitionistic logic (dual LNL model), $\mathcal{L} : H \dashv J : C$, consists of the following:

- i. a symmetric monoidal coclosed category $(\mathcal{L}, \perp, \oplus, \bullet-)$,
- ii. a cocartesian coclosed category $(C, 0, +, -)$, and
- iv. a symmetric comonoidal adjunction $\mathcal{L} : H \dashv J : C$, where $\eta_A : A \longrightarrow JHA$ and $\varepsilon_R : HJR \longrightarrow R$ are the unit and counit of the adjunction respectively.

It is well-known that an adjunction $\mathcal{L} : H \dashv J : C$ induces a monad $H; J : \mathcal{L} \longrightarrow \mathcal{L}$, but when the adjunction is symmetric comonoidal we obtain a symmetric comonoidal monad, in fact, $H; J$ defines the linear exponential why-not denoted $?A = JHA$. By the definition of dual LNL models we know that both H and J are symmetric comonoidal functors, and hence, are equipped with natural transformations $h_{A,B} : H(A \oplus B) \longrightarrow HA + HB$ and $j_{R,S} : J(R + S) \longrightarrow JR \oplus JS$, and maps $h_\perp : H \perp \longrightarrow 0$ and $j_0 : J0 \longrightarrow \perp$. We will make heavy use of these maps throughout the sequel.

Compare this definition with that of Bellin's dual linear category from [3], and we can easily see that the definition of dual LNL models – much like LNL models – is more succinct.

Definition 22. A dual linear category, \mathcal{L} , consists of the following data:

- i. A symmetric monoidal coclosed category $(\mathcal{L}, \oplus, \perp, \bullet-)$ with
- ii. a symmetric co-monoidal monad $(?, \eta, \mu)$ on \mathcal{L} such that
 - a. each free $?$ -algebra carries naturally the structure of a commutative \oplus -monoid. This implies that there are distinguished symmetric monoidal natural transformations $w_A : \perp \longrightarrow ?A$ and $c_A : ?A \oplus ?A \longrightarrow ?A$ which form a commutative monoid and are $?$ -algebra morphisms.
 - b. whenever $f : (?A, \mu_A) \longrightarrow (?B, \mu_B)$ is a morphism of free $?$ -algebras, then it is also a monoid morphism.

2.3.1. A Useful Isomorphism. One useful property of Benton's LNL model is that the maps associated with the symmetric monoidal left adjoint in the model are isomorphisms. Since dual LNL models are dual we obtain similar isomorphisms with respect to the right adjoint.

Lemma 23 (Symmetric Comonoidal Isomorphisms). Given any dual LNL model $\mathcal{L} : H \dashv J : C$, then there are the following isomorphisms:

$$J(R + S) \cong JR \oplus JS \quad \text{and} \quad J0 \cong \perp$$

Furthermore, the former is natural in R and S .

Proof. Suppose $\mathcal{L} : H \dashv J : C$ is a dual LNL model. Then we can define the following family of maps:

$$\begin{aligned} j_{R,S}^{-1} &:= JR \oplus JS \xrightarrow{\eta} JH(JR \oplus JS) \xrightarrow{Jh_{A,B}} J(HJR + HJS) \xrightarrow{J(\varepsilon_R + \varepsilon_S)} J(R + S) \\ j_0^{-1} &:= \perp \xrightarrow{\eta} JH \perp \xrightarrow{Jh_\perp} J0 \end{aligned}$$

It is easy to see that $j_{R,S}^{-1}$ is natural, because it is defined in terms of a composition of natural transformations. All that is left to be shown is that $j_{R,S}^{-1}$ and j_0^{-1} are mutual inverses with $j_{R,S}$ and j_0 ; for the details see Appendix B.1. \square

Just as Benton we also do not have similar isomorphisms with respect to the functor H . One fact that we can point out, that Benton did not make explicit – because he did not use the notion of symmetric comonoidal functor – is that j^{-1} makes J also a symmetric monoidal functor.

Corollary 24. Given any dual LNL model $\mathcal{L} : H \dashv J : C$, the functor (J, j^{-1}) is symmetric monoidal.

Proof. This holds by straightforwardly reducing the diagrams defining a symmetric monoidal functor, Definition 7, to the diagrams defining a symmetric comonoidal functor, Definition 8, using the fact that j^{-1} is an isomorphism. \square

2.3.2. Dual LNL Model Implies Dual Linear Category. The next result shows that any dual LNL model induces a symmetric comonoidal monad.

Lemma 25 (Symmetric Comonoidal Monad). Given a dual LNL model $\mathcal{L} : H \dashv J : C$, the functor, $? = H; J$, defines a symmetric comonoidal monad.

Proof. Suppose (H, h) and (J, j) are two symmetric comonoidal functors, such that, $\mathcal{L} : H \dashv J : C$ is a dual LNL model. We can easily show that $?A = JHA$ is symmetric monoidal by defining the following maps:

$$\begin{aligned} r_{\perp} &:= ?\perp \equiv JH\perp \xrightarrow{j_{H\perp}} J0 \xrightarrow{j_{\perp}} \perp \\ r_{A,B} &:= ?(A \oplus B) \equiv JH(A \oplus B) \xrightarrow{j_{HA,B}} J(HA + HB) \xrightarrow{j_{HA,HB}} JHA \oplus JHB \equiv ?A \oplus ?B \end{aligned}$$

The fact that these maps satisfy the appropriate symmetric comonoidal functor diagrams from Definition 8 is obvious, because symmetric comonoidal functors are closed under composition.

We have a dual LNL model, and hence, we have the symmetric comonoidal natural transformations $\eta_A : A \longrightarrow JHA$ and $\varepsilon_R : HJR \longrightarrow R$ which correspond to the unit and counit of the adjunction respectively. Define $\mu_A := J\varepsilon_{HA} : JHJHA \longrightarrow JHA$. This implies that we have maps $\eta_A : A \longrightarrow ?A$ and $\mu_A : ??A \longrightarrow ?A$, and thus, we can show that $(?, \eta, \mu)$ is a symmetric comonoidal monad. All the diagrams defining a symmetric comonoidal monad hold by the structure given by the adjunction. For the complete proof see Appendix B.2. \square

The monad from the previous result must be equipped with the additional structure to model the right weakening and contraction structural rules.

Lemma 26 (Right Weakening and Contraction). Given a dual LNL model $\mathcal{L} : H \dashv J : C$, then for any $?A$ there are distinguished symmetric comonoidal natural transformations $w_A : \perp \longrightarrow ?A$ and $c_A : ?A \oplus ?A \longrightarrow ?A$ that form a commutative monoid, and are $?$ -algebra morphisms with respect to the canonical definitions of the algebras $?A, \perp, ?A \oplus ?A$.

Proof. Suppose (H, h) and (J, j) are two symmetric comonoidal functors, such that, $\mathcal{L} : H \dashv J : C$ is a dual LNL model. Again, we know $?A = H; J : \mathcal{L} \longrightarrow \mathcal{L}$ is a symmetric comonoidal monad by Lemma 25.

We define the following morphisms:

$$\begin{aligned} w_A &:= \perp \xrightarrow{j_{\perp}^{-1}} J0 \xrightarrow{j_{\circ HA}} JHA \equiv ?A \\ c_A &:= ?A \oplus ?A \equiv JHA \oplus JHA \xrightarrow{j_{HA,HA}^{-1}} J(HA + HA) \xrightarrow{j_{\nabla HA}} JHA \equiv ?A \end{aligned}$$

The remainder of the proof is by carefully checking all of the required diagrams. Please see Appendix B.3 for the complete proof. \square

Lemma 27 (*?-Monoid Morphisms*). Suppose $\mathcal{L} : \mathbf{H} \dashv \mathbf{J} : \mathbf{C}$ is a dual LNL model. Then if $f : (?A, \mu_A) \longrightarrow (?B, \mu_B)$ is a morphism of free ?-algebras, then it is a monoid morphism.

Proof. Suppose $\mathcal{L} : \mathbf{H} \dashv \mathbf{J} : \mathbf{C}$ is a dual LNL model. Then we know $?A = \mathbf{JHA}$ is a symmetric comonoidal monad by Lemma 25. Bellin [3] remarks that by Maietti, Maneggia de Paiva and Ritter's Proposition 25 [15], it suffices to show that $\mu_A : ??A \longrightarrow ?A$ is a monoid morphism. For the details see the complete proof in Appendix B.4. \square

Finally, we may now conclude the following corollary.

Corollary 28. Every dual LNL model is a dual linear category.

2.3.3. Dual Linear Category implies Dual LNL Model. This section shows essentially the inverse to the result from the previous section. That is, that from any dual linear category we may construct a dual LNL model. By exploiting the duality between LNL models and dual LNL models this result follows straightforwardly from Benton's result. The proof of this result must first find a symmetric monoid coclosed category, a cocartesian coclosed category, and finally, a symmetric comonoidal adjunction between them. Take the symmetric monoid coclosed category to be an arbitrary dual linear category \mathcal{L} . Then we may define the following categories.

- The Eilenberg-Moore category, $\mathcal{L}^?$, has as objects all ?-algebras, $(A, h_A : ?A \longrightarrow A)$, and as morphisms all ?-algebra morphisms.
- The Kleisli category, $\mathcal{L}_?$, is the full subcategory of $\mathcal{L}^?$ of all free ?-algebras $(?A, \mu_A : ??A \longrightarrow ?A)$.

The previous three categories are related by a pair of adjunctions:

$$\begin{array}{ccc}
 \mathcal{L} & \xrightleftharpoons[F]{U} & \mathcal{L}^? \\
 \parallel & & \uparrow i \\
 \mathcal{L} & \xrightleftharpoons[U]{F} & \mathcal{L}_?
 \end{array}$$

The functor $F(A) = (?A, \mu_A)$ is the free functor, and the functor $U(A, h_A) = A$ is the forgetful functor. Note that we, just as Benton did, are overloading the symbols F and U . Lastly, the functor $i : \mathcal{L}_? \longrightarrow \mathcal{L}^?$ is the injection of the subcategory of free ?-algebras into its parent category.

We are now going to show that both $\mathcal{L}^?$ and $\mathcal{L}_?$ are induce two cocartesian coclosed categories. Then we could take either of those when constructing a dual LNL model from a dual linear category. First, we show $\mathcal{L}^?$ is cocartesian.

Lemma 29. If \mathcal{L} is a dual linear category, then $\mathcal{L}^?$ has finite coproducts.

Proof. We give a proof sketch of this result, because the proof is essentially by duality of Benton's corresponding proof for LNL models (see Lemma 9, [4]). Suppose \mathcal{L} is a dual linear category. Then we first need to identify the initial object which is defined by the ?-algebra $(\perp, r_\perp : ?\perp \longrightarrow \perp)$. The unique map between the initial map and any other ?-algebra $(A, h_A : ?A \longrightarrow A)$ is defined by $\perp \xrightarrow{w_A} ?A \xrightarrow{h_A} A$. The coproduct of the ?-algebras $(A, h_A : ?A \longrightarrow A)$ and $(B, h_B : ?B \longrightarrow B)$ is $(A \oplus B, r_{A,B}; (h_A \oplus h_B))$. Injections and the codiagonal map are defined as follows:

- Injections:

$$\begin{aligned}\iota_1 &:= A \xrightarrow{\rho_A} A \oplus \perp \xrightarrow{\text{id}_A \oplus w_B} A \oplus ?B \xrightarrow{\text{id} \oplus h_b} A \oplus B \\ \iota_2 &:= B \xrightarrow{\lambda_A} \perp \oplus B \xrightarrow{w_A \oplus \text{id}_B} ?A \oplus B \xrightarrow{h_A \oplus \text{id}_B} A \oplus B\end{aligned}$$

- Codagonal map:

$$\nabla := A \oplus A \xrightarrow{\eta_A \oplus \eta_A} ?A \oplus ?A \xrightarrow{c_A} ?A \xrightarrow{h_A} A$$

Showing that these respect the appropriate diagrams is straightforward. \square

Notice as a direct consequence of the previous result we know the following.

Corollary 30. The Kleisli category, $\mathcal{L}_?$, has finite coproducts.

Thus, both $\mathcal{L}^?$ and $\mathcal{L}_?$ are cocartesian, but we need a cocartesian coclosed category, and in general these are not coclosed, and so we follow Benton's lead and show that there are actually two subcategories of $\mathcal{L}^?$ that are coclosed.

Definition 31. We call an object, A , of a category, \mathcal{L} , **subtractable** if for any object B of \mathcal{L} , the internal cohom $A \bullet - B$ exists.

We now have the following results:

Lemma 32. In $\mathcal{L}^?$, all the free $?$ -algebras are subtractable, and the internal cohom is a free $?$ -algebra.

Proof. The internal cohom is defined as follows:

$$(?A, \delta_A) \bullet - (B, h_B) := (?A \bullet - B, \delta_{A \bullet - B})$$

We can capitalize on the adjunctions involving F and U from above to lift the internal cohom of \mathcal{L} into $\mathcal{L}^?$:

$$\begin{aligned}\text{Hom}_{\mathcal{L}^?}((?A \bullet - B), \delta_{A \bullet - B}), (C, h_C)) &= \text{Hom}_{\mathcal{L}^?}(F(A \bullet - B), (C, h_C)) \\ &\cong \text{Hom}_{\mathcal{L}}(A \bullet - B, U(C, h_C)) \\ &= \text{Hom}_{\mathcal{L}}(A \bullet - B, C) \\ &\cong \text{Hom}_{\mathcal{L}}(A, C \oplus B) \\ &= \text{Hom}_{\mathcal{L}}(A, U(C \oplus B, h_{C \oplus B})) \\ &\cong \text{Hom}_{\mathcal{L}^?}(FA, (C \oplus B, h_{C \oplus B})) \\ &= \text{Hom}_{\mathcal{L}^?}((?A, \delta_A), (C \oplus B, h_{C \oplus B}))\end{aligned}$$

The previous equation holds for any $h_{C \oplus B}$ making $C \oplus B$ a $?$ -algebra, in particular, the co-product in $\mathcal{L}^?$ (Lemma 29), and hence, we may instantiate the final line of the previous equation with the following:

$$\text{Hom}_{\mathcal{L}^?}((?A, \delta_A), (C, h_C) + (B, \delta_B))$$

Thus, we obtain our result. \square

Lemma 33. We have the following cocartesian coclosed categories:

- i. The full subcategory, $\text{Sub}(\mathcal{L}^?)$, of $\mathcal{L}^?$ consisting of objects the subtractable $?$ -algebras is cocartesian coclosed, and contains the Kleisli category.
- ii. The full subcategory, \mathcal{L}_γ^* , of $\text{Sub}(\mathcal{L}^?)$ consisting of finite coproducts of free $?$ -algebras is cocartesian coclosed.

Let C be either of the previous two categories. Then we must exhibit an adjunction between C and \mathcal{L} , but this is easily done.

Lemma 34. The adjunction $\mathcal{L} : F \vdash U : C$, with the free functor, F , and the forgetful functor, U , is symmetric comonoidal.

Proof. Showing that F and U are symmetric comonoidal follows similar reasoning to Benton's result, but in the opposite; see Lemma 13 and Lemma 14 of [4]. Lastly, showing that the unit and the counit of the adjunction are comonoidal natural transformations is straightforward, and we leave it to the reader. The reasoning is similar to Benton's, but in the opposite; see Lemma 15 and Lemma 16 of [4]. \square

Corollary 35. Any dual linear category gives rise to a dual LNL model.

2.4. A Mixed Bilinear/Non-Linear Model. The main goal of our research program is to give a non-trivial categorical model of bi-intuitionistic logic. In this section we give an introduction of the model we have in mind, but leave the details and the study of the logical and programmatic sides to future work.

The naive approach would be to try and define a LNL-style model of bi-intuitionistic logic as an adjunction between a bilinear category and a bi-cartesian bi-closed category, but this results in a few problems. First, should the adjunction be monoidal or comonoidal? Furthermore, we know bi-cartesian bi-closed categories are trivial (Theorem 20), and hence, this model is not very interesting nor incorrect. We must separate the two worlds using two dual adjunctions, and hence, we arrive at the following definition.

Definition 36. A **mixed bilinear/non-linear model** consists of the following:

- i. a bilinear category $(\mathcal{L}, \top, \otimes, \multimap, \perp, \oplus, \multimap)$,
- ii. a cartesian closed category $(\mathcal{I}, 1, \times, \rightarrow)$,
- iii. a cocartesian coclosed category $(C, 0, +, -)$,
- iv. a LNL model $\mathcal{I} : F \dashv G : \mathcal{L}$, and
- v. a dual LNL model $\mathcal{L} : H \dashv J : C$.

Since \mathcal{L} is a bilinear category then it is also a linear category, and a dual linear category. Thus, the LNL model intuitively corresponds to an adjunction between \mathcal{I} and the linear subcategory of \mathcal{L} , and the dual LNL model corresponds to an adjunction between the dual linear subcategory of \mathcal{L} and C . In addition, both intuitionistic logic and cointuitionistic logic can be embedded into \mathcal{L} via the linear modalities of-course, $!A$, and why-not, $?A$, using the well-known Girard embeddings. This implies that we have a very controlled way of mixing \mathcal{I} and C within \mathcal{L} , and hence, linear logic is the key.

$\frac{}{S \vdash_C S} \text{C_id}$	$\frac{S \vdash_C \Psi}{S \vdash_C T, \Psi} \text{C_weak}$	$\frac{S \vdash_C T, T, \Psi}{S \vdash_C T, \Psi} \text{C_contr}$	$\frac{R \vdash_C \Psi_1, S, T, \Psi_2}{R \vdash_C \Psi_1, T, S, \Psi_2} \text{C_ex}$
$\frac{}{0 \vdash_C \Psi} \text{C_0}$	$\frac{T_1 \vdash_C \Psi_1 \quad T_2 \vdash_C \Psi_2}{T_1 + T_2 \vdash_C \Psi_1, \Psi_2} \text{C_+L}$	$\frac{R \vdash_C \Psi, T_1}{R \vdash_C \Psi, T_1 + T_2} \text{C_+R}_1$	
$\frac{R \vdash_C \Psi, T_2}{R \vdash_C \Psi, T_1 + T_2} \text{C_+R}_2$	$\frac{T_1 \vdash_C T_2, \Psi}{T_1 - T_2 \vdash_C \Psi} \text{C_+L}$	$\frac{S \vdash_C \Psi_1, T_1 \quad T_2 \vdash_C \Psi_2}{S \vdash_C \Psi_1, \Psi_2, T_1 - T_2} \text{C_+R}$	
	$\frac{S \vdash_C \Psi_1, T \quad T \vdash_C \Psi_2}{S \vdash_C \Psi_1, \Psi_2} \text{C_cut}$	$\frac{A \vdash_L \cdot; \Psi}{HA \vdash_C \Psi} \text{H}_L$	

Figure 1: Non-linear fragment of the DLNL logic

3. DUAL LNL LOGIC

We now turn to developing the syntactic side of dual LNL models called dual LNL logic (DLNL). First, we give a sequent calculus formalization which we will simply refer to as DLNL logic, then a natural deduction formalization called DND logic, and finally a term assignment to the natural deduction version. Each of these systems will consistently use the same syntax and naming conventions for formulas, types, and contexts given by the following definition.

Definition 37. The the syntax for formulas, types, and contexts are given as follows:

$$\begin{aligned}
 & \text{(non-linear formulas/types)} \quad R, S, T ::= 0 \mid S + T \mid S - T \mid HA \\
 & \text{(linear formulas/types)} \quad A, B, C ::= \perp \mid A \oplus B \mid A \multimap B \mid JS \\
 & \text{(non-linear contexts)} \quad \Psi, \Theta ::= \cdot \mid T \mid \Psi, \Theta \\
 & \text{(linear contexts)} \quad \Gamma, \Delta ::= \cdot \mid A \mid \Gamma, \Delta
 \end{aligned}$$

The term assignment will index contexts by terms, but we will maintain the same naming convention throughout.

3.1. The Sequent Calculus for Dual LNL Logic. In this section we take the dual of Benton's [4] sequent calculus for LNL logic to obtain the sequent calculus for dual LNL logic. The inference rules for the non-linear fragment can be found in Figure 1 and the linear fragment in Figure 2. The remainder of this section is devoted to proving cut-elimination. However, the proof is simply a dualization of Benton's [4] proof of cut-elimination for LNL logic.

Just as Benton we use n -ary cuts:

$$\frac{S \vdash_C \Psi, S^n \quad S \vdash_C \Psi'}{S \vdash_C \Psi, \Psi'} \text{C_cut}_n \quad \frac{A \vdash_L \Delta; \Psi, S^n \quad S \vdash_C \Psi'}{A \vdash_L \Delta; \Psi, \Psi'} \text{LC_cut}_n$$

where $S^n = S, \dots, S$ n -times. We call DLNL^+ the system DLNL with n -cuts replacing ordinary 1-cuts. Such cuts are admissible in DLNL and cut-elimination for DLNL^+ implies cut-elimination for DLNL.

We begin with the a few standard definitions. The *rank* of a formula, denoted by $|A|$ or $|S|$, is the number of the logical symbols in the given formula. The *cut-rank* of a derivation Π , denoted by $c(\Pi)$, is the maximum of the ranks of the cut formulas in Π plus one; if Π is cut-free its cut rank is

$$\begin{array}{c}
\frac{}{A \vdash_L A; \cdot} \text{LL_id} \quad \frac{A \vdash_L \Delta; \Psi}{A \vdash_L \Delta; T, \Psi} \text{LC_weak} \quad \frac{A \vdash_L \Delta; T, T, \Psi}{A \vdash_L \Delta; T, \Psi} \text{LC_contr} \\
\\
\frac{A \vdash_L \Delta_1, A, B, \Delta_2; \Psi}{A \vdash_L \Delta_1, B, A, \Delta_2; \Psi} \text{LL_ex} \quad \frac{A \vdash_L \Delta; \Psi_1, S, T, \Psi_2}{A \vdash_L \Delta; \Psi_1, T, S, \Psi_2} \text{LC_ex} \\
\\
\frac{A \vdash_L \Delta_1, B; \Psi_1 \quad B \vdash_L \Delta_2; \Psi_2}{A \vdash_L \Delta_1, \Delta_2; \Psi_1, \Psi_2} \text{LL_cut} \quad \frac{A \vdash_L \Delta; \Psi_1, T \quad T \vdash_C \Psi_2}{A \vdash_L \Delta; \Psi_1, \Psi_2} \text{LC_cut} \\
\\
\frac{}{\perp \vdash_L \cdot; \cdot} \text{LL_}\perp_L \quad \frac{A \vdash_L \Delta; \Psi}{A \vdash_L \perp, \Delta; \Psi} \text{LL_}\perp_R \quad \frac{A \vdash_L \Delta; \Psi, T_1}{A \vdash_L \Delta; \Psi, T_1 + T_2} \text{LC_}\perp_{R_1} \\
\\
\frac{A \vdash_L \Delta; \Psi, T_2}{A \vdash_L \Delta; \Psi, T_1 + T_2} \text{LC_}\perp_{R_2} \quad \frac{B_1 \vdash_L \Delta_1; \Psi_1 \quad B_2 \vdash_L \Delta_2; \Psi_2}{B_1 \oplus B_2 \vdash_L \Delta_1, \Delta_2; \Psi_1, \Psi_2} \text{LL_}\oplus_L \\
\\
\frac{A \vdash_L \Delta, B, C; \Psi}{A \vdash_L \Delta, B \oplus C; \Psi} \text{LL_}\oplus_R \quad \frac{B_1 \vdash_L B_2, \Delta; \Psi}{B_1 \bullet B_2 \vdash_L \Delta; \Psi} \text{LL_}\bullet_L \\
\\
\frac{A \vdash_L B_1, \Delta_1; \Psi_1 \quad B_2 \vdash_L \Delta_2; \Psi_2}{A \vdash_L B \bullet C, \Delta_1, \Delta_2; \Psi_1, \Psi_2} \text{LL_}\bullet_R \quad \frac{A \vdash_L \Delta; \Psi_1, T_1 \quad T_2 \vdash_C \Psi_2}{A \vdash_L \Delta; \Psi_1, \Psi_2, T_1 - T_2} \text{LL_}\neg_R \\
\\
\frac{T \vdash_C \Psi}{JT \vdash_L \cdot; \Psi} J_L \quad \frac{A \vdash_L \Delta; T, \Psi}{A \vdash_L \Delta, JT; \Psi} J_R \quad \frac{A \vdash_L \Delta, B; \Psi}{A \vdash_L \Delta; HB, \Psi} H_R
\end{array}$$

Figure 2: Linear fragment of the DLNL logic

0. Finally, the *depth* of a derivation Π , denoted by $d(\Pi)$, is the length of the longest path in Π . The following three results establishes cut elimination.

Lemma 38 (Cut Reduction). The following defines the cut reduction procedure:

- (1) If Π_1 is a derivation of $T \vdash_C \Psi, S^n$ and Π_2 is a derivation of $S \vdash_C \Psi'$ with $c(\Pi_1), c(\Pi_2) \leq |S|$, then there exists a derivation Π of $T \vdash_C \Psi, \Psi'$ with $c(\Pi) \leq |S|$;
- (2) If Π_1 is a derivation of $T \vdash_L \Delta; \Psi, S^n$ and Π_2 is a derivation of $S \vdash_C \Psi'$ with $c(\Pi_1), c(\Pi_2) \leq |S|$, then there exists a derivation Π of $T \vdash_L \Delta; \Psi, \Psi'$ with $c(\Pi) \leq |S|$;
- (3) If Π_1 is a derivation of $B \vdash_L \Delta; \Psi, A^n$ and Π_2 is a derivation of $A \vdash_L \Delta', \Psi'$ with $c(\Pi_1), c(\Pi_2) \leq |S|$, then there exists a derivation Π of $B \vdash_C \Delta, \Delta', \Psi, \Psi'$ with $c(\Pi) \leq |A|$.

Proof. By induction on $d(\Pi_1) + d(\Pi_2)$. We give one case where the last inferences of Π_1 and Π_2 are logical inferences; please see Appendix B.5 for the complete proof.

\bullet — right / \bullet — left. We have

$$\Pi_1 = \frac{\frac{\pi_1}{A \vdash_L \Delta_1; \Psi_1, B_1} \quad \frac{\pi_2}{B_2 \vdash_L \Delta_2; \Psi_2}}{A \vdash_L B_1 \bullet B_2, \Delta_1, \Delta_2; \Psi_1, \Psi_2} \text{LL-}\bullet_R \quad \Pi_2 = \frac{\frac{\pi_3}{B_1 \vdash_L B_2, \Delta; \Psi}}{B_1 \bullet B_2 \vdash_L \Delta; \Psi} \text{LL-}\bullet_L}{A \vdash_L \Delta_1, \Delta_2, \Delta; \Psi_1, \Psi_2, \Psi} \text{LL-cut}$$

reduces to Π

$$\frac{\frac{\pi_1}{A \vdash_L \Delta_1, B_1; \Psi_1} \quad \frac{\pi_3}{B_1 \vdash_L B_2, \Delta; \Psi}}{A \vdash_L \Delta_1, \Delta, B_2; \Psi_1, \Psi} \text{LL-cut} \quad \frac{\pi_2}{B_2 \vdash_L \Delta_2; \Psi_2}}{A \vdash_L \Delta_1, \Delta_2, \Delta; \Psi_1, \Psi_2, \Psi} \text{LL-cut}$$

The resulting derivation Π has cut rank $c(\Pi) = \max(|B_1| + 1, c(\pi_1), c(\pi_2), |B_2| + 1, c(\pi_3)) \leq |B_1 \bullet B_2|$. \square

Lemma 39 (Decrease in Cut-Rank). Let Π be a DLNL^+ proof of a sequent $S \vdash_C \Psi$ or $A \vdash_L \Delta; \Psi$ with $c(\Pi) > 0$. Then there exists a proof Π' of the same sequent with $c(\Pi') < c(\Pi)$.

Proof. By induction on $d(\Pi)$. If the last inference is not a cut, then we apply the induction hypothesis. If the last inference is a cut on a formula A , but A is not of maximal rank among the cut formulas, so that $c(\Pi) > |A| + 1$, then we apply the induction hypothesis. Finally, if the last inference is a cut on A and $c(\Pi) = |A| + 1$ we have the following situation:

$$\Pi = \frac{\frac{\Pi_1}{B \vdash_L \Delta, A; \Psi} \quad \frac{\Pi_2}{A \vdash_L \Delta'; \Psi'}}{B \vdash_L \Delta, \Delta'; \Psi, \Psi'} \text{LL-cut}$$

Now since $c(\Pi_1), c(\Pi_2) \leq |A| + 1$ then by applying the induction hypothesis to the premises of the previous derivation we can construct derivations Π'_1 and Π'_2 with $c(\Pi'_1) \leq |A|$ and $c(\Pi'_2) \leq |A|$. Then by cut reduction we can construct a derivation Π' proving $B \vdash_L \Delta, \Delta'; \Psi, \Psi'$ with $c(\Pi') \leq |A|$ as required. \square

Theorem 40 (Cut Elimination). Let Π be a proof of a sequent $S \vdash_C \Psi$ or $A \vdash_L \Delta; \Psi$ such that $c(\Pi) > 0$. There is an algorithm which yields a cut free proof of the same sequent.

Proof. By induction on $c(\Pi)$ using the previous lemma. \square

3.2. Sequent-style Natural Deduction. The inference rules for the non-linear and linear fragments of the sequent-style natural deduction formalization of DLNL (DND) can be found in Figure 3 and Figure 4 respectively.

Remark 41. In DLNL logic contexts are treated multiplicatively and so are in DND. Non-linear context could also be treated additively. In the case of the minor premises of non-linear disjunction elimination (rule NLL_{+E} of Figure 3 an additive interpretation is required, namely, both minor premises must have the same right context, to match the categorical interpretation of disjunction as coproduct. The same holds for the term assignment in the rule TC_{+E} of Figure 5. Of course additive contexts can be simulated using weakening and contraction. This is what we do in the case of disjunction elimination.

We now recall a correspondence between DND and DLNL logic. First, we need the admissible rule of cut, i.e., substitution.

$$\begin{array}{c}
\frac{}{S \vdash_C S} \text{NC_id} \quad \frac{S \vdash_C \Psi}{S \vdash_C T, \Psi} \text{NC_weak} \quad \frac{S \vdash_C T, T, \Psi}{S \vdash_C T, \Psi} \text{NC_contr} \\
\\
\frac{S \vdash_C 0, \Psi \quad S_1 \vdash_C \Psi_1, \dots, S_n \vdash_C \Psi_n}{S \vdash_C \Psi, \Psi_1, \dots, \Psi_n} \text{NC_0}_E \quad \frac{S \vdash_C \Psi, T_1}{S \vdash_C \Psi, T_1 + T_2} \text{NC_+}_{I_1} \\
\\
\frac{S \vdash_C \Psi, T_2}{S \vdash_C \Psi, T_1 + T_2} \text{NC_+}_{I_2} \quad \frac{S \vdash_C \Psi_1, T_1 + T_2 \quad T_1 \vdash_C \Psi_2 \quad T_2 \vdash_C \Psi_2}{S \vdash_C \Psi_1, \Psi_2} \text{NC_+}_E \\
\\
\frac{S \vdash_C \Psi_1, T_1 \quad T_2 \vdash_C \Psi_2}{S \vdash_C \Psi_1, \Psi_2, T_1 - T_2} \text{NC_+}_I \quad \frac{S \vdash_C \Psi_1, T_1 - T_2 \quad T_1 \vdash_C T_2, \Psi_2}{S \vdash_C \Psi_1, \Psi_2} \text{NC_+}_E \\
\\
\frac{S \vdash_C \Psi_1, \text{HA} \quad A \vdash_L \cdot; \Psi_2}{S \vdash_C \Psi_1, \Psi_2} \text{NC_H}_E
\end{array}$$

Figure 3: Non-linear fragment of DND logic

$$\begin{array}{c}
\frac{}{A \vdash_L A; \cdot} \text{NLL_id} \quad \frac{A \vdash_L \Delta; \Psi}{A \vdash_L \Delta; T, \Psi} \text{NLC_weak} \quad \frac{A \vdash_L \Delta; T, T, \Psi}{A \vdash_L \Delta; T, \Psi} \text{NLC_contr} \\
\\
\frac{A \vdash_L \Delta; \Psi}{A \vdash_L \Delta, \perp; \Psi} \text{NLL_}\perp_I \quad \frac{A \vdash_L \perp, \Delta; \cdot}{A \vdash_L \Delta; \cdot} \text{NLL_}\perp_E \quad \frac{A \vdash_L \Delta, B_1, B_2; \Psi}{A \vdash_L \Delta, B_1 \oplus B_2; \Psi} \text{NLL_}\oplus_I \\
\\
\frac{A \vdash_L \Delta, B_1 \oplus B_2; \Psi \quad B_1 \vdash_L \Delta_1; \Psi_1 \quad B_2 \vdash_L \Delta_2; \Psi_2}{A \vdash_L \Delta, \Delta_1, \Delta_2; \Psi, \Psi_1, \Psi_2} \text{NLL_}\oplus_E \\
\\
\frac{A \vdash_L \Delta_1, B_1; \Psi_1 \quad B_2 \vdash_L \Delta_2; \Psi_2}{A \vdash_L B_1 \bullet B_2, \Delta_1, \Delta_2; \Psi_1, \Psi_2} \text{NLL_}\bullet_I \\
\\
\frac{A \vdash_L \Delta_1, B_1 \bullet B_2; \Psi_1 \quad B_1 \vdash_L B_1, \Delta_2; \Psi_2}{A \vdash_L \Delta_1, \Delta_2; \Psi_1, \Psi_2} \text{NLL_}\bullet_E \quad \frac{A \vdash_L \Delta; T, \Psi}{A \vdash_L \Delta, \text{JT}; \Psi} \text{NLL_J}_I \\
\\
\frac{A \vdash_L \Delta, \text{JT}; \Psi_1 \quad T \vdash_C \Psi_2}{A \vdash_L \Delta; \Psi_1, \Psi_2} \text{NLL_J}_E \quad \frac{A \vdash_L \Delta, B; \Psi}{A \vdash_L \Delta; \text{HB}, \Psi} \text{NLL_H}_I \\
\\
\frac{A \vdash_L \Delta; \Psi_1, \text{HA} \quad A \vdash_L \cdot; \Psi_2}{A \vdash_L \Delta; \Psi_1, \Psi_2} \text{NLL_H}_E
\end{array}$$

Figure 4: Linear fragment of DND logic

Lemma 42 (Admissible Rules in DND). The following rules are admissible in DND:

$$\frac{S \vdash_C \Psi_1, T \quad T \vdash_C \Psi_2}{S \vdash_C \Psi_1, \Psi_2} \text{NC_cut} \qquad \frac{A \vdash_L \Delta; \Psi_1, T \quad T \vdash_C \Psi_2}{A \vdash_L \Delta; \Psi_1, \Psi_2} \text{NLC_cut}$$

$$\frac{A \vdash_L \Delta_1, B; \Psi_1 \quad B \vdash_L \Delta_2; \Psi_2}{A \vdash_L \Delta_1, \Delta_2; \Psi_1, \Psi_2} \text{NLL_cut}$$

Using these admissible rules we can construct a proof preserving translation between DND and DLNL logic.

Lemma 43 (Translations between DND and DLNL logic). There are functions $S : DND \rightarrow DLNL$ and $N : DLNL \rightarrow DND$ from natural deduction to sequent calculus derivations.

Notice that the right rules of the sequent calculus and the introductions of natural deduction have the same form. Elimination rules are derivable from left rules with *cut* and left rules are derivable using the admissible cut rule in DND. For instance, the NC.0_E rule

$$\frac{S \vdash_C 0, \Psi \quad S_1 \vdash_C \Psi_1, \dots, S_n \vdash_C \Psi_n}{S \vdash_C \Psi, \Psi_1, \dots, \Psi_n} \text{NC.0}_E$$

is derivable in the sequent calculus as follows:

$$\frac{\frac{S \vdash_C 0, \Psi \quad 0 \vdash S_1, \dots, S_n}{S \vdash_C \Psi, S_1, \dots, S_n} \text{C_cut} \quad \frac{S_1 \vdash_C \Psi_1}{S \vdash_C \Psi, \Psi_1, S_2, \dots, S_n} \text{C_cut} \quad \vdots}{\frac{S \vdash_C \Psi, \Psi_1, \dots, \Psi_{n-1}, S_n}{S \vdash_C \Psi, \Psi_1, \dots, \Psi_{n-1}, \Psi_n} \text{C_cut} \quad S_n \vdash \Psi_n} \text{C_cut}$$

3.3. Term Assignment. We now turn to giving a term assignment to DND logic called TND, which is greatly influenced by Crolard's term assignment for subtractive logic in the paper *A formulae-as-types interpretation of subtractive logic* JLC 2004. Crolard based his term assignment on Parigot's [16] $\lambda\mu$ -calculus. He then shows that a type theory of coroutines can be given by subtractive types and it is this result we pull inspiration from.

TND pushes beyond Crolard's paper *A formulae-as-types interpretation of subtractive logic* JLC 2004, where he gives a term assignment to bi-intuitionistic logic within a variant of Parigot's [16] classical $\lambda\mu$ -calculus and interprets the term assigned to subtraction in terms of coroutines. He then restricts the calculus to provide a constructive version of subtraction called *safe coroutines*. The second author used a variant of his constructive calculus as a term assignment to co-intuitionistic logic and to linear co-intuitionistic logic [3] without using the $\lambda\mu$ -calculus. In this formulation, distinct terms are assigned to distinct formulas in the context and the reduction of a term in context may impact other terms in the context.

The syntax of TND terms is defined by the following definition.

Definition 44. The syntax for TND terms and typing judgments are given by the following grammar:

(non-linear terms) $s, t ::= x \mid \varepsilon \mid t_1 \cdot t_2 \mid \text{false } t \mid x(t) \mid \text{mkc}(t, x) \mid \text{inl } t \mid \text{inr } t \mid$
 $\text{case } t \text{ of } x.t_1, y.t_2 \mid \text{H } e \mid \text{let } J x = e \text{ in } t \mid \text{postp}(x \mapsto t_1, t_2) \mid$
 $\text{let } H x = t_1 \text{ in } t_2$

(linear terms) $e, u ::= x \mid \text{connect}_\perp \text{ to } e \mid \text{postp}_\perp e \mid \text{postp}(x \mapsto e_1, e_2) \mid \text{mkc}(e, x) \mid x(e) \mid$
 $e_1 \oplus e_2 \mid \text{casel } e \mid \text{caser } e \mid J t$

(non-linear judgment) $x : R \vdash_C \Psi$

(linear judgment) $x : A \vdash_L \Delta; \Psi$

Contexts, Δ and Ψ , are the straightforward extension where each type is annotated with a term from the respective fragment.

To aid the reader in understanding the variable structure, which variable annotations are bound, deployed throughout the TND term syntax we give the definitions of the free variable functions in the following definition.

Definition 45. The free variable functions, $FV(t)$ and $FV(e)$, for linear and non-linear terms t and e are defined by mutual recursion as follows:

linear terms:

$FV(x) = \{x\}$
 $FV(\text{connect}_\perp \text{ to } e) = FV(e)$
 $FV(x(e)) = FV(e)$
 $FV(\text{mkc}(e, y)) = FV(e)$
 $FV(e_1 \oplus e_2) = FV(e_1) \cup FV(e_2)$
 $FV(\text{casel } e) = FV(e)$
 $FV(\text{caser } e) = FV(e)$
 $FV(J t) = FV(t)$

non-linear terms:

$FV(x) = \{x\}$
 $FV(\varepsilon) = \emptyset$
 $FV(t_1 \cdot t_2) = FV(t_1) \cup FV(t_2)$
 $FV(\text{false } t) = FV(t)$
 $FV(x(t)) = FV(t)$
 $FV(\text{mkc}(t, y)) = FV(t)$
 $FV(\text{inl } t) = FV(\text{inr } t) = FV(t)$
 $FV(\text{case } t_1 \text{ of } x.t_2, y.t_3) =$
 $FV(t_1) \cup FV(t_2) \setminus \{x\} \cup FV(t_3) \setminus \{y\}$
 $FV(\text{let } J y = e \text{ in } t) = FV(e) \cup FV(t) \setminus \{y\}$
 $FV(\text{let } H y = t_1 \text{ in } t_2) = FV(t_1) \cup FV(t_2) \setminus \{y\}$
 $FV(H e) = FV(e)$

The free variables of a p -term are defined as follows:

$$\begin{aligned} FV(\text{postp}_\perp e) &= FV(e) \\ FV(\text{postp}(x \mapsto e_1, e_2)) &= FV(e_1) \setminus \{x\} \cup FV(e_2) \end{aligned}$$

and similarly for terms $\text{postp}(x \mapsto t_1, t_2)$.

Terms are then typed by annotating the previous term structure over DND derivations, and this is accomplished by annotating the DND inference rules. The typing rules for the non-linear fragment of TND can be found in Figure 5, and the typing rules for the linear fragment of TND can be found in Figure 6.

Remark 46. Let us call terms of the form $\text{postp}(x \mapsto t_1, t_2)$, $\text{postp}(x \mapsto e_1, e_2)$, and $\text{postp}_\perp e$ p -terms. Then say that a term t is p -normal if t does not contain any p -term as a proper subterm. In a typed calculus, linear p -terms can be typed with \perp . Non-linear p terms can be typed with 0: in presence of the TC_0_E rule this yields instances of the *ex falso* rule. This is what happens in Crolard's calculus, where the analogue of the $\text{postp}(x \mapsto t_1, t_2)$, namely, *resume t_2 with $x \mapsto t_1$* , always goes with a *weakening* operation. The term ε is the identity of the contraction binary operator $t_1 \cdot t_2$.

$$\begin{array}{c}
\frac{}{x : S \vdash_C x : S} \text{TC_id} \quad \frac{x : S \vdash_C \Psi}{x : S \vdash_C \Psi, \varepsilon : T} \text{TC_weak} \quad \frac{x : S \vdash_C t_1 : T, t_2 : T, \Psi}{x : S \vdash_C (t_1 \cdot t_2) : T, \Psi} \text{TC_contr} \\
\\
\frac{x : S \vdash_C t : 0, \Psi \quad x_1 : S_1 \vdash_C \Psi_1, \dots, x_n : S_n \vdash_C \Psi_n}{x : S \vdash_C \Psi, [\text{false } t/x_1]\Psi_1, \dots, [\text{false } t/x_n]\Psi_n} \text{TC_0}_E \\
\\
\frac{x : S \vdash_C \Psi, t : T_1}{x : S \vdash_C \Psi, \text{inl } t : T_1 + T_2} \text{TC_+}_{I_1} \quad \frac{x : S \vdash_C \Psi, t : T_2}{x : S \vdash_C \Psi, \text{inr } t : T_1 + T_2} \text{TC_+}_{I_2} \\
\\
\frac{x : S \vdash_C \Psi_1, t : T_1 + T_2 \quad y : T_1 \vdash_C \Psi_2 \quad z : T_2 \vdash_C \Psi_3 \quad |\Psi_2| = |\Psi_3|}{x : S \vdash_C \Psi_1, \text{case } t \text{ of } y.\Psi_2, z.\Psi_3} \text{TC_+}_E \\
\\
\frac{x : S \vdash_C \Psi_1, t : T_1 \quad y : T_2 \vdash_C \Psi_2}{x : S \vdash_C \Psi_1, \text{mkc}(t, y) : T_1 - T_2, [y(t)/y]\Psi_2} \text{TC_}_I \\
\\
\frac{x : S \vdash_C \Psi_1, s : T_1 - T_2 \quad y : T_1 \vdash_C t : T_2, \Psi_2}{x : S \vdash_C \Psi_1, \text{postp}(y \mapsto t, s), [y(s)/y]\Psi_2} \text{TC_}_E \\
\\
\frac{x : S \vdash_C \Psi_1, t : \text{HA} \quad y : A \vdash_L \cdot; \Psi_2}{x : S \vdash_C \Psi_1, \text{let } H y = t \text{ in } \Psi_2} \text{TC_H}_E
\end{array}$$

Figure 5: Non-linear fragment of the term assignment for TND

However when within a non- p -normal term an expression of the form $\text{postp}(x \mapsto t_1, t_2)$ is eliminated as a β redex, there is a choice of the syntax for the contextual reduction. In absence of a more detailed analysis of the matter, we prefer to leave the typing of p terms implicit in the syntax, to enforce the requirement of p -normality and to use the admissible rule of weakening in place of the TC_0_E rule in this context.

The typing rules depend on the extension of let and case expressions to typing contexts. We use the following notation for *parallel composition* of typing contexts:

$$\Delta = e_1 : A_1 \parallel \dots \parallel e_n : A_n$$

This operation should be regarded as associative, commutative and having the empty context as its identity. The extension of let expressions to contexts is given as follows:

$$\begin{aligned}
\text{let } p = t \text{ in } \cdot &= \cdot \\
\text{let } p = t_1 \text{ in } (t_2 : A) &= \text{let } p = t_1 \text{ in } t_2 : A \\
\text{let } p = t \text{ in } (\Psi_1 \parallel \Psi_2) &= (\text{let } p = t \text{ in } \Psi_1) \parallel (\text{let } p = t \text{ in } \Psi_2)
\end{aligned}$$

where $p = H y$ or $p = J y$. Case expressions are handled similarly.

Similarly to DND logic we have the following admissible rules.

$$\begin{array}{c}
\frac{}{x : A \vdash_L x : A; \cdot} \text{TLL_id} \qquad \frac{x : A \vdash_L \Delta; \Psi}{x : A \vdash_L \Delta; \Psi, \varepsilon : T} \text{TL_weak} \\
\\
\frac{x : A \vdash_L \Delta; t_1 : T, t_2 : T, \Psi}{x : A \vdash_L \Delta; t_1 \cdot t_2 : T, \Psi} \text{TLC_contr} \qquad \frac{x : A \vdash_L \Delta; \Psi \quad e : B \in \Delta}{x : A \vdash_L \Delta, \text{connect}_\perp \text{ to } e : \perp; \Psi} \text{TLL_}\perp_I \\
\\
\frac{x : A \vdash_L e : \perp, \Delta; \cdot}{x : A \vdash_L \text{postp}_\perp e, \Delta; \cdot} \text{TLL_}\perp_E \qquad \frac{x : A \vdash_L \Delta, e_1 : B_1, e_2 : B_2; \Psi}{x : A \vdash_L \Delta, e_1 \oplus e_2 : B_1 \oplus B_2; \Psi} \text{TLL_}\oplus_I \\
\\
\frac{x : A \vdash_L \Delta, e : B_1 \oplus B_2; \Psi \quad y : B_1 \vdash_L \Delta_1; \Psi_1 \quad z : B_2 \vdash_L \Delta_2; \Psi_2}{x : A \vdash_L \Delta, [\text{case}_l(e)/y]\Delta_1, [\text{case}_r(e)/z]\Delta_2; \Psi, [\text{case}_l(e)/y]\Psi_1, [\text{case}_r(e)/z]\Psi_2} \text{TLL_}\oplus_E \\
\\
\frac{x : A \vdash_L \Delta_1, e : B_1; \Psi_1 \quad y : B_2 \vdash_L \Delta_2; \Psi_2}{x : A \vdash_L \text{mkc}(e, y) : B_1 \bullet B_2, \Delta_1, [y(e)/y]\Delta_2; \Psi_1, [y(e)/y]\Psi_2} \text{TLL_}\bullet_I \\
\\
\frac{x : A \vdash_L \Delta_1, e_1 : B_1 \bullet B_2; \Psi_1 \quad y : B_1 \vdash_L e_2 : B_1, \Delta_2; \Psi_2}{x : A \vdash_L \Delta_1, \text{postp}(y \mapsto e_2, e_1), \Delta_2; \Psi_1, \Psi_2} \text{TLL_}\bullet_E \\
\\
\frac{x : A \vdash_L \Delta; t : T, \Psi}{x : A \vdash_L \Delta, \text{J } t : \text{J } T; \Psi} \text{TLL_J}_I \qquad \frac{x : A \vdash_L \Delta, e : \text{J } T; \Psi_1 \quad y : T \vdash_C \Psi_2}{x : A \vdash_L \Delta; \Psi_1, \text{let J } y = e \text{ in } \Psi_2} \text{TLL_J}_E \\
\\
\frac{x : A \vdash_L \Delta, e : B; \Psi}{x : A \vdash_L \Delta; \text{H } e : \text{H } B, \Psi} \text{TLL_H}_I \qquad \frac{x : A \vdash_L \Delta; \Psi_1, t : \text{H } A \quad y : A \vdash_L \cdot; \Psi_2}{x : A \vdash_L \Delta; \Psi_1, \text{let H } y = t \text{ in } \Psi_2} \text{TLL_H}_E
\end{array}$$

Figure 6: Linear fragment of the term assignment for TND

Lemma 47 (Admissible Typing Rules). The term assignment for the admissible rules of the calculus is as follows:

$$\begin{array}{c}
\frac{x : S \vdash_C \Psi_1, t : T \quad y : T \vdash_C \Psi_2}{x : S \vdash_C \Psi_1, [t/y]\Psi_2} \text{TC_cut} \qquad \frac{x : A \vdash_L \Delta; \Psi_1, t : T \quad y : T \vdash_C \Psi_2}{x : A \vdash_L \Delta; \Psi_1, [t/y]\Psi_2} \text{NLC_cut} \\
\\
\frac{x : A \vdash_L \Delta_1, e : B; \Psi_1 \quad y : B \vdash_L \Delta_2; \Psi_2}{x : A \vdash_L \Delta_1, [e/y]\Delta_2; \Psi_1, [e/y]\Psi_2} \text{TLL_cut}
\end{array}$$

We generalize the rule of contraction on the non-linear side to contexts. Let m_1 and m_2 be multisets of terms, then we denote by $m_1 \cdot m_2$ the sum of multisets; if multisets are represented as lists, then the sum is representable as the appending of the lists. We denote singleton multisets, $\{t\}$, by the term that inhabits it, e.g. t . We extend this to contexts, $\Psi_1 \cdot \Psi_2$, recursively as follows:

$$\begin{aligned}
(\cdot) \cdot (\cdot) &= (\cdot) \\
(t_1 : S) \cdot (t_2 : S) &= t_1 \cdot t_2 : S \\
(\Psi_1 \parallel \Psi_3) \cdot (\Psi_2 \parallel \Psi_4) &= (\Psi_1 \cdot \Psi_2) \parallel (\Psi_3 \cdot \Psi_4)
\end{aligned}$$

where $|\Psi_1| = |\Psi_3|$ and $|\Psi_2| = |\Psi_4|$.

At this point we are now ready to turn to computing in TND by specifying the reduction relation. This definition is perhaps the most interesting aspect of the theory, because reducing one term may affect others.



Figure 7: Reductions for Linear Terms

β -Reduction in TND. As we discussed above cointuitionistic logic can be interpreted as a theory of coroutines that manipulate local context. Thus, reducing one term in a typing context could affect other terms in the context. This implies that the definition of the reduction relation for TND must account for more than a single term. We accomplish this by defining the reduction relation of terms in context, $x : S \vdash_C \Psi_1, t : T, \Psi_2$ and $x : A \vdash_{\perp} \Delta_1, e : B, \Delta_2; \Psi$, so that the manipulation of the context is made explicit.

The reduction rules for the linear and non-linear fragments can be found in Figure 7 and Figure 8 respectively. We denote the judgments for reduction by $x : S \vdash_C \Psi_1 \rightsquigarrow x : S \vdash_C \Psi_2$ and $x : A \vdash_{\perp} \Delta_1; \Psi_1 \rightsquigarrow x : A \vdash_{\perp} \Delta_2; \Psi_2$. In the interest of readability we do not show full derivations, but it should be noted that it is assumed that every term mentioned in a reduction rule is typable with the expected type given where it occurs in the judgment. Furthermore, the reduction relation depends on a few standard definitions and non-standard binding operations.

The non-standard binding operations concern the variable y in $\text{mkc}(t, y)$ and in $\text{postp}(y \mapsto t, s)$ and the related expressions $y(t)$ and $y(s)$, respectively, occurring in the non-linear context; similar operations occur in the linear case. Consider term assignment to the rule subtraction introduction TC_{-I} in Figure 5. The variable y is the unique free variable occurring in the sequent $y : T_2 \vdash_C \Psi_2$, the minor premise of the inference. In the conclusion $x : S \vdash_{\Psi_1}, (t, y) : T_1 - T_2, [y(t)/y] \Psi_2$ the variable y is bound in $\text{mkc}(t, y)$; moreover, the occurrences of the free variable y have been substituted simultaneously in the context Ψ by the expression $y(t)$ which denotes a bound variable, indexed



Figure 8: Reductions for Non-linear Terms

with t . Similar explanations apply to the term assignment for subtraction elimination, and to the corresponding linear rules in Figure 6.

An analogue of the capture of a free variable by a binder in the λ -calculus, is an occurrence of a bound variable $y(t)$ whose binder is ambiguous, for instance in a context where there were two



Figure 9: Reductions for Non-linear Terms Continued

occurrences of $\text{mkc}(t, y)$, as a result of a contraction/cut reduction in a derivation. Such a context may be the conclusion of the following derivation, if $x_1 = x_2 \text{ m } y_1 = y_2$; here $t_1 = \text{false } x_1$, $t_2 = \text{false } x_2$:

$$\frac{z : 0 \vdash_C z : 0 \quad \frac{x_1 : S \vdash_C x_1 : S \quad y_1 : T \vdash_C y_1 : T}{x_1 : S \vdash_C \text{mkc}(x_1, y_1) : S - T, y_1(x_1) : T} \quad \frac{x_2 : A \vdash_C x_2 : A \quad y_2 : B \vdash_C y_2 : B}{x_2 : S \vdash_C \text{mkc}(x_2, y_2) : S - T, y_2(x_2) : T}}{z : 0 \vdash_C \text{mkc}(t_1, y_1) : S - T, \text{mkc}(t_2, y_2) : S - T, y_1(t_1) : T, y_2(t_2) : T}$$

A formal notion of α conversion has been proposed for this notion of binding in untyped linear contexts in [3]. Here (capture-avoiding) substitution, denoted by $[t_1/x]t_2$, $[e/x]t$, $[t/x]e$, and $[e_1/x]e_2$, is defined in the usual way. We extend capture-avoiding substitution to multisets in the following way:

- $[t_1 \cdot \dots \cdot t_n/z]s = [t_1/z]s \cdot \dots \cdot [t_n/z]s$
- $[t_1 \cdot \dots \cdot t_n/z]p = [t_1/z]p \parallel \dots \parallel [t_n/z]p$, where p is a p -term

The extension of the other flavors of substitution to multisets are similar. Standard extension of substitution to contexts was also necessary.

Finally, there are several commuting conversions that are required for reduction, for example, the following is one:

$$\frac{\frac{x : S \vdash_C \Psi_1, t : T_2 + T_3 \quad z : T_3 \vdash_C \Psi_3, t_2 : T_4 + T_5}{x : S \vdash_C \Psi_1, \text{case } t \text{ of } y.t_1, z.t_2 : T_4 + T_5} \quad v_1 : T_4 \vdash_C \Psi_4 \quad v_2 : T_5 \vdash_C \Psi_5}{x : S \vdash_C \Psi_1, \text{case } (\text{case } t \text{ of } y.t_1, z.t_2) \text{ of } v_1.\Psi_4, v_2.\Psi_5}$$

commutes to

$$\frac{x : S \vdash_{\mathcal{C}} \Psi_1, t : T_2 + T_3 \quad \Pi_1 \quad \Pi_2}{x : S \vdash_{\mathcal{C}} \Psi_1, \text{case } t \text{ of } y_2.(\Psi_2, \text{case } t_1 \text{ of } v_1.\Psi_4, v_2.\Psi_5), y_3.(\Psi_3, \text{case } t_2 \text{ of } v_1.\Psi_4, v_2.\Psi_5)}$$

where

$$\Pi_1 : \frac{y_2 : T_2 \vdash_{\mathcal{C}} \Psi_2, t_1 : T_4 + T_5 \quad v_1 : T_4 \vdash_{\mathcal{C}} \Psi_4 \quad v_2 : T_5 \vdash_{\mathcal{C}} \Psi_5}{y_2 : T_2 \vdash_{\mathcal{C}} \Psi_2, \text{case } t_1 \text{ of } v_1.\Psi_4, v_2.\Psi_5}$$

$$\Pi_2 : \frac{y_3 : T_3 \vdash_{\mathcal{C}} \Psi_3, t_2 : T_4 + T_5 \quad v_1 : T_4 \vdash_{\mathcal{C}} \Psi_4 \quad v_2 : T_5 \vdash_{\mathcal{C}} \Psi_5}{y_3 : T_3 \vdash_{\mathcal{C}} \Psi_3, \text{case } t_2 \text{ of } v_1.\Psi_4, v_2.\Psi_5}$$

If $t_1 = \text{inl } s_1$ and $t_2 = \text{inr } s_2$ then after commutation

$$y_2 : T_2 \vdash_{\mathcal{C}} \Psi_2, \text{case } (\text{inl } s_1) \text{ of } v_1.\Psi_4, v_2.\Psi_5 \rightsquigarrow_{\beta} y_2 : T_2 \vdash_{\mathcal{C}} \Psi_2, [s_1/v_1]\Psi_4$$

and

$$y_3 : T_3 \vdash_{\mathcal{C}} \Psi_3, \text{case } (\text{inr } s_2) \text{ of } v_1.\Psi_4, v_2.\Psi_5 \rightsquigarrow_{\beta} y_2 : T_2 \vdash_{\mathcal{C}} \Psi_3, [s_2/v_2]\Psi_5$$

There are other commuting conversions as well, but as one can see, due to the complexities introduced in reduction arising from the fact that multiple terms in the context are affected during reduction results in the commuting conversions from being very compact. The remainder of the commuting conversions can be found in Appendix A. In the next section we give the interpretation of TND into the categorical model.

3.4. Categorical interpretation of rules. We now turn to the interpretation of Dual LNL Logic into our categorical model given in Section 2. We structure the proof similarly to Bierman [5], but the proof itself follows similarly to Benton's [4] proof for LNL Logic.

Given a *signature* \mathbf{Sg} , consisting of a collection of types σ_i , where $\sigma_i = A$ or S , and a collection of *sorted function symbols* $f_j : \sigma_1, \dots, \sigma_n \rightarrow \tau$ and given a Symmetric Monoidal Category (SMC) $(\mathcal{C}, \bullet, 1, \alpha, \lambda, \rho, \gamma)$, a *structure* \mathcal{M} for \mathbf{Sg} is an assignment of an object $\llbracket \sigma \rrbracket$ of \mathcal{C} for each type σ and of a morphism $\llbracket f \rrbracket : \llbracket \sigma_1 \rrbracket \bullet \dots \bullet \llbracket \sigma_n \rrbracket \rightarrow \llbracket \tau \rrbracket$ for each function $f : \sigma_1, \dots, \sigma_n \rightarrow \tau$ of \mathbf{Sg} .¹ The types of terms in context $\Delta = [e_1 : A_1, \dots, e_n : A_n]$ or $\Delta = [t_1 : T_1, \dots, t_n : T_n]$ are interpreted into the SMC as $\llbracket \sigma_1, \sigma_2, \dots, \sigma_n \rrbracket = (\dots (\llbracket \sigma_1 \rrbracket \bullet \llbracket \sigma_2 \rrbracket) \dots) \bullet \llbracket \sigma_n \rrbracket$; left associativity is also intended for concatenations of type sequences Γ, Δ . Thus, we need the “book-keeping” functions $\text{Split}(\Gamma, \Delta) : \llbracket \Gamma, \Delta \rrbracket \rightarrow \llbracket \Gamma \rrbracket \bullet \llbracket \Delta \rrbracket$ and $\text{Join}(\Gamma, \Delta) : \llbracket \Gamma \rrbracket \bullet \llbracket \Delta \rrbracket \rightarrow \llbracket \Gamma, \Delta \rrbracket$ inductively defined using the associativity laws α and its inverse α^{-1} (cfr Bierman 1994, Bellin 2015).

The semantics of terms in context is then specified by induction on terms:

$$\llbracket x : A \vdash_{\mathcal{L}} x : A \rrbracket =_{df} id_{\llbracket \sigma \rrbracket}$$

$$\llbracket x : A \vdash_{\mathcal{L}} f(e_1, \dots, e_n) : B \rrbracket =_{df} \llbracket x : A \vdash e_1 : A_1 \rrbracket \bullet \dots \bullet \llbracket x : \sigma \vdash e_n : A_n \rrbracket; \llbracket f \rrbracket$$

and similarly with non-linear types. Following this one then proves by induction on the type derivation that substitution in the term calculus corresponds to composition in the category ([5], Lemma 13).

¹ In this subsection only we use the symbol \bullet and 1 for the monoidal binary operation and its unit in the categorical structure, distinguished from the \oplus and \perp symbols in the formal language. We shall show that the interpretation of \oplus is isomorphic to the operation \bullet , so we shall be able to identify them (and similarly for \perp and 1).

In the mixed sequents $x : A \vdash_L \Delta; \Psi, t : T$ of TND non-linear terms are interpreted through the functor $J : \mathcal{C} \rightarrow \mathcal{L}$. Thus, we have the following:

$$x : A \vdash_L \Delta; \Psi, t : T = x : A \vdash_L \Delta, J\Psi, Jt : JT; \cdot$$

Let \mathcal{M} be a structure for a signature Sg in a SMC \mathcal{L} . Equations in context will be denoted by $x : A \vdash_L \Gamma, e_1 = e_2 : B; \Psi$ and $x : S \vdash_C \Psi, t_1 = t_2 : T$, and are both defined to be the reflexive, symmetric, and transitive closure of the reduction relations defined by the rules in Figure 7 and Figure 8 respectively. Given such an equation:

$$x : A \vdash_L \Gamma, e_1 = e_2 : B; \Psi$$

we say that the structure *satisfies* the equation if it assigns the same morphisms to $x : A \vdash_L \Gamma, e_1 : B; \Psi$ and to $x : A \vdash_L \Gamma, e_2 : B; \Psi$. Similarly, \mathcal{M} satisfies $x : S \vdash_C \Psi, t_1 = t_2 : T$ if it assigns the same morphism to $x : S \vdash_C \Psi, t_1 : T$ and to $x : S \vdash_C \Psi, t_2 : T$. Then given an algebraic theory $\text{Th} = (\text{Sg}, \text{Ax})$, a structure \mathcal{M} for Sg is a *model* for Th if it satisfies all the axioms in Ax .

We now go through some cases of the rules in TND to specify their categorical interpretation so as to satisfy the equations in context and to prove consistency of TND, and hence, DLNL logic in the model. We do not give every case, but the ones we do not give are similar to the ones given here. We analyze the linear connectives, giving an argument for co-ILL that is analogue to Bierman's for ILL. We conclude that as expected:

- the cotensor *par* can be identified with the bifunctor \bullet of the structure;
- linear subtraction $\bullet -$ is the left adjoint to the bifunctor \bullet ;
- the unit \perp can be identified with 1.

3.4.1. *Linear Disjunction.* The introduction rule for Par is of the form

$$\frac{x : A \vdash_L \Delta, e_1 : B, e_2 : C; \Psi}{x : A \vdash_L \Delta, e_1 \oplus e_2 : B \oplus C; \Psi} \text{TLL-}\oplus_I$$

This suggests an operation on Hom-sets of the form:

$$\Phi_{A, \Delta, J\Psi} : \mathcal{L}(A, \Delta \bullet (B \bullet C) \bullet J\Psi) \rightarrow \mathcal{L}(A, \Delta \bullet B \oplus C \bullet J\Psi)$$

natural in Δ, A and $J\Psi$. Given $e : A \rightarrow \Delta \bullet (B \bullet C) \bullet J\Psi$, $a : A' \rightarrow A$, $h : \Delta \rightarrow \Delta'$, and $p : J\Psi \rightarrow J\Psi'$, naturality yields:

$$\Phi_{A', \Delta', J\Psi'}(a; e; h \bullet (id_B \bullet id_C) \bullet p) = a; \Phi_{A, \Delta, J\Psi}(e); h \bullet id_{B \oplus C} \bullet p$$

In particular, suppose we have $d : A \rightarrow \Delta \bullet (B \bullet C) \bullet J\Psi$, and let $e = id_\Delta \bullet (id_B \bullet id_C) \bullet id_{J\Psi}$, $h = id_\Delta$, and $p = id_{J\Psi}$. Then we have $\Phi_{A, \Delta, J\Psi}(d) = d$; $\Phi_{(\Delta \bullet (B \bullet C) \bullet J\Psi), \Delta, J\Psi}(id_\Delta \bullet (id_B \bullet id_C) \bullet id_{J\Psi})$. By functoriality of \bullet we have $id_B \bullet id_C = id_{B \bullet C}$. Hence, writing \bigoplus for $\Phi_{(\Delta \bullet (B \bullet C) \bullet J\Psi), \Delta, J\Psi}(id_\Delta \bullet id_{B \bullet C} \bullet id_{J\Psi})$ we have $\Phi_{A, \Delta, J\Psi}(d) = d; \bigoplus$. Finally, given the morphism $\psi_{\Delta, B, C, P} : ((\Delta \bullet B) \bullet C) \bullet J\Psi \rightarrow \Delta \bullet (B \bullet C) \bullet J\Psi$, which is natural in all arguments and is definable using Split and Join, we define:

$$\llbracket x : A \vdash_L \Delta, e_1 \oplus e_2 : B \oplus C, J\Psi; \cdot \rrbracket =_{df} \llbracket x : A \vdash_L \Delta, e_1 : B, e_2 : C, J\Psi; \cdot \rrbracket; \psi; \bigoplus.$$

The Par elimination rule has the form

$$\frac{z : A \vdash_L \Delta_1, e : B \oplus C; \Psi_1 \quad x : B \vdash_L \Delta_2; \Psi_2 \quad y : C \vdash_L \Delta_3; \Psi_3}{z : A \vdash_L \Delta_1, [\text{case}_l(e)/x]\Delta_2, [\text{case}_r(e)/y]\Delta_3; \Psi_1, [\text{case}_l(e)/x]\Psi_2, [\text{case}_r(e)/y]\Psi_3} \text{TLL-}\oplus_E$$

This suggests an operation on Hom-sets of the form

$$\Psi_{A,\Delta,J\Psi} : \mathcal{L}(A, B \oplus C \bullet \Delta_1 \bullet J\Psi_1) \times \mathcal{L}(B, \Delta_2 \bullet J\Psi_2) \times \mathcal{L}(C, \Delta_3 \bullet J\Psi_3) \rightarrow \mathcal{L}(A, \Delta \bullet J\Psi)$$

natural in $A, \Delta, J\Psi$ where we write $\Delta = \Delta_1 \bullet \Delta_2 \bullet \Delta_3$ and $J\Psi = J\Psi_1 \bullet J\Psi_2 \bullet J\Psi_3$. Given the following morphisms:

$$\begin{array}{lll} g : A \rightarrow B \oplus C \bullet \Delta_1 \bullet J\Psi_1 & d_1 : \Delta_1 \rightarrow \Delta'_1 & p_1 : J\Psi_1 \rightarrow J\Psi'_1 \\ e : B \rightarrow \Delta_2 \bullet J\Psi_2 & d_2 : \Delta_2 \rightarrow \Delta'_2 & p_2 : J\Psi_2 \rightarrow J\Psi'_2 \\ f : C \rightarrow \Delta_3 \bullet J\Psi_3 & d_3 : \Delta_3 \rightarrow \Delta'_3 & p_3 : J\Psi_3 \rightarrow J\Psi'_3 \\ a : A' \rightarrow A \end{array}$$

naturality yields:

$$\Psi_{A',\Delta',J\Psi'}((a; g; id_{B \oplus C} \bullet d_1 \bullet p_1), (e; d_2 \bullet p_2), (f; d_3 \bullet p_3)) = a; \Psi_{A,\Delta,J\Psi}(g, e, f); d_1 \bullet d_2 \bullet d_3 \bullet p_1 \bullet p_2 \bullet p_3; \text{Join}(\Delta', J\Psi').$$

In particular, set $e = id_B, f = id_C, a = id_A, d_i = id_{\Delta_i}$, and $p_i = id_{J\Psi_i}$, and we get

$$\Psi_{A,\Delta,J\Psi}(g, e, f) = \Psi_{A,\Delta,J\Psi}(g, id_B, id_C); id_{\Delta} \bullet c \bullet d; \text{Join}(\Delta, J\Psi)$$

where the operation Join implements the required associativity. Writing $(x)^*$ for $\Psi_{D,\Delta}(x, id_B, id_C)$ we define

$$\begin{aligned} \llbracket z : A \vdash_L \Delta_1, [\text{casel } e/x] \Delta_2, [\text{caser } e/y] \Delta_3; \Psi_1, [\text{casel } e/x] \Psi_2, [\text{caser } e/y] \Psi_3 \rrbracket &=_{df} \\ \llbracket z : A \vdash_L \Delta_1, e : B \oplus C; \Psi_1 \rrbracket^*; (id_{\Delta_1} \bullet \llbracket x : B \vdash_L \Delta_2; \Psi_2 \rrbracket \bullet \llbracket y : C \vdash_L \Delta_3; \Psi_3 \rrbracket); \text{Join}(\Delta, J\Psi). \end{aligned}$$

We now turn to the equations in context. Consider the following case:

$$\frac{\begin{array}{lll} e \equiv \text{casel}(e_1 \oplus e_2) & & \\ e' \equiv \text{caser}(e_1 \oplus e_2) & & \\ |\Delta_1| = |\Delta'_1| & |\Psi_1| = |\Psi'_1| & y : A_2 \vdash_L \Delta_3; J\Psi_3 = \Delta'_3; J\Psi'_3 \\ |\Delta_2| = |\Delta'_2| & |\Psi_2| = |\Psi'_2| & x : A_1 \vdash_L \Delta_2; J\Psi_2 = \Delta'_2; J\Psi'_2 \\ |\Delta_3| = |\Delta'_3| & |\Psi_3| = |\Psi'_3| & z : B \vdash_L e_1 : A_1, e_2 : A_2, \Delta_1; J\Psi_1 = e'_1 : A_1, e'_2 : A_2, \Delta'_1; J\Psi'_1 \end{array}}{z : B \vdash_L \Delta_1, [e/x] \Delta_2, [e'/x] \Delta_3; \Psi_1, [e/x] \Psi_2, [e'/x] \Psi_3 = \Delta'_1, [e'_1/x] \Delta_2, [e'_2/x] \Delta_3; \Psi_1, [e'_1/x] \Psi'_2, [e'_2/x] \Psi'_3} \oplus \beta$$

Let

$$q : B \rightarrow A_1 \bullet A_2 \bullet \Delta_1 \bullet J\Psi, \quad m : A_1 \rightarrow \Delta_2 \bullet J\Psi_2 \quad \text{and} \quad n : A_2 \rightarrow \Delta_3 \bullet J\Psi_3.$$

Then to satisfy the above equations in context we need that the following diagram commutes:

$$\begin{array}{ccc} B \xrightarrow{q} \Delta_1 \bullet J\Psi_1 \bullet (A_1 \bullet A_2) & \xrightarrow{id_{\Delta_1} \bullet m \bullet n} & \Delta_1 \bullet J\Psi_1 \bullet \Delta_2 \bullet J\Psi_2 \bullet \Delta_3 \bullet J\Psi_3 \\ \downarrow \oplus & & \uparrow id_{\Delta_1} \bullet m \bullet n \\ \Delta_1 \bullet J\Psi_1 \bullet A \oplus B & \xrightarrow{*} & \Delta_1 \bullet J\Psi_1 \bullet A \bullet B \end{array}$$

We make the assumption that the above decomposition is unique. Moreover, supposing Δ_1 to be empty and $m = id_A, n = id_B, q = id_A \bullet id_B = id_{A \bullet B}$ we obtain $(id_A \bullet id_B; \oplus)^* = id_A \bullet id_B$ and similarly $(id_{A \oplus B})^*; \oplus = id_{A \oplus B}$; hence we may conclude that there is a natural isomorphism

$$\frac{D \rightarrow \Gamma \bullet A \bullet B}{D \rightarrow \Gamma \bullet A \oplus B}$$

so we can identify \bullet and \oplus . Finally we see that the following η equation in context is also satisfied:

$$\frac{\begin{array}{c} \oplus - \eta \text{ rule} \\ |\Delta| = |\Delta'| \quad |\Psi| = |\Psi'| \quad z : B \vdash_L \Delta; \Psi = \Delta'; \Psi' \end{array}}{z : B \vdash_L (\text{casel } e \oplus \text{caser } e) : A_1 \oplus A_2, \Delta; \Psi = e : A_1 \oplus A_2, \Delta'; \Psi'}$$

(3.1)

3.4.2. *Linear subtraction.* 3.4.2.1. *Subtraction introduction.* The introduction rule for subtraction has the form:

$$\frac{x : A \vdash_L \Delta_1, e : B; \Psi_1 \quad y : C \vdash_L \Delta_2; \Psi_2 \quad |\Psi_1| = |\Psi_2|}{x : A \vdash_L \Delta_1, \text{mkc}(e, y) : B \multimap C, [y(e)/y]\Delta_2; \Psi_1, [y(e)/y]\Psi_2} \text{TLL}_{\multimap-I}$$

This suggests a natural transformation with components:

$$\Phi_{A, \Delta, J\Psi} : \mathcal{L}(A, \Delta_1 \bullet B \bullet J\Psi_1) \times \mathcal{L}(C, \Delta_2 \bullet J\Psi_2) \rightarrow \mathcal{L}(A, \Delta_1 \bullet (B \multimap C) \bullet \Delta_2 \bullet J\Psi_1 \bullet J\Psi_2)$$

natural in $A, \Delta_1, \Delta_2, J\Psi_1, J\Psi_2$. Taking morphisms

$$e : A \rightarrow \Delta_1 \bullet B \bullet J\Psi_1, \quad f : C \rightarrow \Delta_2 \bullet J\Psi_2$$

and also $a : A' \rightarrow A, d_1 : \Delta_1 \rightarrow \Delta'_1, d_2 : \Delta_2 \rightarrow \Delta'_2, p_1 : J\Psi_1 \rightarrow J\Psi'_1, p_2 : J\Psi_2 \rightarrow J\Psi'_2$, by naturality we have

$$\begin{aligned} & \Phi_{A', \Delta'_1, \Delta'_2, J\Psi'_1, J\Psi'_2}((a; e; d_1 \bullet \text{id}_B \bullet p_1), (f; d_2; p_2)) = \\ & = a; \Phi_{A, \Delta, J\Psi}(e, f); d_1 \bullet d_2 \bullet \text{id}_{B \multimap C}; \text{Join}(\Delta'_1, \Delta'_2, B \multimap C, J\Psi'_1, J\Psi'_2) \end{aligned}$$

In particular, taking $a = \text{id}_A, d_1 = \text{id}_{\Delta_1}, p_1 = \text{id}_{J\Psi_1}, p_2 = \text{id}_{J\Psi_2}$ but $d_2 : C \rightarrow \Delta_2 \bullet J\Psi_2$ and $f = \text{id}_C$ we have:

$$\begin{aligned} \Phi_{A, \Delta_1, \Delta_2, J\Psi_1, J\Psi_2}(e, d_2) &= \Phi_{A, \Delta_1}(e, \text{id}_C); \text{id}_{\Delta_1} \bullet d_2 \bullet \text{id}_{A \multimap B} \bullet \text{id}_{J\Psi_1} \bullet \text{id}_{J\Psi_2}; \\ & \quad \text{Join}(\Delta_1, \Delta_2, A \multimap B, J\Psi_1, J\Psi_2) \end{aligned}$$

Writing $\text{MKC}_{A, \Delta_1, J\Psi_1}^C(e)$ for $\Phi_{A, \Delta_1, J\Psi_1}(e, \text{id}_C)$, $\Phi_{A, \Delta, J\Psi}(e, d_2)$ can be expressed as the composition

$$\text{MKC}_{A, \Delta_1, J\Psi_1}^C(e); \text{id}_{\Delta_1} \bullet d_2 \bullet \text{id}_{B \multimap C}$$

where $\text{MKC}_{A, \Delta_1, J\Psi_1}^C$ is a natural transformation with components

$$\mathcal{L}(A, \Delta_1 \bullet B \bullet J\Psi) \times \mathcal{L}(C, C) \rightarrow \mathcal{L}(A, \Delta_1 \bullet C \bullet C \multimap C)$$

so we make the definition

$$\begin{aligned} & \llbracket x : A \vdash_L \Delta_1, \text{mkc}(e, y) : B \multimap C, [y(e)/y]\Delta_2; \Psi_1 \cdot [y(e)/y]\Psi_2 \rrbracket =_{df} \\ & \text{MKC}_{A, \Delta_1, J\Psi_1}^C \llbracket x : A \vdash_L \Delta_1, e_1 : B \rrbracket; \text{id}_{\Delta_1} \bullet \llbracket y : C \vdash_L \Delta_2; \Psi_2 \rrbracket \bullet \text{id}_{B \multimap C}; \\ & \quad \text{Join}(\Delta_1, \Delta_2, B \multimap C, J\Psi_1, J\Psi_2) \end{aligned}$$

Notice that $\text{MKC}_{A, \Delta_1, J\Psi_1}^C$ corresponds to the one-premise form of the subtraction introduction rule

$$\frac{x : A \vdash_L \Delta_1, e : B; \Psi_1}{x : A \vdash_L \Delta_1, \text{mkc}(e, y) : B \multimap C, y(e) : C; \Psi_1} \text{TLL}_{\multimap-I}$$

which is equivalent in terms of provability to the more general form considered here [11].

The subtraction elimination rule has the form:

$$\frac{x : A \vdash_L \Delta_1, e_1 : B \multimap C; \Psi_1 \quad y : B \vdash_L e_2 : C, \Delta_2; \Psi_2 \quad |\Psi_1| = |\Psi_2|}{x : A \vdash_L \text{postp}(y \mapsto e_2, e_1), \Delta_1, [y(e_1)/y]\Delta_2; \Psi_1, [y(e_1)/y]\Psi_2} \text{TLL}_{\multimap-E}$$

This suggests a natural transformation with components

$$\Psi_{A, \Delta_1, \Delta_2, J\Psi_1, J\Psi_2} : \mathcal{L}(A, \Delta_1 \bullet (B \multimap C) \bullet J\Psi_1) \times \mathcal{L}(B, C \bullet \Delta_2 \bullet J\Psi_2) \rightarrow \mathcal{L}(A, \Delta_1 \bullet \Delta_2 \bullet J\Psi_1 \bullet J\Psi_2)$$

natural in $A, \Delta_1, \Delta_2, J\Psi_1, J\Psi_2$. Here $\text{postp}(y \mapsto e_2, e_1)$ is given type 1 and an application of left identity λ_{1, Δ_2} is assumed implicitly.

Given

$$e : A \rightarrow \Delta_1 \bullet (B \multimap C) \bullet J\Psi_1, \quad f : B \rightarrow C \bullet \Delta_2 \bullet J\Psi_2$$

and also $a : A' \rightarrow A, d_1 : \Delta_1 \rightarrow \Delta'_1, d_2 : \Delta_2 \rightarrow \Delta'_2, p_1 : J\Psi_1 \rightarrow J\Psi'_1, p_2 : J\Psi_2 \rightarrow J\Psi'_2$ naturality yields

$$\Psi_{A', \Delta'_1, \Delta'_2, J\Psi'_1, J\Psi'_2}((a; e; d_1 \bullet id_{B \bullet C} \bullet p_1), (f; id_C \bullet d_2 \bullet p_2)) =$$

$$a; \Psi_{A, \Delta_1, \Delta_2, J\Psi_1, J\Psi_2}(e, f); \lambda_{1, \Delta_1} \bullet d_1 \bullet d_2 \bullet p_1 \bullet p_2; \text{Join}(\Delta'_1, \Delta'_2, J\Psi_1, J\Psi_2)$$

In particular, taking $a : A \rightarrow \Delta_1 \bullet (B \bullet C)$, $e = id_{\Delta_1 \bullet (B \bullet C)}$, $d_1 = id_{\Delta_1}$, $d_2 : id_{\Delta_2}$, $p_1 = id_{J\Psi_1}$, $p_2 = id_{J\Psi_2}$ we obtain

$$\Psi_{A, \Delta_1, \Delta_2}(a, f) = a; \Psi_{A, \Delta_1, \Delta_2}(id_{\Delta_1 \bullet (C \bullet D)} \bullet id_{J\Psi_1}, f); \text{Join}(\Delta_1, \Delta_2, J\Psi_1, J\Psi_2)$$

Writing **POSTP**(f) for $\Psi_{A, \Delta_1, \Delta_2, J\Psi_1, J\Psi_2}(id_{\Delta_1 \bullet (B \bullet C)} \bullet id_{J\Psi_1}, f)$ we define

$$\llbracket x : A \vdash_L \Delta_1, \text{postp}(y \mapsto e_2, e_1), [y(e_1)/y]\Delta_2; \Psi_1, [y(e_1)/y], \Psi_2 \rrbracket =_{df}$$

$$\llbracket x : A \vdash_L \Delta_1, e_1 : B \bullet C \rrbracket; id_{\Delta_1} \bullet \text{POSTP} \llbracket y : B \vdash_L e_2 : C, \Delta_2; \Psi_2 \rrbracket; \text{Join}(\Delta_1, \Delta_2, J\Psi_1, J\Psi_2)$$

3.4.2.3. *Equations in context.* We have equations in context of the form

$\begin{array}{l} \Delta_1 = \Delta'_1 \\ \Delta_2 = \Delta'_2 \\ \Psi_1 = \Psi'_1 \\ \Psi_2 = \Psi'_2 \end{array}$	$\begin{array}{l} e_z \equiv z(\text{mkc}(e_1, y)) \\ e_p \equiv \text{postp}(z \mapsto e_2, \text{mkc}(e_1, y)) \\ x : B \vdash_L e_1 : A_1, \Delta_1; \Psi_1 = e'_1 : A_1, \Delta'_1; \Psi'_1 \\ y : A_2 \vdash_L \Delta_2; \Psi_2 = \Delta'_2; \Psi'_2 \\ z : A_1 \vdash_L e_2 : A_2, \Delta_3; \Psi_3 = e'_2 : A_2, \Delta'_3; \Psi'_3 \end{array}$
$\frac{x : B \vdash_L \Delta_1, e_p, [y(e_1)/y]\Delta_2, [e_z/z]\Delta_3; \Psi_1, [y(e_1)/y]\Psi_2, [e_z/z]\Psi_3 = \Delta'_1, [[e'_1/z]e'_2/y]\Delta_2, [e'_1/z]\Delta'_3; \Psi'_1, [[e'_1/z]e'_2/y]\Psi'_2, [e'_1/z]\Psi'_3}{\bullet - \beta}$	

We repeat the derivations of the redex and of the redutum.

Redex:

$$\frac{x : B \vdash_L e_1 : A_1, \Delta_1; \Psi_1 \quad y : A_2 \vdash_L \Delta_2; \Psi_2}{x : B \vdash_L \text{mkc}(e_1, y) : A_1 \bullet A_2, \Delta_1, [y(e_1)/y]\Delta_2; \Psi_1, [y(e_1)/y]\Psi_2 \quad z : A_1 \vdash_L e_2 : A_2, \Delta_3; \Psi_3}$$

$$x : B \vdash_L \Delta_1, \overbrace{\text{postp}(z \mapsto e_2, \text{mkc}(e_1, y))}^{e_p}, [y(e_1)/y]\Delta_2, \overbrace{[z(\text{mkc}(e_1, y)) / z]\Delta_3}^{e_z}; \Psi_1, [y(e_1)/y]\Psi_2, [z(\text{mkc}(e_1, y)/z)] : \Psi_3$$

Redutum:

$$\frac{x : B \vdash_L e'_1 : A_1, \Delta'_1; \Psi'_1 \quad z : A_1 \vdash_L e'_2 : A_2, \Delta'_3; \Psi'_3}{x : B \vdash_L \Delta'_1, [e'_1/z]\Delta'_3, [e'_1/z]e'_2 : A_2; \Psi'_1, [e'_1/z]\Psi'_3 \quad y : A_2 \vdash_L \Delta_2; \Psi_2}$$

$$x : B \vdash_L \Delta'_1, [[e'_1/z]e'_2/y]\Delta'_2, [e'_1/z]\Delta'_3; \Psi'_1, [[e'_1/z]e'_2/y]\Psi'_2, [e'_1/z]\Psi'_3$$

Given morphisms $n : B \rightarrow \Delta_1 \bullet A_1$ and $m : A_1 \rightarrow \Delta_3 \bullet A_2$, for these equations to be satisfied we need the following diagram to commute (omitting non-linear terms):

$$\begin{array}{ccc} B & \xrightarrow{n} & \Delta_1 \bullet A_1 \\ \text{MKC}^{A_2}(n) \downarrow & & \downarrow id_{\Delta_1} \bullet m \\ \Delta_1 \bullet (A_1 \bullet A_2) \bullet A_2 & \xrightarrow{\text{POSTP}(m) \bullet id_{A_2}} & \Delta_1 \bullet \Delta_3 \bullet A_2 \end{array}$$

in particular, taking $n = id_{A_1}$ we have

$$\begin{array}{ccc} A_1 & \xrightarrow{m} & \Delta_3 \bullet A_2 \\ \text{MKC}_2^A(id_{A_1}) \downarrow & \nearrow \text{POSTP}(m) \bullet id_{A_2} & \\ (A_1 \bullet A_2) \bullet A_2 & & \end{array}$$

Assuming the above decomposition to be unique, we can show that the η equation in context is also satisfied:

$$\frac{|\Delta| = |\Delta'| \quad |\Psi| = |\Psi'| \quad z : B \vdash_L \Delta, \Psi = \Delta', \Psi'}{z : B \vdash_L \text{postp}(x \mapsto y, e), \text{mkc}(x(e), y) : A_1 \bullet A_2, \Delta, \Psi = e : A_1 \bullet A_2, \Delta'; \Psi'} \quad (3.2)$$

and conclude that there is a natural isomorphism between the maps

$$\frac{A \rightarrow \Delta \bullet B}{A \bullet B \rightarrow \Delta}$$

i.e., that \bullet is the left adjoint to the bifunctor \bullet .

3.4.3. Functors. Recall that a model of Linear-Non Linear co-intuitionistic logic consists of a symmetric comonoidal adjunction $\mathcal{L} : H \dashv J : C$ where $\mathcal{L} = (\mathcal{L}, \perp, \oplus, \multimap)$ is a symmetric monoidal coclosed category and $C = (C, 0, +, -)$ is a cocartesian coclosed category.

We use the same symbols for the functors $J : C \rightarrow \mathcal{L}$ and $\mathcal{H} : \mathcal{L} \rightarrow C$ in the models and for the operators that represent them in the language.

3.4.3.1 rules for $J : C \rightarrow \mathcal{L}$.

$$\boxed{\begin{array}{c} TL_J_I \\ \frac{x : A \vdash_{\mathcal{L}} \Delta; t : T, \Psi}{x : A \vdash_{\mathcal{L}} \Delta, Jt : JT; \Psi} \end{array}} \quad (3.3)$$

If $\Delta = \bar{R} : \Delta$ and $\Psi = \bar{S} : \Psi$, then the categorical interpretation of the rule is an application of α^{-1} :

$$\begin{array}{c} A \xrightarrow{\bar{R} \bullet Jt \bullet \bar{S}} \Delta \bullet JT \bullet J\Psi \\ \hline A \xrightarrow{(\bar{R} \bullet Jt) \bullet \bar{S}} (\Delta \bullet JT) \bullet J\Psi \end{array}$$

$$\boxed{\begin{array}{c} TL_J_E \text{ elimination} \\ \frac{x : A \vdash_{\mathcal{L}} \Delta, e : JT; \Psi_1 \quad y : T \vdash_C \Psi_2 \text{ where } |\Psi_1| = |\Psi_2|}{x : A \vdash_{\mathcal{L}} \Delta; \Psi_1 \cdot \text{let } Jy = e \text{ in } \Psi_2} \end{array}} \quad (3.4)$$

If $\Delta = \bar{R} : \Delta$, $\Psi_1 = \bar{R}' : \Psi_1$, $\Psi_2 = \bar{S} : \Psi_2$, then the categorical interpretation of the rule is given by an operation of the form

$$\mathcal{L}(A, \Delta \bullet JT \bullet J\Psi_1) \times C(T, \Psi_2) \rightarrow \mathcal{L}(A, \Delta \bullet J\Psi_1 \bullet J\Psi_2)$$

given by the following compositions

$$\begin{array}{c} \frac{\frac{A \xrightarrow{\bar{R} \bullet e \bullet J\bar{R}'} \Delta \bullet JT \bullet J\Psi_1 \quad JT \xrightarrow{\bar{S}} J\Psi_2 \text{ in } \mathcal{L}}{A \xrightarrow{\bar{R} \bullet e \bullet J(\bar{R}')} \Delta \bullet J(T) \bullet J(\Psi_1)} \quad \frac{T \xrightarrow{\bar{S}} \Psi_2 \text{ in } C}{JT \xrightarrow{\bar{S}} J\Psi_2 \text{ in } \mathcal{L}}}{A \xrightarrow{\bar{R} \bullet e \bullet J(\bar{R}')} \Delta \bullet J(T) \bullet J(\Psi_1) \xrightarrow{id_{\Delta} \bullet J(\bar{S}) \bullet id_{J(\Psi_1)}} \Delta \bullet J(\Psi_1) \bullet J(\Psi_2) \xrightarrow{id_{\Delta} \bullet j_{\Psi_1 \Psi_2}^{-1}} \Delta \bullet J(\Psi_1 + \Psi_2)} \\ \text{since } |\Psi_1| = |\Psi_2|, \quad \xrightarrow{id_{\Delta} \bullet \nabla_{\Psi_1}} \Delta \bullet J(\Psi_1) \end{array}$$

3.4.3.2 rules for $H : \mathcal{L} \rightarrow C$.

$$\boxed{\begin{array}{c} TC_H_I \\ \frac{x : A \vdash_{\mathcal{L}} \bar{R} : \Delta, e : B; \Psi}{x : A \vdash_{\mathcal{L}} \bar{R} : \Delta; He : HB, \Psi} \end{array}} \quad (3.5)$$

Let $\Delta = \bar{R} : \Delta$ and $\Psi = \bar{S} : \Psi$: then

$$\begin{array}{c}
\frac{A \xrightarrow{\bar{R} \oplus e \oplus J(\bar{S})} \Delta \bullet B \bullet J(\Psi)}{A \xrightarrow{\bar{R} \oplus JH(e) \bullet J(\bar{S})} \Delta \bullet JH(B) \bullet J(\Psi)} \text{ using } \eta_B : B \rightarrow JHB \\
\\
\boxed{
\begin{array}{c}
TC_H_E \\
\frac{x : B \vdash_L \Delta; t : HA, \Psi_1 \quad y : A \vdash_L \Psi_2}{x : B \vdash_L \Delta; \Psi_1, \text{let } Hy = t \text{ in } \Psi_2}
\end{array}
} \tag{3.6}
\end{array}$$

The categorical interpretation of $H \text{ elim}_1$ is as follows: Let $\Psi_1 = \bar{R} : \Psi_1$ and $\Psi_2 = \bar{S} : \Psi_2$. Then we have the following compositions:

$$\begin{array}{c}
\frac{S \xrightarrow{t+\bar{R}} H(A) + \Psi \quad \frac{A \xrightarrow{J(\bar{S})} J(\Psi) \quad \text{in } \mathcal{L}}{HA \xrightarrow{HJ(\bar{S})} HJ(\Psi) \xrightarrow{\epsilon_\Psi} \Psi \quad \text{in } C}}{S \xrightarrow{t+\bar{R}} H(A) + \Psi \xrightarrow{HJ(\bar{S})+id_\Psi} \Psi + \Psi \xrightarrow{\nabla_\Psi} \Psi}
\end{array}$$

4. RELATED AND FUTURE WORK

The most comprehensive treatment of ILL is in Gavin Bierman's thesis [5]. There one finds the Proof Theory (Chapter 2), i.e., the sequent calculus with cut-elimination, natural deduction and axiomatic versions of ILL. Then (Chapter 3) a term assignment to the natural deduction and to the sequent calculus versions are presented with β -reductions and commutative conversions, and strong normalization and confluence are proved for the resulting calculus. A painstaking analysis of the rules of the labeled calculus leads to the construction of a categorical model of ILL, a *linear category*, in particular of the exponential part, a main contribution of Bierman and of the Cambridge school of the 1990s with respect to previous models by Seely and Lafont. Bellin [3] presents a categorical model of co-intuitionistic linear logic based on a dualization of Bierman [5] construction for ILL..

Benton's work [4] on LNL logic, which we read in a TeX report, presents the categorical model for Linear-Non-Linear Intuitionistic logic LNL. Chapter 2 shows how to obtain a LNL model from a Linear Category and viceversa. Versions of the sequent calculus for LNL are considered and cut-elimination is proved for one such version. Then Natural Deduction is given with term assignment and the categorical interpretation of a fragment of the natural deduction system. β reductions and commuting conversions are presented. The present work follows Benton's paper aiming at a (non-trivial) dualization of it.

Bi-intuitionistic logic was introduced by C.Rauszer [19] (Semi-boolean algebras and their applications to intuitionistic logic with dual operations. In *Fundamenta Mathematicae*, 83, pp. 219 249, 1974) with an algebraic and Kripke semantics [20] (An algebraic and Kripke-style approach to a certain extension of intuitionistic logic, *Dissertationes Mathematicae* 168, 1980) and a Gentzen style sequent calculus [?] (A formalization of the propositional calculus of H-B logic. *Studia Logica*, 33(1):23 34, 1974). Co-intuitionistic logic requires a multiple conclusion system, because of the cotensor in the linear case and of contraction right in the non-linear one. This raises the problem of the relations between intuitionistic implication and disjunction, and, dually, between subtraction and conjunction. In the case of the logic FILL that extends ILL with the cotensor (*par*) applying Maheara and Dragalin's restriction that only one formula occurs in the succedent of the premise of

an implication right, yields a calculus that does not satisfies cut-elimination, as noticed by Schellinx [21]. Similarly, in the logic BILL (*Bi-Intuitionistic Linear Logic*) requiring that only one formula occurs in the antecedent of the premise of a subtraction left yields a system that does not satisfy cut-elimination.

$$\frac{\Gamma, A \vdash B}{\Gamma \vdash A \multimap B} \multimap R \qquad \frac{A \vdash B, \Delta}{A \multimap B \vdash \Delta} \multimap E$$

As a simple counterexample, consider the sequent $p \Rightarrow q, r \rightarrow ((p - q) \wedge r)$ given by Pinto and Uusatu around 2003 [17], which is provable with cut but not cut-free with Dragalin's restrictions.

Hyland and De Paiva introduced a sequent calculus for FILL labeled with terms

$$\frac{\bar{y} : \Gamma, x : A \vdash t : B, \bar{u} : \Delta}{\bar{y} : \Gamma \vdash \lambda x : TA \multimap B, \bar{u} : \Delta} \multimap R$$

where $x : A$ occurs in $t : B$ if and only if there is an “essential dependency” of B from A . The restriction on the $\multimap I$ is that x does not occur in the terms $\bar{u} : \Delta$. The original term assignment did not guarantee cut-elimination, as noticed by Bierman [6]; the assignment to *par left* ($\oplus L$) had to be fine tuned, as indicated by Bellin [1]

A detailed presentation of the term calculus for FILL with a full proof of cut elimination, is in [14], where the correctness for a categorical semantics for FILL is also proved. Another correct formalization of FILL, a sequent calculus with a relational annotation, was given by Braüner and De Paiva [7], with a proof of cut-elimination. The second author [1] gave a system of proof nets for FILL which sequentialize in the sequent calculus with term assignment; the essential fact here is that $x : A$ occurs in $t : B$ if and only if there is a “directed chain” between A and B in the proof structure. Here cut elimination is proved by reduction to cut-elimination for proof nets.

A system of two-sided proof nets (in the style of natural deduction) was given by Cockett and Seely [9]. For Bi-Intuitionistic Linear Logic, they gave also a system of proof nets, corresponding to a sequent calculus without annotations and restrictions that therefore collapses into classical MLL. Recently, Clouston, Dawson, Gore and Tiu [?] gave an annotation-free formalization for BILL, alternative to sequent calculi, in the form of deep-inference and display calculi for BILL. This calculus enjoys cut-elimination and is relevant to the categorical semantics bi-intuitionistic linear logic. Annotation-free formalizations of Bi-Intuitionistic Logic use the display calculus [12], nested sequents [13] and deep inference [18].

Tristan Crolard [10, 11] made an in-depth study of Rauszer's logic. In [10] he showed that models of Rauszer logic (called “subtractive logic”) based on bi-cartesian closed categories (with co-exponents) collapse to preorders. He also studied models of subtractive logic and showed that its first order theory is constant-domain logic, thus it is not a conservative extension of intuitionistic logic.

Crolard [11] develops the type theory for subtractive logic, extending a system of multiple conclusion classical natural deduction with a connective of subtraction and then decorating proofs with a system of annotations of dependencies that allows us to identify “constructive proofs”: these are derivations where only the premise of an implication introduction depends on the discharged assumption and only the premise of a subtraction elimination depends on the discharged conclusion. Therefore Crolard's sequent calculus with annotations is not affected by the counterexamples to cut-eliminations.

The type theory is Parigot $\lambda\mu$ -calculus extended with operators for sums, products and subtraction, where the operators for subtraction introduction and elimination are understood as a calculus of co-routines. A constructive system of co-routines is then obtained by imposing restrictions on terms corresponding to the restrictions on constructive proofs.

In a series of papers the second author gave a “pragmatic” interpretation of bi-intuitionism, where intuitionistic and co-intuitionistic logic are interpreted as logics of the acts of assertion and making a hypothesis, respectively, the interactions between the two sides depending on negations (see, e.g., [2] Assertions, hypotheses, conjectures, expectations: Rough-sets semantics and proof-theory.) Here the separation between intuitionistic and co-intuitionistic logic and their models is given a linguistic motivation. Writing $\vdash p$ for the type of assertions that p is true and using intuitionistic connectives with the BHK interpretation, one gives a “pragmatic interpretation” of *ILL*, where an expression A is *justified* or *unjustified* (Dalla Pozza and Garola, A pragmatic interpretation of intuitionistic propositional logic, *Erkenntnis*, 43, 1995, pp.81-109.). Similarly, writing $\mathcal{H} p$ for the type of hypotheses that p is true, and using co-intuitionistic connectives, one builds a co-intuitionistic language, for which an analogue “pragmatic interpretation” has been attempted. Both languages may be given a modal interpretation in *S4*, with $(\vdash p)^M = \Box p$ and $(\mathcal{H} p)^M = \Diamond p$. Notice that here there is a semantic duality between an assertion $\vdash p$ and a hypothesis $\mathcal{H} \neg p$, as $\Box p$ and $\Diamond \neg p$ are contradictory. Similarly there is a semantic duality between $\mathcal{H} p$ and $\vdash \neg p$, but not between $\vdash p$ and the hypothesis $\mathcal{H} p$. A useful direction of research in the proof theory of bi-intuitionism may be the investigation the relations between co-intuitionistic proofs and intuitionistic refutations.

It is in this context that a term assignment for co-intuitionistic logic has been developed, starting from Crolard’s definition but independently of the $\lambda\mu$ framework. This calculus was used here as a term assignment of Dual LNL logic.

Finally, to achieve the project outlined in the introduction of putting together intuitionistic and co-intuitionistic adjoint logic in the environment of BILL the definition of a suitable syntax for BILL will play a key role.

5. CONCLUSION

TODO

REFERENCES

- [1] Gianluigi Bellin. Subnets of proof-nets in multiplicative linear logic with MIX. *Mathematical Structures in Computer Science*, 7(6):663–699, 1997. URL: <http://journals.cambridge.org/action/displayAbstract?aid=44699>.
- [2] Gianluigi Bellin. *Assertions, Hypotheses, Conjectures, Expectations: Rough-Sets Semantics and Proof Theory*, pages 193–241. Springer Netherlands, Dordrecht, 2014. URL: http://dx.doi.org/10.1007/978-94-007-7548-0_10, doi:10.1007/978-94-007-7548-0_10.
- [3] Gianluigi Bellin. Categorical proof theory of co-intuitionistic linear logic. *Logical Methods in Computer Science*, 10(3):Paper 16, September 2014.
- [4] Nick Benton. A mixed linear and non-linear logic: Proofs, terms and models (preliminary report). Technical Report UCAM-CL-TR-352, University of Cambridge Computer Laboratory, 1994.
- [5] G. M. Bierman. *On Intuitionistic Linear Logic*. PhD thesis, Wolfson College, Cambridge, December 1993.
- [6] Gavin M. Bierman. A note on full intuitionistic linear logic. *Ann. Pure Appl. Logic*, 79(3):281–287, 1996. URL: [https://doi.org/10.1016/0168-0072\(96\)00004-8](https://doi.org/10.1016/0168-0072(96)00004-8), doi:10.1016/0168-0072(96)00004-8.
- [7] Torben Braüner and Valeria de Paiva. A formulation of linear logic based on dependency-relations. In *Computer Science Logic, 11th International Workshop, CSL ’97, Annual Conference of the EACSL, Aarhus, Denmark, August 23-29, 1997, Selected Papers*, pages 129–148, 1997. URL: <https://doi.org/10.1007/BFb0028011>, doi:10.1007/BFb0028011.
- [8] J.R.B. Cockett and R.A.G. Seely. Proof theory for full intuitionistic linear logic, bilinear logic, and mix categories. *Theory and Applications of Categories*, 3(5):85–131, 1997.
- [9] J.R.B. Cockett and R.A.G. Seely. Weakly distributive categories. *Journal of Pure and Applied Algebra*, 114(2):133 – 173, 1997.
- [10] Tristan Crolard. Subtractive logic. *Theoretical Computer Science*, 254(1-2):151–185, 2001.

- [11] Tristan Crolard. A formulae-as-types interpretation of subtractive logic. *J. Log. Comput.*, 14(4):529–570, 2004. URL: <https://doi.org/10.1093/logcom/14.4.529>.
- [12] Rajeev Goré. Dual intuitionistic logic revisited. In *Automated Reasoning with Analytic Tableaux and Related Methods, International Conference, TABLEUX 2000, St Andrews, Scotland, UK, July 3-7, 2000, Proceedings*, pages 252–267, 2000. URL: https://doi.org/10.1007/10722086_21, doi:10.1007/10722086_21.
- [13] Rajeev Goré, Linda Postniece, and Alwen Tiu. Cut-elimination and proof-search for bi-intuitionistic logic using nested sequents. In *Advances in Modal Logic 7, papers from the seventh conference on "Advances in Modal Logic," held in Nancy, France, 9-12 September 2008*, pages 43–66, 2008. URL: <http://www.aiml.net/volumes/volume7/Gore-Postniece-Tiu.pdf>.
- [14] Harley Eades III and Valeria de Paiva. Multiple conclusion linear logic: Cut elimination and more. In *Logical Foundations of Computer Science - International Symposium, LFCS 2016, Deerfield Beach, FL, USA, January 4-7, 2016. Proceedings*, pages 90–105, 2016. URL: https://doi.org/10.1007/978-3-319-27683-0_7, doi:10.1007/978-3-319-27683-0_7.
- [15] Maria Emilia Maietti, Paola Maneggia, Valeria de Paiva, and Eike Ritter. Relating categorical semantics for intuitionistic linear logic. *Applied Categorical Structures*, 13(1):1–36, 2005. URL: <http://dx.doi.org/10.1007/s10485-004-3134-z>, doi:10.1007/s10485-004-3134-z.
- [16] Michel Parigot. Lambda-mu-calculus: An algorithmic interpretation of classical natural deduction. In Andrei Voronkov, editor, *Logic Programming and Automated Reasoning*, volume 624 of *Lecture Notes in Computer Science*, pages 190–201. Springer Berlin / Heidelberg, 1992.
- [17] Luis Pinto and Tarmo Uustalu. Relating sequent calculi for bi-intuitionistic propositional logic. In *Proceedings Third International Workshop on Classical Logic and Computation, CL&C 2010, Brno, Czech Republic, 21-22 August 2010.*, pages 57–72, 2010. URL: <https://doi.org/10.4204/EPTCS.47.7>, doi:10.4204/EPTCS.47.7.
- [18] Linda Postniece. Deep inference in bi-intuitionistic logic. In *Logic, Language, Information and Computation, 16th International Workshop, WoLLIC 2009, Tokyo, Japan, June 21-24, 2009. Proceedings*, pages 320–334, 2009. URL: https://doi.org/10.1007/978-3-642-02261-6_26, doi:10.1007/978-3-642-02261-6_26.
- [19] Cecylia Rauszer. Semi-boolean algebras and their applications to intuitionistic logic with dual operations. *Fundamenta Mathematicae*, 83:219–249, 1974.
- [20] Cecylia Rauszer. An algebraic and kripke-style approach to a certain extension of intuitionistic logic. In *Dissertationes Mathematicae*, 167:1–67, Institut Mathématique de l’Académie Polonaise des Sciences 1980.
- [21] Harold Schellinx. Some syntactical observations on linear logic. *Journal of Logic and Computation*, 1(4):537–559, 1991.

APPENDIX A. COMMUTING CONVERSIONS

Non linear rules.

- (1) *disjunction intro* TC_{+I_1} and TC_{+I_2} commute upwards with every inference and the terms obtained are the same.
- (2) *disjunction elim* TC_{+E} commutes upwrds with inferences in the derivation of the major premise, the terms assigned to the resulting subderivations are equated. For instance

$$\frac{\begin{array}{c} y : T_2 \vdash_C \Psi_2, t_1 : T_4 + T_5 \\ z : T_3 \vdash_C \Psi_3, t_2 : T_4 + T_5 \end{array} \quad \begin{array}{c} x : S \vdash_C \Psi_1, t : T_2 + T_3 \end{array} \quad |\Psi_2| = |\Psi_3|}{\begin{array}{c} x : S \vdash_C \Psi_1, \text{case } t \text{ of } y.\Psi_2, z.\Psi_3, \text{case } t \text{ of } y.t_1, z.t_2 : T_4 + T_5 \end{array} \quad \begin{array}{c} v_1 : T_4 \vdash_C \Psi_4 \quad v_2 : T_5 \vdash_C \Psi_5 \end{array} \quad |\Psi_4| = |\Psi_5|}{x : S \vdash_C \Psi_1, \text{case } t \text{ of } y.\Psi_2, z.\Psi_3, \text{case } (\text{case } t \text{ of } y.t_1, z.t_2) \text{ of } v_1.\Psi_4, v_2.\Psi_5}$$

commutes to

$$\frac{\begin{array}{c} |\Psi_4| = |\Psi_5| \\ v_1 : T_4 \vdash_C \Psi_4 \\ v_2 : T_5 \vdash_C \Psi_5 \end{array} \quad \begin{array}{c} y : T_2 \vdash_C \Psi_2, t_1 : T_4 + T_5 \end{array} \quad \begin{array}{c} |\Psi_4| = |\Psi_5| \\ v_1 : T_4 \vdash_C \Psi_4 \\ v_2 : T_5 \vdash_C \Psi_5 \end{array} \quad \begin{array}{c} z : T_3 \vdash_C \Psi_3, t_2 : T_4 + T_5 \end{array}}{\begin{array}{c} x : S \vdash_C \Psi_1, t : T_2 + T_3 \end{array} \quad \begin{array}{c} y : T_2 \vdash_C \Psi_2, \text{case } t_1 \text{ of } v_1.\Psi_4, v_2.\Psi_5 \end{array} \quad \begin{array}{c} z : T_3 \vdash_C \Psi_3, \text{case } t_2 \text{ of } v_1.\Psi_4, v_2.\Psi_5 \end{array}}{x : S \vdash_C \Psi_1, \text{case } t \text{ of } y.(\Psi_2, \text{case } t_1 \text{ of } v_1.\Psi_4, v_2.\Psi_5), z.(\Psi_3, \text{case } t_2 \text{ of } v_1.\Psi_4, v_2.\Psi_5)}$$

Remark 48. If $t_1 = \text{inl } s_1$ and $u_2 = \text{inr } s_2$ then after commutation

$$y : T_2 \vdash_C \Psi_2, \text{case } \text{inl } s_1 \text{ of } v_1.\Psi_4, v_2.\Psi_5 \rightsquigarrow_\beta y : T_2 \vdash_C \Psi_2, [s_1/v_1]\Psi_4$$

$$y : T_3 \vdash_C \Psi_3, \text{case inr } s_2 \text{ of } v_1.\Psi_4, v_2.\Psi_5 \rightsquigarrow_\beta y : T_2 \vdash_C \Psi_3, [s_1/v_2]\Psi_5$$

- (3) Subtraction introduction TC_{-I} commutes upwards with inferences in both branches with any inference \mathcal{I} :

$$\frac{\frac{x : S \vdash_C t'_1 : T_1, \Psi'_1}{x : S \vdash_C t_1 : T_1, \Psi_1} \mathcal{I} \quad y : T_2 \vdash_C \Psi_2}{x : S \vdash_C \Psi_1, \text{mkc}(t, y) : T_1 - T_2, [y(t)/y]\Psi_2} \text{TC}_{-I} \quad \frac{\frac{x : S \vdash_C t_1 : T_1, \Psi_1 \quad \frac{y : T_2 \vdash_C \Psi'_2}{y : T_2 \vdash_C \Psi_2} \mathcal{I}}{x : S \vdash_C \Psi_1, \text{mkc}(t_1, y) : T_1 - T_2, [y(t)/y]\Psi_2} \mathcal{I} \quad y : T_2 \vdash_C \Psi_2}{x : S \vdash_C \Psi_1, \text{mkc}(t_1, y) : T_1 - T_2, [y(t)/y]\Psi_2} \text{TC}_{-I}$$

commutes to

$$\frac{\frac{x : S \vdash_C t_1 : T_1, \Psi'_1 \quad y : T_2 \vdash_C \Psi_2}{x : S \vdash_C \Psi'_1, \text{mkc}(t_1, y) : T_1 - T_2, [y(t')/y]\Psi_2} \text{TC}_{-I} \quad y : T_2 \vdash_C \Psi_2}{x : S \vdash_C \Psi_1, \text{mkc}(t_1, y) : T_1 - T_2, [y(t)/y]\Psi_2} \mathcal{I} \quad \frac{\frac{x : S \vdash_C t_1 : T_1, \Psi_1 \quad y : T_2 \vdash_C \Psi'_2}{x : S \vdash_C \Psi_1, \text{mkc}(t_1, y) : T_1 - T_2, \Psi'_2} \text{TC}_{-I} \quad y : T_2 \vdash_C \Psi_2}{x : S \vdash_C \Psi_1, \text{mkc}(t_1, y) : T_1 - T_2, \Psi_2} \mathcal{I}$$

- (4) Subtraction elimination TC_{-E} commutes upwards. For instance,

$$\frac{\frac{x : S \vdash_C w : S_1, \Psi_1 \quad z : S_2 \vdash_C \Psi_2, t_1 : T_1 - T_2}{x : S \vdash_C \Psi_1, \text{mkc}(w, z) : S_1 - S_2, [z(w)/z]\Psi_2, [z(w)/z]t_1 : T_1 - T_2} \quad y : T_1 \vdash_C t : T_2, \Psi_3}{x : S \vdash_C \Psi_1, \text{mkc}(w, z) : S_1 - S_2, [z(w)/z]\Psi_2, \text{postp}(y \mapsto t, [z(w)/z]t_1), [y([z(w)/z]t_1)/y]\Psi_3} \text{TC}_{-E}$$

commutes to

$$\frac{x : S \vdash_C w : S_1, \Psi_1 \quad \frac{z : S_2 \vdash_C \Psi_2, t_1 : T_1 - T_2 \quad y : T_1 \vdash_C t : T_2, \Psi_3}{z : S_2 \vdash_C \Psi_2, \text{postp}(y \mapsto t, t_1), [y(t_1)/y]\Psi_3}}{x : S \vdash_C \Psi_1, \text{mkc}(w, z) : S_1 - S_2, [z(w)/z]\Psi_2, \text{postp}(y \mapsto t, [z(w)/z]t_1), [z(w)/z][y(t_1)/y]\Psi_3}$$

where

$$[z(w)/z][y(t_1)/y]\Psi_3 = [y([z(w)/z]t_1)/y]\Psi_3 \quad (\text{A.1})$$

since $z \notin \Psi_2$.

Linear rules.

- (5) The \perp introduction rule $\text{TILL}_{\perp I}$ rule commutes with any inference, as $\text{connect}_{\perp \text{toe}}$ can be “rewired” to any term in the context.
- (6) The commutations of the rules for linear subtraction $\text{TILL}_{\bullet - I}$ and $\text{TILL}_{\bullet - E}$ are similar to those for non-linear subtraction.
- (7) Linear disjunction (*par*) introduction ($\text{TLL}_{\oplus I}$) commutes with any inference. Linear disjunction elimination ($\text{TLL}_{\oplus E}$) also commutes upwards. For example (writing a proof without non-linear parts for simplicity) we have the following:

$$\frac{\frac{x : A \vdash_L \Delta, e : (B_1 \oplus B_2) \quad \frac{y : B_1 \vdash_L \Delta_1, e_1 : C_1 \oplus C_2 \quad v : C_1 \vdash_L \Delta_3 \quad w : C_2 \vdash_L \Delta_4}{y : B_1 \vdash_L \Delta_1, [\text{casel } e_1/v]\Delta_3, [\text{caser } e_1/w]\Delta_4} \oplus E}{x : A \vdash_L \Delta, [\text{casel } e/y]\Delta_1, [\text{casel } e/y][\text{casel } e_1/v]\Delta_3, [\text{casel } e/y][\text{caser } e_1/w]\Delta_4, [\text{caser } e/z]\Delta_2} \oplus E$$

commutes to

$$\frac{\frac{x : A \vdash_L \Delta, e : (B_1 \oplus B_2) \quad y : B_1 \vdash_L \Delta_1, e_1 : C_1 \oplus C_2 \quad z : B_2 \vdash_L \Delta_2}{x : A \vdash_L \Delta, [\text{casel } e/y]\Delta_1, [\text{casel } e/y]e_1 : C_1 \oplus C_2, [\text{caser } e/z]\Delta_2} \oplus E \quad \frac{v : C_1 \vdash_L \Delta_3 \quad w : C_2 \vdash_L \Delta_4}{x : A \vdash_L \Delta, [\text{casel } e/y]\Delta_1, [\text{casel } [\text{casel } e/y]e_1/v]\Delta_3, [\text{caser } [\text{casel } e/y]e_1/w]\Delta_4, [\text{caser } e/z]\Delta_2} \oplus E}{x : A \vdash_L \Delta, [\text{casel } e/y][\text{casel } e_1/v]\Delta_3, [\text{caser } [\text{casel } e/y]e_1/w]\Delta_4, [\text{caser } e/z]\Delta_2} \oplus E$$

Now

$$[\text{casel } e/y][\text{casel } e_1/v]\Delta_3 = [\text{casel } [\text{casel } e/y]e_1/v]\Delta_3 \quad (\text{A.2})$$

because y does not occur in Δ_3 , only in e_1 and

$$[\text{casel } e/y][\text{casel } e_1/w]\Delta_4 = [\text{caser } [\text{casel } e/y]e_1/w]\Delta_4$$

because y does not occur in Δ_4 , only in e_1 .

APPENDIX B. PROOFS

B.1. Proof of Lemma 23. We show that both of the maps:

$$j_{R,S}^{-1} := JR \oplus JS \xrightarrow{\eta} JH(JR \oplus JS) \xrightarrow{Jh_{A,B}} J(HJR + HJS) \xrightarrow{J(\varepsilon_R + \varepsilon_S)} J(R + S)$$

$$j_0^{-1} := \perp \xrightarrow{\eta} JH \perp \xrightarrow{Jh_{\perp}} J0$$

are mutual inverses with $j_{R,S} : J(R + S) \rightarrow JR \oplus JS$ and $j_0 : \perp \rightarrow J0$ respectively.

Case. The following diagram implies that $j_{R,S}^{-1} ; j_{R,S} = \text{id}$:

$$\begin{array}{ccccc}
 JR \oplus JS & \xrightarrow{\eta} & JH(JR \oplus JS) & & \\
 \downarrow \eta \oplus \eta & & \downarrow Jh & & \\
 JR \oplus JS & \xleftarrow{J\varepsilon \oplus J\varepsilon} & JHJR \oplus JHJS & \xleftarrow{j} & J(HJR + HJS) \\
 & \searrow j & & \downarrow J(\varepsilon + \varepsilon) & \\
 & & & & J(R + S)
 \end{array}$$

The two top diagrams both commute because η and ε are the unit and counit of the adjunction respectively, and the bottom diagram commutes by naturality of j .

Case. The following diagram implies that $j_{R,S} ; j_{R,S}^{-1} = \text{id}$:

$$\begin{array}{ccccc}
 J(R + S) & \xrightarrow{j} & JR \oplus JS & & \\
 \downarrow \eta & & \downarrow \eta & & \\
 J(R + S) & \xleftarrow{J\varepsilon} & JHJ(R + S) & \xrightarrow{JHj} & JH(JR \oplus JS) \\
 & \searrow J(\varepsilon + \varepsilon) & & \downarrow Jh & \\
 & & & & J(HJR + HJS)
 \end{array}$$

The top left and bottom diagrams both commute because η and ε are the unit and counit of the adjunction respectively, and the top right diagram commutes by naturality of η .

Case. The following diagram implies that $j_0^{-1} ; j_0 = \text{id}$:

$$\begin{array}{ccc}
 \perp & \xrightarrow{\eta} & \mathbf{JH} \perp \\
 \parallel & & \downarrow \mathbf{Jh}_{\perp} \\
 \perp & \xleftarrow{j_0} & \mathbf{J0}
 \end{array}$$

This diagram holds because η is the unit of the adjunction.

Case. The following diagram implies that $j_0; j_0^{-1} = \text{id}$:

$$\begin{array}{ccccc}
 \mathbf{J0} & & \xrightarrow{j_0} & & \perp \\
 & \searrow \eta & & & \downarrow \eta \\
 & & \mathbf{JHJ0} & & \\
 & \swarrow \mathbf{Jh}_{\perp} & & \swarrow \mathbf{JHj_0} & \\
 \mathbf{J0} & & \xleftarrow{\mathbf{Jh}_{\perp}} & & \mathbf{JH} \perp
 \end{array}$$

The top-left and bottom diagrams commute because η and ε are the unit and counit of the adjunction respectively, and the top-right diagram commutes by naturality of η .

B.2. Proof of Lemma 25. Since $?$ is the composition of two symmetric comonoidal functors we know it is also symmetric comonoidal, and hence, the following diagrams all hold:

$$\begin{array}{ccc}
 ?((A \oplus B) \oplus C) & \xrightarrow{r_{A \oplus B, C}} & ?(A \oplus B) \oplus ?C \\
 \downarrow ?\alpha_{A, B, C} & & \downarrow r_{A, B} \oplus \text{id}_{?C} \\
 ?(A \oplus (B \oplus C)) & & (?A \oplus ?B) \oplus ?C \\
 \downarrow r_{A, B \oplus C} & & \downarrow \alpha_{?A, ?B, ?C} \\
 ?A \oplus ?(B \oplus C) & \xrightarrow{\text{id}_{?A} \oplus r_{B, C}} & ?A \oplus (?B \oplus ?C)
 \end{array}$$

$$\begin{array}{ccc}
 ?(\perp \oplus A) & \xrightarrow{r_{\perp, A}} & ?\perp \oplus ?A \\
 \downarrow ?\lambda_A & & \downarrow r_{\perp} \oplus \text{id}_{?A} \\
 ?A & \xrightarrow{\lambda^{-1}_{?A}} & \perp \oplus ?A
 \end{array}$$

$$\begin{array}{ccc}
 ?(A \oplus \perp) & \xrightarrow{r_{A, \perp}} & ?A \oplus ?\perp \\
 \downarrow ?\rho_A & & \downarrow \text{id}_{?A} \oplus r_{\perp} \\
 ?A & \xrightarrow{\rho^{-1}_{?A}} & ?A \oplus \perp
 \end{array}$$

$$\begin{array}{ccc}
 ?(A \oplus B) & \xrightarrow{r_{A, B}} & ?A \oplus ?B \\
 \downarrow ?\beta_{A, B} & & \downarrow \beta_{?A, ?B} \\
 ?(B \oplus A) & \xrightarrow{r_{B, A}} & ?B \oplus ?A
 \end{array}$$

Next we show that $(?, \eta, \mu)$ defines a monad where $\eta_A : A \longrightarrow ?A$ is the unit of the adjunction, and $\mu_A = J\varepsilon_{HA} : ??A \longrightarrow ?A$. It suffices to show that every diagram of Definition 13 holds.

Case.

$$\begin{array}{ccc}
 ?^3 A & \xrightarrow{\mu_{?A}} & ?^2 A \\
 \downarrow ?\mu_A & & \downarrow \mu_A \\
 ?^2 A & \xrightarrow{\mu_A} & ?A
 \end{array}$$

It suffices to show that the following diagram commutes:

$$\begin{array}{ccc}
 J(H(?^2 A)) & \xrightarrow{J\varepsilon_{H ?A}} & J(H ?A) \\
 \downarrow J(H\mu_A) & & \downarrow J\varepsilon_{HA} \\
 J(H ?A) & \xrightarrow{J\varepsilon_{HA}} & J(HA)
 \end{array}$$

But this diagram is equivalent to the following:

$$\begin{array}{ccc}
 HJHJHA & \xrightarrow{\varepsilon_{HJHA}} & HJHA \\
 \downarrow HJ\varepsilon_{HA} & & \downarrow \varepsilon_{HA} \\
 HJHA & \xrightarrow{\varepsilon_{HA}} & HA
 \end{array}$$

The previous diagram commutes by naturality of ε .

Case.

$$\begin{array}{ccccc}
 & & ?A & & \\
 & \swarrow & \uparrow \mu_A & \searrow & \\
 ?A & \xrightarrow{\eta_{?A}} & ?^2 A & \xleftarrow{? \eta_A} & ?A
 \end{array}$$

It suffices to show that the following diagrams commutes:

$$\begin{array}{ccccc}
 & & JHA & & \\
 & \swarrow & \uparrow J\varepsilon_{HA} & \searrow & \\
 JHA & \xrightarrow{\eta_{JHA}} & JHJHA & \xleftarrow{JH\eta_A} & JHA
 \end{array}$$

Both of these diagrams commute because η and ε are the unit and counit of an adjunction.

It remains to be shown that η and μ are both symmetric comonoidal natural transformations, but this easily follows from the fact that we know η is by assumption, and that μ is because it is defined in terms of ε which is a symmetric comonoidal natural transformation. Thus, all of the following diagrams commute:

$$\begin{array}{ccc}
 A \oplus B & \xrightarrow{\eta_A \oplus \eta_B} & ?A \oplus ?B \\
 \downarrow \eta_A & \nearrow r_{A,B} & \\
 ?(A \oplus B) & &
 \end{array}
 \qquad
 \begin{array}{ccc}
 \perp & \xrightarrow{\eta_\perp} & ?\perp \\
 \downarrow \eta_\perp & \nearrow r_\perp & \\
 \perp & &
 \end{array}$$

$$\begin{array}{ccccc}
 ?^2(A \oplus B) & \xrightarrow{?r_{A,B}} & ?(?A \oplus ?B) & \xrightarrow{r_{?A,?B}} & ?^2 A \oplus ?^2 B \\
 \downarrow \mu_{A \oplus B} & & & & \downarrow \mu_A \oplus \mu_B \\
 ?(A \oplus B) & \xrightarrow{r_{A,B}} & ?A \oplus ?B & &
 \end{array}
 \qquad
 \begin{array}{ccc}
 ?^2 \perp & \xrightarrow{?r_\perp} & ?\perp \\
 \downarrow \mu_\perp & \nearrow r_\perp & \\
 ?\perp & \xrightarrow{r_\perp} & \perp
 \end{array}$$

B.3. Proof of Lemma 26. Suppose (H, h) and (J, j) are two symmetric comonoidal functors, such that, $\mathcal{L} : H \vdash J : C$ is a dual LNL model. Again, we know $?A = H; J : \mathcal{L} \longrightarrow \mathcal{L}$ is a symmetric comonoidal monad by Lemma 25.

We define the following morphisms:

$$\begin{aligned}
 w_A &:= \perp \xrightarrow{j_0^{-1}} J0 \xrightarrow{J\circ_{HA}} JHA = ?A \\
 c_A &:= ?A \oplus ?A = JHA \oplus JHA \xrightarrow{j_{HA,HA}^{-1}} J(HA + HA) \xrightarrow{J\nabla_{HA}} JHA = ?A
 \end{aligned}$$

Next we show that both of these are symmetric comonoidal natural transformations, but for which functors? Define $W(A) = \perp$ and $C(A) = ?A \oplus ?A$ on objects of \mathcal{L} , and $W(f : A \longrightarrow B) = \text{id}_\perp$ and $C(f : A \longrightarrow B) = ?f \oplus ?f$ on morphisms. So we must show that $w : W \longrightarrow ?$ and $c : C \longrightarrow ?$ are symmetric comonoidal natural transformations. We first show that w is and then we show that c is. Throughout the proof we drop subscripts on natural transformations for readability.

Case. To show w is a natural transformation we must show the following diagram commutes for any morphism $f : A \longrightarrow B$:

$$\begin{array}{ccc}
 W(A) & \xrightarrow{w_A} & ?A \\
 \downarrow W(f) & & \downarrow ?f \\
 W(B) & \xrightarrow{w_B} & ?B
 \end{array}$$

This diagram is equivalent to the following:

$$\begin{array}{ccc}
 \perp & \xrightarrow{w_A} & ?A \\
 \text{id}_\perp \downarrow & & \downarrow ?f \\
 \perp & \xrightarrow{w_B} & ?B
 \end{array}$$

It further expands to the following:

$$\begin{array}{ccccc}
 \perp & \xrightarrow{j_0^{-1}} & \mathbf{J}0 & \xrightarrow{J(\diamond_{HA})} & \mathbf{J}HA \\
 \text{id}_\perp \downarrow & & & & \downarrow JHf \\
 \perp & \xrightarrow{j_0^{-1}} & \mathbf{J}0 & \xrightarrow{J(\diamond_{HB})} & \mathbf{J}HB
 \end{array}$$

This diagram commutes, because $J(\diamond_{HA}); Jf = J(\diamond_{HA}; f) = J(\diamond_{HB})$, by the uniqueness of the initial map.

Case. The functor W is comonoidal itself. To see this we must exhibit a map

$$s_\perp := \text{id}_\perp : W \perp \longrightarrow \perp$$

and a natural transformation

$$s_{A,B} := \rho_\perp^{-1} : W(A \oplus B) \longrightarrow WA \oplus WB$$

subject to the coherence conditions in Definition 8. Clearly, the second map is a natural transformation, but we leave showing they respect the coherence conditions to the reader. Now we can show that w is indeed symmetric comonoidal.

Case.

$$\begin{array}{ccc}
 W(A \oplus B) & \xrightarrow{s_{A,B}} & WA \oplus WB \\
 \downarrow w_{A \oplus B} & & \downarrow w_A \oplus w_B \\
 ?(A \oplus B) & \xrightarrow{r_{A,B}} & ?A \oplus ?B
 \end{array}$$

Expanding the objects of the previous diagram results in the following:

$$\begin{array}{ccc}
 \perp & \xrightarrow{s_{A,B}} & \perp \oplus \perp \\
 \downarrow w_{A \oplus B} & & \downarrow w_A \oplus w_B \\
 ?(A \oplus B) & \xrightarrow{r_{A,B}} & ?A \oplus ?B
 \end{array}$$

This diagram commutes, because the following fully expanded diagram commutes:






Diagram 1 commutes because 0 is the initial object, diagram 2 commutes by naturality of j , diagram 3 commutes because J is a symmetric comonoidal functor, diagram 4 commutes because j_0 is an isomorphism (Lemma 23), diagram 5 commutes by functoriality of J , and diagram 6 commutes by naturality of ρ .


Case.



Expanding the objects in the previous diagram results in the following:



This diagram commutes because the following one does:



The diagram on the left commutes because j_0 is an isomorphism (Lemma 23), and the diagram on the right commutes because 0 is the initial object.

Case. Now we show that $c_A : ?A \oplus ?A \longrightarrow ?A$ is a natural transformation. This requires the following diagram to commute (for any $f : A \longrightarrow B$):

$$\begin{array}{ccc} \mathbf{C}A & \xrightarrow{c_A} & ?A \\ \mathbf{C}f \downarrow & & \downarrow ?f \\ \mathbf{C}B & \xrightarrow{c_B} & ?B \end{array}$$

This expands to the following diagram:

$$\begin{array}{ccc} ?A \oplus ?A & \xrightarrow{c_A} & ?A \\ ?f \oplus ?f \downarrow & & \downarrow ?f \\ ?B \oplus ?B & \xrightarrow{c_B} & ?B \end{array}$$

This diagram commutes because the following diagram does:

$$\begin{array}{ccccc} \mathbf{J}HA \oplus \mathbf{J}HA & \xrightarrow{j_{HA,HA}^{-1}} & \mathbf{J}(HA + HA) & \xrightarrow{\mathbf{J}\nabla_{HA}} & \mathbf{J}HA \\ \mathbf{J}Hf \oplus \mathbf{J}Hf \downarrow & & \downarrow \mathbf{J}(Hf + Hf) & & \downarrow \mathbf{J}Hf \\ \mathbf{J}HB \oplus \mathbf{J}HB & \xrightarrow{j_{HB,HB}^{-1}} & \mathbf{J}(HB + HB) & \xrightarrow{\mathbf{J}\nabla_{HB}} & \mathbf{J}HB \end{array}$$

The left square commutes by naturality of j^{-1} , and the right square commutes by naturality of the codiagonal $\nabla_A : A + A \longrightarrow A$.

Case. The functor $\mathbf{C} : \mathcal{L} \longrightarrow \mathcal{L}$ is indeed symmetric comonoidal where the required maps are defined as follows:

$$t_{\perp} := ?\perp \oplus ?\perp \xrightarrow{j^{-1}} \mathbf{J}H\perp \oplus \mathbf{J}H\perp \xrightarrow{\mathbf{J}\nabla} \mathbf{J}H\perp \xrightarrow{\mathbf{J}h_{\perp}} \mathbf{J}0 \xrightarrow{j_0} \perp$$

$$t_{A,B} := ?(A \oplus B) \oplus ?(A \oplus B) \xrightarrow{r_{A,B} \oplus r_{A,B}} (?A \oplus ?B) \oplus (?A \oplus ?B) \xrightarrow{\text{iso}} (?A \oplus ?A) \oplus (?B \oplus ?B)$$

where iso is a natural isomorphism that can easily be defined using the symmetric monoidal structure of \mathcal{L} . Clearly, t is indeed a natural transformation, but we leave checking that the required diagrams in Definition 8 commute to the reader. We can now show that $c_A : ?A \oplus ?A \longrightarrow ?A$ is symmetric comonoidal. The following diagrams from Definition 10 must commute:

Case.

$$\begin{array}{ccc} \mathbf{C}(A \oplus B) & \xrightarrow{t_{A,B}} & \mathbf{C}A \oplus \mathbf{C}B \\ \mathbf{C}_{A \oplus B} \downarrow & & \downarrow \mathbf{C}_A \oplus \mathbf{C}_B \\ ?(A \oplus B) & \xrightarrow{r_{A,B}} & ?A \oplus ?B \end{array}$$

Expanding the objects in the previous diagram results in the following:


$$\begin{array}{ccc}
?(A \oplus B) \oplus ?(A \oplus B) & \xrightarrow{t_{A,B}} & (?A \oplus ?A) \oplus (?B \oplus ?B) \\
\downarrow c_{A \oplus B} & & \downarrow c_A \oplus c_B \\
?(A \oplus B) & \xrightarrow{r_{A,B}} & ?A \oplus ?B
\end{array}$$

This diagram commutes, because the following fully expanded one does:


$$\begin{array}{c}
\begin{array}{c}
\text{JH}(A \oplus B) \oplus \text{JH}(A \oplus B) \xrightarrow{\text{Jh} \oplus \text{Jh}} \text{J}(\text{HA} + \text{HB}) \oplus \text{J}(\text{HA} + \text{HB}) \xrightarrow{\text{j} \oplus \text{j}} (\text{JHA} \oplus \text{JHB}) \oplus (\text{JHA} \oplus \text{JHB}) \xrightarrow{\text{iso}} (\text{JHA} \oplus \text{JHA}) \oplus (\text{JHB} \oplus \text{JHB}) \\
\downarrow \text{j}^{-1} \quad (2) \quad \downarrow \text{j}^{-1} \quad (4) \quad \downarrow \text{j}^{-1} \oplus \text{j}^{-1} \quad (6) \\
\text{J}(\text{H}(A \oplus B) + \text{H}(A \oplus B)) \xrightarrow{\text{J}(\text{h} + \text{h})} \text{J}((\text{HA} + \text{HB}) + (\text{HA} + \text{HB})) \xrightarrow{\text{Jiso}} \text{J}((\text{HA} + \text{HA}) + (\text{HB} + \text{HB})) \xrightarrow{\text{j}} \text{J}(\text{HA} + \text{HA}) \oplus \text{J}(\text{HB} + \text{HB}) \\
\downarrow \text{J}\nabla \quad (1) \quad \downarrow \text{J}\nabla \quad (3) \quad \downarrow \text{J}(\nabla + \nabla) \quad (5) \quad \downarrow \text{J}\nabla \oplus \nabla \\
\text{JH}(A \oplus B) \xrightarrow{\text{Jh}} \text{J}(\text{HA} + \text{HB}) \xlongequal{\quad} \text{J}(\text{HA} + \text{HB}) \xrightarrow{\text{j}} \text{JHA} \oplus \text{JHB}
\end{array}
\end{array}$$

Diagram 1 commutes by naturality of ∇ , diagram 2 commutes by naturality of j^{-1} , diagram 3 commutes by straightforward reasoning on coproducts, diagram 4 commutes by straightforward reasoning on the symmetric monoidal structure of J after expanding the definition of the two isomorphisms – here $J\text{iso}$ is the corresponding isomorphisms on coproducts – diagram 5 commutes by naturality of j , and diagram 6 commutes because j is an isomorphism (Lemma 23).


Case.



Expanding the objects of this diagram results in the following:



Simply unfolding the morphisms in the previous diagram reveals the following:



Clearly, this diagram commutes.

At this point we have shown that $w_A : \perp \rightarrow ? A$ and $c_A : ? A \oplus ? A \rightarrow ? A$ are symmetric comonoidal naturality transformations. Now we show that for any $? A$ the triple $(? A, w_A, c_A)$ forms a commutative monoid. This means that the following diagrams must commute:

Case.

$$\begin{array}{ccccc}
(?A \oplus ?A) \oplus ?A & \xrightarrow{\alpha_{?A, ?A, ?A}} & ?A \oplus (?A \oplus ?A) & \xrightarrow{\text{id}_{?A} \oplus c_A} & ?A \oplus ?A \\
\downarrow c_A \oplus \text{id}_A & & & & \downarrow c_A \\
?A \oplus ?A & \xrightarrow{c_A} & & & ?A
\end{array}$$

The previous diagram commutes, because the following one does (we omit subscripts for readability):

$$\begin{array}{ccccccc}
(JHA \oplus JHA) \oplus JHA & \xrightarrow{\alpha} & JHA \oplus (JHA \oplus JHA) & \xrightarrow{\text{id} \oplus j^{-1}} & JHA \oplus J(HA + HA) & \xrightarrow{\text{id} \oplus J \nabla} & JHA \oplus JHA \\
\downarrow j^{-1} \oplus \text{id} & & (1) & & \downarrow j^{-1} & (2) & \downarrow j^{-1} \\
J(HA + HA) \oplus JHA & \xrightarrow{j^{-1}} & J((HA + HA) + HA) & \xrightarrow{J\alpha} & J(HA + (HA + HA)) & \xrightarrow{J(\text{id} + \nabla)} & J(HA + HA) \\
\downarrow J \nabla \oplus \text{id} & (3) & \downarrow J(\nabla + \text{id}) & (4) & & & \downarrow J \nabla \\
JHA \oplus JHA & \xrightarrow{j^{-1}} & J(HA + HA) & \xrightarrow{J \nabla} & & & JHA
\end{array}$$

Diagram 1 commutes because J is a symmetric monoidal functor (Corollary 24), diagrams 2 and 3 commute by naturality of j^{-1} , and diagram 4 commutes because (HA, \diamond, ∇) is a commutative monoid in \mathcal{C} , but we leave the proof of this to the reader.

Case.

$$\begin{array}{ccc}
?A \oplus \perp & & \\
\downarrow \text{id}_{?A} \oplus w_A & \searrow \rho_{?A} & \\
?A \oplus ?A & \xrightarrow{c_A} & ?A
\end{array}$$

The previous diagram commutes, because the following one does:

$$\begin{array}{ccccc}
JHA \oplus \perp & \xrightarrow{\rho} & JHA & & \\
\downarrow \text{id} \oplus j_0^{-1} & (1) & & & \parallel \\
JHA \oplus J0 & \xrightarrow{j^{-1}} & J(HA + 0) & \xrightarrow{J\rho} & JHA \\
\downarrow \text{id} \oplus J\diamond & (2) & \downarrow J(\text{id} \oplus \diamond) & (3) & \parallel \\
JHA \oplus JHA & \xrightarrow{j^{-1}} & J(HA + HA) & \xrightarrow{J \nabla} & JHA
\end{array}$$

Diagram 1 commutes because J is a symmetric monoidal functor (Corollary 24), diagram 2 commutes by naturality of j^{-1} , and diagram 3 commutes because (HA, \diamond, ∇) is a commutative monoid in \mathcal{C} , but we leave the proof of this to the reader.

Case.

$$\begin{array}{ccc}
?A \oplus ?A & & \\
\downarrow \beta_{?A, ?A} & \searrow c_A & \\
?A \oplus ?A & \xrightarrow{c_A} & ?A
\end{array}$$

This diagram commutes, because the following one does:

$$\begin{array}{ccccc}
JHA \oplus JHA & \xrightarrow{j^{-1}} & J(HA + HA) & \xrightarrow{J\nabla} & JHA \\
\downarrow \beta & & \downarrow J\beta & & \parallel \\
JHA \oplus JHA & \xrightarrow{j^{-1}} & J(HA + HA) & \xrightarrow{J\nabla} & JHA
\end{array}$$

The left diagram commutes by naturality of j^{-1} , and the right diagram commutes because (HA, \diamond, ∇) is a commutative monoid in \mathcal{C} , but we leave the proof of this to the reader.

Finally, we must show that $w_A : \perp \rightarrow ?A$ and $c_A : ?A \oplus ?A \rightarrow ?A$ are $?$ -algebra morphisms. The algebras in play here are $(?A, \mu : ??A \rightarrow ?A)$, $(\perp, r_\perp : ?\perp \rightarrow \perp)$, and $(?A \oplus ?A, u_A : ?(?A \oplus ?A) \rightarrow ?A \oplus ?A)$, where $u_A := ?(?A \oplus ?A) \xrightarrow{r_{?A, ?A}} ?^2A \oplus ?^2A \xrightarrow{\mu_A \oplus \mu_A} ?A \oplus ?A$. It suffices to show that the following diagrams commute:

Case.

$$\begin{array}{ccc}
?\perp & \xrightarrow{r_\perp} & \perp \\
\downarrow ?w & & \downarrow w \\
??A & \xrightarrow{\mu} & ?A
\end{array}$$

This diagram commutes, because the following fully expanded one does:

$$\begin{array}{ccccc}
JH\perp & \xrightarrow{Jh_\perp} & J0 & \xrightarrow{j_0} & \perp \\
\downarrow JHj_0^{-1} & \searrow JHj_0^{-1} & & & \downarrow j_0^{-1} \\
& & JHJ0 & \xrightarrow{JHj_0} & JH\perp \\
& & & & \downarrow Jh_\perp \\
& & & & J0 \\
\downarrow JHJ_\diamond & & \xrightarrow{J\varepsilon_0} & & \downarrow J_\diamond \\
JHJHA & \xrightarrow{J\varepsilon} & & & JHA
\end{array}$$

(1) (2) (3) (4)




Diagram 1 clearly commutes, diagram 2 commutes by naturality of ε , diagram 3 commutes by naturality of ∇ , diagram 4 commutes because ε is the counit of the symmetric comonoidal adjunction, diagram 5 commutes because j is an isomorphism (Lemma 23), diagram 6 commutes by naturality of j^{-1} , and diagram 7 is the same diagram as 3, but this diagram is redundant for readability.

B.4. Proof of Lemma 27. Suppose $\mathcal{L} : H \dashv J : C$ is a dual LNL model. Then we know $?A = JHA$ is a symmetric comonoidal monad by Lemma 25. Bellin [3] remarks that by Maietti, Maneggia de Paiva and Ritter's Proposition 25 [15], it suffices to show that $\mu_A : ??A \rightarrow ?A$ is a monoid morphism. Thus, the following diagrams must commute:

Case.

$$\begin{array}{ccc} ??A \oplus ??A & \xrightarrow{c_{?A}} & ??A \\ \downarrow \mu_A \oplus \mu_A & & \downarrow \mu_A \\ ?A \oplus ?A & \xrightarrow{c_A} & ?A \end{array}$$

This diagram commutes because the following fully expanded one does:

$$\begin{array}{ccccc} JHJHA \oplus JHJHA & \xrightarrow{j^{-1}} & J(HJHA + HJHA) & \xrightarrow{J\nabla} & JHJHA \\ \downarrow J\varepsilon \oplus J\varepsilon & & \downarrow J(\varepsilon + \varepsilon) & & \downarrow J\varepsilon \\ JHA \oplus JHA & \xrightarrow{j^{-1}} & J(HA + HA) & \xrightarrow{J\nabla} & JHA \end{array}$$

The left square commutes by naturality of j^{-1} and the right square commutes by naturality of the codiagonal.

Case.

$$\begin{array}{ccc} & \perp & \\ w_{?A} \swarrow & & \searrow w_A \\ ??A & \xrightarrow{\mu_A} & ?A \end{array}$$

This diagram commutes because the following fully expanded one does:

$$\begin{array}{ccc} \perp & \xlongequal{\quad} & \perp \\ \downarrow j_0^{-1} & & \downarrow j_0^{-1} \\ J0 & \xlongequal{\quad} & J0 \\ \downarrow J\circ & & \downarrow J\circ \\ JHJHA & \xrightarrow{J\varepsilon} & JHA \end{array}$$

The top square trivially commutes, and the bottom square commutes by uniqueness of the initial map.

B.5. Proof of Cut Reduction (Lemma 38). By induction on $d(\Pi_1) + d(\Pi_2)$. We consider only the case where the last inferences of Π_1 and Π_2 are logical inferences. The other cases are handled mainly by permutation of inferences and use of the inductive hypothesis; we refer to Benton's text for them. Throughout the proof we will add an asterisk to the name of an inference rule to indicate that the rule may be applied zero or more times.

J right / J left. We have

$$\Pi_1 = \frac{\pi_1}{A \vdash_L \Delta, JT^n; T, \Psi} J_R \quad \Pi_2 = \frac{T \vdash_C \Psi'}{JT \vdash_L; \Delta, JT^{n+1}, \Psi'} J_L$$

By the inductive hypothesis applied to Π_2 and π_1 there exists a proof Π' of $A \vdash_L \Delta; T, \Psi, \Psi'$ with $c(\Pi') \leq |JT| = |T| + 1$. Then the following derivation

$$\Pi_1 = \frac{\frac{\Pi' \quad \pi_2}{A \vdash_L \Delta; T, \Psi, \Psi' \quad T \vdash_C \Psi'} LC_cut}{\frac{A \vdash_L \Delta, \Psi, \Psi', \Psi'}{A \vdash_L \Delta, \Psi, \Psi'} C_contr^*}$$

has cut rank $\max(|T| + 1, c(\Pi'), c(\pi_2)) = |T| + 1 = |JT|$.

H right / H left. We have

$$\Pi_1 = \frac{\pi_1}{B \vdash_L \Delta, A; HA^n; \Psi} H_R \quad \Pi_2 = \frac{\pi_2}{A \vdash_L; \Psi'} H_L$$

By the inductive hypothesis applied to Π_2 and π_1 there exists a proof Π' of $B \vdash_L \Delta; A, \Psi, \Psi'$ with $c(\Pi') \leq |HA| = |A| + 1$. Then the following derivation

$$\Pi_1 = \frac{\frac{\Pi' \quad \pi_2}{B \vdash_L \Delta; A, \Psi, \Psi' \quad A \vdash_L; \Psi'} LL_cut}{\frac{B \vdash_L \Delta, \Psi, \Psi', \Psi'}{B \vdash_L \Delta, \Psi, \Psi'} C_contr^*}$$

has cut rank $\max(|A| + 1, c(\Pi'), c(\pi_2)) = |A| + 1 = |HA|$.

+ *right₁ / + left.* We have

$$\Pi_1 = \frac{\pi_1}{S \vdash_C T_1, (T_1 + T_2)^n, \Psi} C_{-+R_1} \quad \Pi_2 = \frac{\pi_2 \quad \pi_3}{T_1 \vdash_C \Psi_1 \quad T_2 \vdash_C \Psi_2} C_{-+L}$$

If $n = 0$, then the reduction is as follows:

$$\Pi_1 = \frac{\pi_1}{S \vdash_C T_1, \Psi} C_{-+R_1} \quad \Pi_2 = \frac{\pi_2 \quad \pi_3}{T_1 \vdash_C \Psi_1 \quad T_2 \vdash_C \Psi_2} C_{-+L}$$

$$\frac{S \vdash_C T_1 + T_2, \Psi}{S \vdash_C \Psi, \Psi_1, \Psi_2} C_cut$$

reduces to

$$\frac{\pi_1 \quad \pi_2}{S \vdash_C T_1, \Psi \quad T_1 \vdash_C \Psi_1} C_cut$$

$$\frac{\Pi}{S \vdash_C \Psi, \Psi_1, \Psi_2} C_weak^*$$

Here $c(\Pi) = \max(|T_1| + 1, c(\pi_1), c(\pi_2)) \leq |T_1 + T_2|$.

If $n > 0$, then by the inductive hypothesis applied to Π_2 and π_1 there exists a proof Π' of $S \vdash_{\mathbf{C}} T_1, \Psi, \Psi_1, \Psi_2$ with $c(\Pi') \leq |T_1 + T_2| = |T_1| + |T_2| + 1$. Then the following derivation

$$\Pi = \frac{\frac{\frac{\Pi'}{S \vdash_{\mathbf{C}} T_1, \Psi, \Psi_1, \Psi_2} \quad \frac{\pi_2}{T_1 \vdash_{\mathbf{C}} \Psi_1}}{S \vdash_{\mathbf{C}} \Psi, \Psi_1, \Psi_1, \Psi_2} \text{C_cut}}{S \vdash_{\mathbf{C}} \Psi, \Psi_1, \Psi_2} \text{C_contr}^*$$

has cut rank $\max(|T_1| + 1, c(\Pi'), c(\pi_2)) \leq |T_1 + T_2|$.

•– right / •– left. We have

$$\Pi_1 = \frac{\frac{\frac{\pi_1}{A \vdash_{\mathbf{L}} \Delta_1; \Psi_1, B_1} \quad \frac{\pi_2}{B_2 \vdash_{\mathbf{L}} \Delta_2; \Psi_2}}{A \vdash_{\mathbf{L}} B_1 \bullet B_2, \Delta_1, \Delta_2; \Psi_1, \Psi_2} \text{LL_}\bullet\text{-}_R \quad \Pi_2 = \frac{\frac{\pi_3}{B_1 \vdash_{\mathbf{L}} B_2, \Delta; \Psi}}{B_1 \bullet B_2 \vdash_{\mathbf{L}} \Delta; \Psi} \text{LL_}\bullet\text{-}_L}{A \vdash_{\mathbf{L}} \Delta_1, \Delta_2, \Delta; \Psi_1, \Psi_2, \Psi} \text{LL_cut}$$

reduces to Π

$$\frac{\frac{\frac{\pi_1}{A \vdash_{\mathbf{L}} \Delta_1, B_1; \Psi_1} \quad \frac{\pi_3}{B_1 \vdash_{\mathbf{L}} B_2, \Delta; \Psi}}{A \vdash_{\mathbf{L}} \Delta_1, \Delta, B_2; \Psi_1, \Psi} \text{LL_cut} \quad \frac{\pi_2}{B_2 \vdash_{\mathbf{L}} \Delta_2; \Psi_2} \text{LL_cut}}{A \vdash_{\mathbf{L}} \Delta_1, \Delta_2, \Delta; \Psi_1, \Psi_2, \Psi} \text{LL_cut}$$

The resulting derivation Π has cut rank $c(\Pi) = \max(|B_1| + 1, c(\pi_1), c(\pi_2), |B_2| + 1, c(\pi_3)) \leq |B_1 \bullet B_2|$.