

The **mandi** Bundle

Paul J. Heafner*

Printed September 12, 2024

mandi version v3.2.2a dated 2024-09-12
mandistudent version v3.2.2a dated 2024-09-12
mandiexp version v3.2.2a dated 2024-09-12

*Email: heafnerj@gmail.com

Contents

Acknowledgements	3
Change History	4
List of Web VPython Programs	5
List of VPython Programs	5
List of Figures	5
1 Introduction	6
2 Getting Help	6
3 Code Availability	6
4 Compatibility with Other Packages and Classes	6
5 Student/Instructor Quick Guide	7
6 The mandi Package	8
6.1 Package Options	8
6.2 The <code>mandisetup</code> Command	8
6.3 Lua \LaTeX is Required	8
6.4 Physical Quantities	9
6.4.1 Typesetting Physical Quantities	9
6.4.2 Checking Physical Quantities	10
6.4.3 Predefined Physical Quantities	10
6.4.4 Defining and Redefining Physical Quantities	22
6.4.5 Changing Units	22
6.5 Physical Constants	23
6.5.1 Typesetting Physical Constants	23
6.5.2 Checking Physical Constants	24
6.5.3 Predefined Physical Constants	24
6.5.4 Defining and Redefining Physical Constants	30
6.5.5 Changing Precision	31
6.6 Predefined Units and Constructs	31
6.7 <code>mandi</code> Source Code	35
7 The mandistudent Package	52
7.1 Traditional Vector Notation	52
7.2 Problems and Annotated Problem Solutions	56
7.3 Coordinate-Free and Index Notation	60
7.4 Web VPython and VPython Program Listings	62
7.5 The <code>webvpythonblock</code> Environment	62
7.6 The <code>vpythonfile</code> Command	68
7.7 The <code>webvpythoninline</code> and <code>vpythoninline</code> Commands	70
7.8 <code>mandistudent</code> Source Code	71
8 The mandiexp Package	83
8.1 The Fundamenal Principles	83
8.2 Other Symbols	85
8.3 <code>mandiexp</code> Source Code	88
Index	95

Acknowledgements

To all of the students who have learned $\text{\LaTeX} 2_{\epsilon}$ in my introductory physics courses over the years, I say a heartfelt thank you. You have contributed directly to the state of this software and to its use in introductory physics courses and to innovating how physics is taught.

I also acknowledge the $\text{\LaTeX} 2_{\epsilon}$ developers who inhabit the [TeX StackExchange](#) site. Entering a new culture is daunting for anyone, especially for newcomers; the $\text{\LaTeX} 2_{\epsilon}$ development culture is no exception. We all share a passion for creating beautiful documents and I learned much over the summers of 2020 and 2021 that improved my ability to do just that. There are too many of you to list individually, and I would surely accidentally omit some were I to try. Collectively, I thank you all for your patience and advice.

Change History

v3.0.0 – 2021-08-22

[mandi](#) ^{→ P.8} [mandistudent](#) ^{→ P.52} [mandiexp](#) ^{→ P.83}
Initial release 6

v3.0.1 – 2021-08-24

[mandi](#) ^{→ P.8} [mandistudent](#) ^{→ P.52}
[mandiexp](#) ^{→ P.83} Minor doc changes 6

v3.1.0 – 2022-01-27

[mandi](#) ^{→ P.8} Added GitHub links to code 6
[mandi](#) ^{→ P.8} Added [\hbar](#) ^{→ P.33} 46
[mandi](#) ^{→ P.8} Added [\lorentzfactor](#) ^{→ P.16} 43
[mandi](#) ^{→ P.8} Added a negative space to
[\lightspeed](#) ^{→ P.32} 37
[mandi](#) ^{→ P.8} Constants' values now use only
[\times](#) 46
[mandi](#) ^{→ P.8} Improved [\CheckConstant](#) ^{→ P.24} 50
[mandi](#) ^{→ P.8} Improved [\CheckQuantity](#) ^{→ P.10} 50
[mandi](#) ^{→ P.8} Unknown package options handled
safely 36
[mandi](#) ^{→ P.8} [mandistudent](#) ^{→ P.52} L^AT_EX3 code now
conforms to formatting standards 6
[mandi](#) ^{→ P.8} [mandistudent](#) ^{→ P.52} [mandiexp](#) ^{→ P.83}
Code formatted for better readability 6
[mandi](#) ^{→ P.8} [mandistudent](#) ^{→ P.52} [mandiexp](#) ^{→ P.83}
xparse is loaded for older formats 6
[mandi](#) ^{→ P.8} [\mivector](#) ^{→ P.34} now requires more
than one component 50
[mandistudent](#) ^{→ P.52} All instances of `GlowScript`
have been changed to `Web VPython` 62
[mandistudent](#) ^{→ P.52} Default URL for
[\vpythonfile](#) ^{→ P.68} is now `vpython.org` 81
[mandistudent](#) ^{→ P.52} Slightly modified
[\image](#) ^{→ P.59} 77
[mandistudent](#) ^{→ P.52} URLs fixed in
[\vpythonfile](#) ^{→ P.68} 81
[mandistudent](#) ^{→ P.52} URLs fixed in
[webvpythonblock](#) ^{→ P.62} 80
[mandistudent](#) ^{→ P.52} Version number works 71
[mandistudent](#) ^{→ P.52} [\diff](#) renamed to [\df](#) ^{→ P.62}
for compatibility with the `numerica` package 62
[mandistudent](#) ^{→ P.52} [\vec](#) ^{→ P.52} and [\dirvec](#) ^{→ P.52}
no longer add `\scriptspace` when no

sub/superscript is given 74

[mandistudent](#) ^{→ P.52} [webvpythonblock](#) ^{→ P.62} is a
variant of [webvpythonblock](#) ^{→ P.62} that omits
the QR code 81

[mandistudent](#) ^{→ P.52} [webvpythonblock](#) ^{→ P.62} now
automatically generates QR codes for
program listings 80

[mandiexp](#) ^{→ P.83} Version number works 88

v3.2.0 – 2023-08-01

[mandi](#) ^{→ P.8} Changed [\unit](#) to [\units](#) 37
[mandi](#) ^{→ P.8} [mandistudent](#) ^{→ P.52} [mandiexp](#) ^{→ P.83}

Source updated 6

[mandistudent](#) ^{→ P.52} Added statement about
BEAMER compatibility 6

[mandistudent](#) ^{→ P.52} [\hilite](#) ^{→ P.58} is not defined
if BEAMER is loaded 77

[mandistudent](#) ^{→ P.52} [\vpythonfile](#) ^{→ P.68} now
shows URL in header 81

[mandistudent](#) ^{→ P.52} [physicsproblem](#) ^{→ P.56} and
[physicsproblem](#) ^{→ P.56} are not defined if
BEAMER is loaded 76

[mandistudent](#) ^{→ P.52} [webvpythonblock](#) ^{→ P.62} and
[webvpythonblock](#) ^{→ P.62} now show URL in
header 80

[mandiexp](#) ^{→ P.83} [mandistudent](#) is loaded for [\vec](#) ^{*} 88

v3.2.1 – 2023-11-22

[mandi](#) ^{→ P.8} Minor changes to `mandi.ins` inside
`mandi.dtx` 6

[mandi](#) ^{→ P.8} [mandistudent](#) ^{→ P.52} [mandiexp](#) ^{→ P.83}
Source updated and bumped version for new
CTAN upload 6

Improved `README.md` at request of CTAN
maintainer 6

v3.2.2 – 2024-02-27

[mandi](#) ^{→ P.8} Ensured all subscripts and
superscripts are braced 6

[mandi](#) ^{→ P.8} Removed unnecessary `\ExplSyntaxOn`
and `\ExplSyntaxOff` pairs 6

v3.2.2a – 2024-09-12

[mandistudent](#) ^{→ P.52} Removed a % to keep a line
from appearing in the `.sty` file. 75

List of Web VPython Programs

1	Example With QR Code	64
2	Example Without QR Code	67

List of VPython Programs

1	A VPython Program	69
---	-----------------------------	----

List of Figures

1	Image shown 20 percent actual size.	60
2	Image shown 20 percent actual size and rotated.	60

1 Introduction

The `mandi`¹ bundle consists of three packages: `mandi`, `mandistudent`, and `mandiexp`. Package `mandi`^{→ P.8} provides the core functionality, namely correctly typesetting physical quantities and constants with their correct SI units as either scalars or vectors, depending on which is appropriate. Package `mandistudent`^{→ P.52} provides other typesetting capability appropriate for written problem solutions. Finally, package `mandiexp`^{→ P.83} provides commands for typesetting expressions from *Matter & Interactions*²

So many changes have been made that I think the best approach for all users is to treat this as a brand new experience. I think the most important thing to keep in mind is that I assume all users will have a relatively recent T_EX distribution (like T_EX Live) that includes an updated kernel. If users report that this is a major problem, I can provide some degree of backwards compatibility. However, I use a fully updated T_EX Live distribution.

2 Getting Help

If you have a question about `mandi`, first please read this documentation to make sure your question is not addressed here. Then if you wish, email me. As a user, you deserve courteous and timely help if you need it. You will never get a response from me saying that because this software is free you are not entitled to help using it. It is sad that some developers take that attitude. Using free software does not absolve the developer of helping users unless the license specifically says so.

3 Code Availability

The `mandi` source repository's `main` branch is at <https://github.com/heafnerj/mandi>. This code will usually coincide with that found on CTAN. The very latest build can be found on the `dev` branch found at <https://github.com/heafnerj/mandi/tree/dev>. Students and other academic users should probably get the `dev` branch code since it is stable and may contain improvements over the `main` branch code.

4 Compatibility with Other Packages and Classes

When using BEAMER, certain commands in the `mandistudent` package are not defined. Because BEAMER uses the `enumitem` package, the `physicsproblem`^{→ P.56} and `physicsproblem*`^{→ P.56} environments are not defined. The `\hilite`^{→ P.58} command is also not defined because it does not work and I do not currently know why.

¹The bundle name can be pronounced either with two syllables, to rhyme with *candy*, or with three syllables, as *M and I*.

²See *Matter & Interactions* and <https://matterandinteractions.org/> for details.

5 Student/Instructor Quick Guide

Use `\vec`^{P.52} to typeset the symbol for a vector. Use `\magnitude`^{P.55} to typeset the symbol for a vector's magnitude. Use `\dirvec`^{P.52} to typeset the symbol for a vector's direction. Use `\changein`^{P.53} to typeset the symbol for the change in a vector or scalar. Use `\zerovec`^{P.53} to typeset the zero vector. Use `\timestento`^{P.34} to typeset scientific notation.

<code>\(\vec{p} \)</code> or <code>\(\vec{*p} \)</code>	<code>\</code>	p or \vec{p}
<code>\(\vec{p}_{\text{final}} \)</code> or <code>\(\vec{*p}_{\text{final}} \)</code>	<code>\</code>	p_{final} or \vec{p}_{final}
<code>\(\text{magnitude}\vec{p} \)</code> or <code>\(\text{magnitude*}\vec{p}_{\text{final}} \)</code>	<code>\</code>	$\ p\ $ or $\ p_{\text{final}}\ $
<code>\(\text{dirvec}\vec{p} \)</code> or <code>\(\text{dirvec*}\vec{p} \)</code>	<code>\</code>	\hat{p} or \hat{p}
<code>\(\text{changein}\vec{p} \)</code> or <code>\(\text{changein } t \)</code>	<code>\</code>	Δp or Δt
<code>\(\text{zerovec} \)</code> or <code>\(\text{zerovec*} \)</code>	<code>\</code>	$\mathbf{0}$ or $\vec{0}$
<code>\(6.02\text{timestento}\{-19\} \)</code>		6.02×10^{-19}

Use a `physical quantity's`^{P.9} name to typeset a magnitude and that quantity's units. If the quantity is a vector, you can add `vector` either to the beginning or the end of the quantity's name. For example, if you want momentum, use `\momentum`^{P.9} and its variants.

<code>\(\text{momentum}\{7.071\} \)</code>	<code>\</code>	$7.071 \text{ kg} \cdot \text{m/s}$
<code>\(\text{vectormomentum}\{3,-4,5\} \)</code>	<code>\</code>	$\langle 3, -4, 5 \rangle \text{ kg} \cdot \text{m/s}$
<code>\(\text{momentumvector}\{3,-4,5\} \)</code>		$\langle 3, -4, 5 \rangle \text{ kg} \cdot \text{m/s}$

Use a `physical constant's`^{P.23} name to typeset its numerical value and units. Append `mathsymbol` to the constant's name to get its mathematical symbol. For example, if you want to typeset the vacuum permittivity, use `\vacuumpermittivity`^{P.30} and its variant.

<code>\(\text{vacuumpermittivitymathsymbol} = \text{vacuumpermittivity} \)</code>	<code>\</code>	$\epsilon_0 = 9 \times 10^{-12} \text{ C}^2/\text{N} \cdot \text{m}^2$
--	----------------	--

Use `\mivector`^{P.34} to typeset symbolic vectors with components. Use the alias `\direction`^{P.13} to typeset a direction or unit vector.

<code>\(\text{mivector}\{\text{slot},\text{slot},\text{slot}\} \)</code> or <code>\(\text{mivector}\{p_x,p_y,p_z\} \)</code>	<code>\</code>	
<code>\(\text{direction}\{\frac{1}{\sqrt{3}},\frac{1}{\sqrt{3}},\frac{1}{\sqrt{3}}\} \)</code> or		
$\langle \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}} \rangle$ or $\langle p_x, p_y, p_z \rangle$		

Use `physicsproblem`^{P.56} and `parts`^{P.56} and `\problempart`^{P.56} for problems. For step-by-step mathematical solutions use `physicssolution`^{P.57}. Use `webvpythonblock`^{P.62} to typeset Web VPython programs. Use `\vpythonfile`^{P.68} to typeset VPython program files.

6 The `mandi` Package

Load `mandi` as you would any package in your preamble.

```
\usepackage[options]{mandi}
```

`\mandiversion`

Typesets the current version and build date.

The version is `\mandiversion{}` and is a stable build.

The version is v3.2.2a dated 2024-09-12 and is a stable build.

6.1 Package Options

N 2021-01-30

`units`=*<type of unit>* (initially unspecified, set to **alternate**)

N 2021-01-30

`preciseconstants`=*<boolean>* (initially unspecified, set to **false**)

Now `mandi` uses a key-value interface for options. The `units` key can be set to **base**, **derived**, or **alternate**. The `preciseconstants` key is always either **true** or **false**.

6.2 The `mandisetaup` Command

N 2021-02-17

`\mandisetaup{<options>}`

Command to set package options on the fly after loadtime. This can be done in the preamble or inside the `\begin{document}... \end{document}` environment.

```
\mandisetaup{units=base}
```

```
\mandisetaup{preciseconstants}
```

```
\mandisetaup{preciseconstants = false}
```

6.3 Lua^AT_EX is Required

In order to make use of better fonts and Unicode features, `mandi` now requires the Lua^AT_EX engine for processing documents. It will not work with other engines.

6.4 Physical Quantities

6.4.1 Typesetting Physical Quantities

Typesetting physical quantities and constants using semantically appropriate names, along with the correct [SI units](#), is the core function of `mandi`. Take momentum as the prototypical physical quantity in an introductory physics course.

```
\momentum{\langle magnitude \rangle}
\momentumvector{\langle c_1, \dots, c_n \rangle}
\vectormomentum{\langle c_1, \dots, c_n \rangle}
```

Command for momentum and its vector variants. The default units will depend on the options passed to `mandi` at load time. Alternate units are the default. Other units can be forced as demonstrated. The vector variants can take more than three components. Note the other variants for the quantity's value and units.

<code>\momentum{5}</code>	<code>\momentumvalue{5}</code>	<code>\momentumbaseunits{5}</code>	<code>\momentumderivedunits{5}</code>	<code>\momentumalternateunits{5}</code>	<code>\momentumvector{2,3,4}</code>	<code>\vectormomentum{2,3,4}</code>	<code>\momentum{\mivector{2,3,4}}</code>	<code>\momentumonlybaseunits</code>	<code>\momentumonlyderivedunits</code>	<code>\momentumonlyalternateunits</code>	<code>\momentumvectorvalue{2,3,4}</code>	<code>\vectormomentumvalue{2,3,4}</code>	<code>\momentumvectorbaseunits{2,3,4}</code>	<code>\vectormomentumbaseunits{2,3,4}</code>	<code>\momentumvectorderivedunits{2,3,4}</code>	<code>\vectormomentumderivedunits{2,3,4}</code>	<code>\momentumvectoralternateunits{2,3,4}</code>	<code>\vectormomentumalternateunits{2,3,4}</code>	<code>\momentumvectoronlybaseunits</code>	<code>\vectormomentumonlybaseunits</code>	<code>\momentumvectoronlyderivedunits</code>	<code>\vectormomentumonlyderivedunits</code>	<code>\momentumvectoronlyalternateunits</code>	<code>\vectormomentumonlyalternateunits</code>
<code>\momentum{5}</code>	<code>\momentumvalue{5}</code>	<code>\momentumbaseunits{5}</code>	<code>\momentumderivedunits{5}</code>	<code>\momentumalternateunits{5}</code>	<code>\momentumvector{2,3,4}</code>	<code>\vectormomentum{2,3,4}</code>	<code>\momentum{\mivector{2,3,4}}</code>	<code>\momentumonlybaseunits</code>	<code>\momentumonlyderivedunits</code>	<code>\momentumonlyalternateunits</code>	<code>\momentumvectorvalue{2,3,4}</code>	<code>\vectormomentumvalue{2,3,4}</code>	<code>\momentumvectorbaseunits{2,3,4}</code>	<code>\vectormomentumbaseunits{2,3,4}</code>	<code>\momentumvectorderivedunits{2,3,4}</code>	<code>\vectormomentumderivedunits{2,3,4}</code>	<code>\momentumvectoralternateunits{2,3,4}</code>	<code>\vectormomentumalternateunits{2,3,4}</code>	<code>\momentumvectoronlybaseunits</code>	<code>\vectormomentumonlybaseunits</code>	<code>\momentumvectoronlyderivedunits</code>	<code>\vectormomentumonlyderivedunits</code>	<code>\momentumvectoronlyalternateunits</code>	<code>\vectormomentumonlyalternateunits</code>
$5 \text{ kg} \cdot \text{m/s}$	5	$5 \text{ kg} \cdot \text{m} \cdot \text{s}^{-1}$	$5 \text{ kg} \cdot \text{m/s}$	$5 \text{ kg} \cdot \text{m/s}$	$\langle 2, 3, 4 \rangle \text{ kg} \cdot \text{m/s}$	$\langle 2, 3, 4 \rangle \text{ kg} \cdot \text{m/s}$	$\text{kg} \cdot \text{m} \cdot \text{s}^{-1}$	$\text{kg} \cdot \text{m/s}$	$\text{kg} \cdot \text{m/s}$	$\langle 2, 3, 4 \rangle$	$\langle 2, 3, 4 \rangle$	$\langle 2, 3, 4 \rangle \text{ kg} \cdot \text{m} \cdot \text{s}^{-1}$	$\langle 2, 3, 4 \rangle \text{ kg} \cdot \text{m} \cdot \text{s}^{-1}$	$\langle 2, 3, 4 \rangle \text{ kg} \cdot \text{m/s}$	$\langle 2, 3, 4 \rangle \text{ kg} \cdot \text{m/s}$	$\langle 2, 3, 4 \rangle \text{ kg} \cdot \text{m/s}$	$\langle 2, 3, 4 \rangle \text{ kg} \cdot \text{m/s}$	$\langle 2, 3, 4 \rangle \text{ kg} \cdot \text{m/s}$	$\text{kg} \cdot \text{m} \cdot \text{s}^{-1}$	$\text{kg} \cdot \text{m} \cdot \text{s}^{-1}$	$\text{kg} \cdot \text{m/s}$	$\text{kg} \cdot \text{m/s}$	$\text{kg} \cdot \text{m/s}$	$\text{kg} \cdot \text{m/s}$

Commands that include the name of a physical quantity typeset units, so they shouldn't be used for algebraic or symbolic values of components. Don't use `\vectormomentum{mv_{x},mv_{y},mv_{z}}` but instead the generic `\mivector{mv_{x},mv_{y},mv_{z}}` instead.

6.4.2 Checking Physical Quantities

U 2022-01-27

\CheckQuantity{ $\langle name \rangle$ }

Command to check and typeset the command, base units, derived units, and alternate units of a defined physical quantity.

6.4.3 Predefined Physical Quantities

Every other defined physical quantity can be treated similarly. Just replace `momentum` with the quantity's name. Obviously, the variants that begin with `\vector` will not be defined for scalar quantities. Here are all the physical quantities, with all their units, defined in `mandi`. Remember that units are not present with symbolic (algebraic) quantities, so do not use the `\vector` variants of these commands for symbolic components. Use `\mivector`^{P.34} instead.

N 2021-02-24

\acceleration{ $\langle magnitude \rangle$ }

\accelerationvector{ $\langle c_1, \dots, c_n \rangle$ }

\vectoracceleration{ $\langle c_1, \dots, c_n \rangle$ }

command

base

$\text{m} \cdot \text{s}^{-2}$

\acceleration

derived

N/kg

alternate

m/s^2

\amount{ $\langle magnitude \rangle$ }

command

base

mol

\amount

derived

mol

alternate

mol

N 2021-02-24

\angularacceleration{ $\langle magnitude \rangle$ }

\angularaccelerationvector{ $\langle c_1, \dots, c_n \rangle$ }

\vectorangularacceleration{ $\langle c_1, \dots, c_n \rangle$ }

command

base

$\text{rad} \cdot \text{s}^{-2}$

\angularacceleration

derived

rad/s^2

alternate

rad/s^2

\angularfrequency{ $\langle magnitude \rangle$ }

command

base

$\text{rad} \cdot \text{s}^{-1}$

\angularfrequency

derived

rad/s

alternate

rad/s

N 2021-02-24

\angularimpulse{ $\langle magnitude \rangle$ }

\angularimpulsevector{ $\langle c_1, \dots, c_n \rangle$ }

\vectorangularimpulse{ $\langle c_1, \dots, c_n \rangle$ }

command

base

$\text{kg} \cdot \text{m}^2 \cdot \text{s}^{-1}$

\angularimpulse

derived

$\text{kg} \cdot \text{m}^2/\text{s}$

alternate

$\text{kg} \cdot \text{m}^2/\text{s}$

N 2021-02-24

`\angularmomentum{\langle magnitude \rangle}`
`\angularmomentumvector{\langle c_1, \dots, c_n \rangle}`
`\vectorangularmomentum{\langle c_1, \dots, c_n \rangle}`

command	<code>\angularmomentum</code>	
base	derived	alternate
$\text{kg} \cdot \text{m}^2 \cdot \text{s}^{-1}$	$\text{kg} \cdot \text{m}^2/\text{s}$	$\text{kg} \cdot \text{m}^2/\text{s}$

N 2021-02-24

`\angularvelocity{\langle magnitude \rangle}`
`\angularvelocityvector{\langle c_1, \dots, c_n \rangle}`
`\vectorangularvelocity{\langle c_1, \dots, c_n \rangle}`

command	<code>\angularvelocity</code>	
base	derived	alternate
$\text{rad} \cdot \text{s}^{-1}$	rad/s	rad/s

`\area{\langle magnitude \rangle}`

command	<code>\area</code>	
base	derived	alternate
m^2	m^2	m^2

`\areachargedensity{\langle magnitude \rangle}`

command	<code>\areachargedensity</code>	
base	derived	alternate
$\text{A} \cdot \text{s} \cdot \text{m}^{-2}$	C/m^2	C/m^2

`\areamassdensity{\langle magnitude \rangle}`

command	<code>\areamassdensity</code>	
base	derived	alternate
$\text{kg} \cdot \text{m}^{-2}$	kg/m^2	kg/m^2

`\capacitance{\langle magnitude \rangle}`

command	<code>\capacitance</code>	
base	derived	alternate
$\text{A}^2 \cdot \text{s}^4 \cdot \text{kg}^{-1} \cdot \text{m}^{-2}$	F	C/V

`\charge{\langle magnitude \rangle}`

command	<code>\charge</code>	
base	derived	alternate
$\text{A} \cdot \text{s}$	C	C

\cmagneticfield $\{\langle magnitude \rangle\}$ **\cmagneticfieldvector** $\{\langle c_1, \dots, c_n \rangle\}$ **\vectorcmagneticfield** $\{\langle c_1, \dots, c_n \rangle\}$ **command****base** $\text{kg} \cdot \text{m} \cdot \text{A}^{-1} \cdot \text{s}^{-3}$ **\cmagneticfield****derived**

N/C

alternate

N/C

\conductance $\{\langle magnitude \rangle\}$ **command****base** $\text{A}^2 \cdot \text{s}^3 \cdot \text{kg}^{-1} \cdot \text{m}^{-2}$ **\conductance****derived**

S

alternate

A/V

\conductivity $\{\langle magnitude \rangle\}$ **command****base** $\text{A}^2 \cdot \text{s}^3 \cdot \text{kg}^{-1} \cdot \text{m}^{-3}$ **\conductivity****derived**

S/m

alternate

A/V · m

\conventionalcurrent $\{\langle magnitude \rangle\}$ **command****base**

A

\conventionalcurrent**derived**

C/s

alternate

A

\current $\{\langle magnitude \rangle\}$ **command****base**

A

\current**derived**

A

alternate

A

\currentdensity $\{\langle magnitude \rangle\}$ **\currentdensityvector** $\{\langle c_1, \dots, c_n \rangle\}$ **\vectorcurrentdensity** $\{\langle c_1, \dots, c_n \rangle\}$ **command****base** $\text{A} \cdot \text{m}^{-2}$ **\currentdensity****derived**C/s · m²**alternate**A/m²**\dielectricconstant** $\{\langle magnitude \rangle\}$ **command****base****\dielectricconstant****derived****alternate**

N 2021-02-24

`\direction{\langle magnitude \rangle}`
`\directionvector{\langle c_1, \dots, c_n \rangle}`
`\vectordirection{\langle c_1, \dots, c_n \rangle}`

command	<code>\direction</code>	
base	derived	alternate

N 2021-02-24

`\displacement{\langle magnitude \rangle}`
`\displacementvector{\langle c_1, \dots, c_n \rangle}`
`\vectordisplacement{\langle c_1, \dots, c_n \rangle}`

command	<code>\displacement</code>	
base	derived	alternate
m	m	m

`\duration{\langle magnitude \rangle}`

command	<code>\duration</code>	
base	derived	alternate
s	s	s

N 2021-02-24

`\electricdipolemoment{\langle magnitude \rangle}`
`\electricdipolemomentvector{\langle c_1, \dots, c_n \rangle}`
`\vectorelectricdipolemoment{\langle c_1, \dots, c_n \rangle}`

command	<code>\electricdipolemoment</code>	
base	derived	alternate
A · s · m	C · m	C · m

N 2021-02-24

`\electricfield{\langle magnitude \rangle}`
`\electricfieldvector{\langle c_1, \dots, c_n \rangle}`
`\vectorelectricfield{\langle c_1, \dots, c_n \rangle}`

command	<code>\electricfield</code>	
base	derived	alternate
kg · m · A ⁻¹ · s ⁻³	V/m	N/C

`\electricflux{\langle magnitude \rangle}`

command	<code>\electricflux</code>	
base	derived	alternate
kg · m ³ · A ⁻¹ · s ⁻³	V · m	N · m ² /C

`\electricpotential{\langle magnitude \rangle}`

command	<code>\electricpotential</code>	
base	derived	alternate
$\text{kg} \cdot \text{m}^2 \cdot \text{A}^{-1} \cdot \text{s}^{-3}$	V	V

N 2021-05-01

`\electricpotentialdifference`{ $\langle magnitude \rangle$ }

command	<code>\electricpotentialdifference</code>	
base	derived	alternate
$\text{kg} \cdot \text{m}^2 \cdot \text{A}^{-1} \cdot \text{s}^{-3}$	V	V

`\electroncurrent`{ $\langle magnitude \rangle$ }

command	<code>\electroncurrent</code>	
base	derived	alternate
s^{-1}	e/s	e/s

`\emf`{ $\langle magnitude \rangle$ }

command	<code>\emf</code>	
base	derived	alternate
$\text{kg} \cdot \text{m}^2 \cdot \text{A}^{-1} \cdot \text{s}^{-3}$	V	V

`\energy`{ $\langle magnitude \rangle$ }

command	<code>\energy</code>	
base	derived	alternate
$\text{kg} \cdot \text{m}^2 \cdot \text{s}^{-2}$	J	J

N 2021-04-15

`\energyinev`{ $\langle magnitude \rangle$ }

command	<code>\energyinev</code>	
base	derived	alternate
eV	eV	eV

N 2021-04-15

`\energyinkev`{ $\langle magnitude \rangle$ }

command	<code>\energyinkev</code>	
base	derived	alternate
keV	keV	keV

N 2021-04-15

`\energyinmev`{ $\langle magnitude \rangle$ }

command	<code>\energyinmev</code>	
base	derived	alternate
MeV	MeV	MeV

`\energydensity`{ $\langle magnitude \rangle$ }

command	\energydensity	
base	derived	alternate
$\text{kg} \cdot \text{m}^{-1} \cdot \text{s}^{-2}$	J/m^3	J/m^3

\energyflux $\{\langle magnitude \rangle\}$
\energyfluxvector $\{\langle c_1, \dots, c_n \rangle\}$
\vectorenergyflux $\{\langle c_1, \dots, c_n \rangle\}$

N 2021-02-24

command	\energyflux	
base	derived	alternate
$\text{kg} \cdot \text{s}^{-3}$	W/m^2	W/m^2

\entropy $\{\langle magnitude \rangle\}$

command	\entropy	
base	derived	alternate
$\text{kg} \cdot \text{m}^2 \cdot \text{s}^{-2} \cdot \text{K}^{-1}$	J/K	J/K

\force $\{\langle magnitude \rangle\}$
\forcevector $\{\langle c_1, \dots, c_n \rangle\}$
\vectorforce $\{\langle c_1, \dots, c_n \rangle\}$

N 2021-02-24

command	\force	
base	derived	alternate
$\text{kg} \cdot \text{m} \cdot \text{s}^{-2}$	N	N

\frequency $\{\langle magnitude \rangle\}$

command	\frequency	
base	derived	alternate
s^{-1}	Hz	Hz

\gravitationalfield $\{\langle magnitude \rangle\}$
\gravitationalfieldvector $\{\langle c_1, \dots, c_n \rangle\}$
\vectorgravitationalfield $\{\langle c_1, \dots, c_n \rangle\}$

N 2021-02-24

command	\gravitationalfield	
base	derived	alternate
$\text{m} \cdot \text{s}^{-2}$	N/kg	N/kg

\gravitationalpotential $\{\langle magnitude \rangle\}$

command	\gravitationalpotential	
base	derived	alternate
$\text{m}^2 \cdot \text{s}^{-2}$	J/kg	J/kg

\gravitationalpotentialdifference $\{\langle magnitude \rangle\}$

N 2021-05-01

command	<code>\gravitationalpotentialdifference</code>	
base	derived	alternate
$\text{m}^2 \cdot \text{s}^{-2}$	J/kg	J/kg

N 2021-02-24

`\impulse{\langle magnitude \rangle}`
`\impulsevector{\langle c_1, \dots, c_n \rangle}`
`\vectorimpulse{\langle c_1, \dots, c_n \rangle}`

command	<code>\impulse</code>	
base	derived	alternate
$\text{kg} \cdot \text{m} \cdot \text{s}^{-1}$	N · s	N · s

`\indexofrefraction{\langle magnitude \rangle}`

command	<code>\indexofrefraction</code>	
base	derived	alternate

`\inductance{\langle magnitude \rangle}`

command	<code>\inductance</code>	
base	derived	alternate
$\text{kg} \cdot \text{m}^2 \cdot \text{A}^{-2} \cdot \text{s}^{-2}$	H	V · s/A

`\linearchargedensity{\langle magnitude \rangle}`

command	<code>\linearchargedensity</code>	
base	derived	alternate
$\text{A} \cdot \text{s} \cdot \text{m}^{-1}$	C/m	C/m

`\linearmassdensity{\langle magnitude \rangle}`

command	<code>\linearmassdensity</code>	
base	derived	alternate
$\text{kg} \cdot \text{m}^{-1}$	kg/m	kg/m

N 2022-01-27

`\lorentzfactor{\langle magnitude \rangle}`

command	<code>\lorentzfactor</code>	
base	derived	alternate

U 2021-05-02

`\luminousintensity{\langle magnitude \rangle}`

command	<code>\luminousintensity</code>	
base	derived	alternate
cd	cd	cd

\magneticcharge{ $\langle magnitude \rangle$ }

command	\magneticcharge	
base	derived	alternate
A · m	A · m	A · m

\magneticdipolemoment{ $\langle magnitude \rangle$ }

\magneticdipolemomentvector{ $\langle c_1, \dots, c_n \rangle$ }

\vectormagneticdipolemoment{ $\langle c_1, \dots, c_n \rangle$ }

command	\magneticdipolemoment	
base	derived	alternate
A · m ²	A · m ²	J/T

\magneticfield{ $\langle magnitude \rangle$ }

\magneticfieldvector{ $\langle c_1, \dots, c_n \rangle$ }

\vectormagneticfield{ $\langle c_1, \dots, c_n \rangle$ }

command	\magneticfield	
base	derived	alternate
kg · A ⁻¹ · s ⁻²	N/A · m	T

\magneticflux{ $\langle magnitude \rangle$ }

command	\magneticflux	
base	derived	alternate
kg · m ² · A ⁻¹ · s ⁻²	T · m ²	V · s

\mass{ $\langle magnitude \rangle$ }

command	\mass	
base	derived	alternate
kg	kg	kg

\mobility{ $\langle magnitude \rangle$ }

command	\mobility	
base	derived	alternate
kg · m ² · A ⁻¹ · s ⁻⁴	m ² /V · s	C · m/N · s

\momentofinertia{ $\langle magnitude \rangle$ }

command	\momentofinertia	
base	derived	alternate
kg · m ²	J · s ²	kg · m ²

N 2021-02-24

\momentum $\{\langle magnitude \rangle\}$
\momentumvector $\{\langle c_1, \dots, c_n \rangle\}$
\vectormomentum $\{\langle c_1, \dots, c_n \rangle\}$

command	\momentum	
base	derived	alternate
$\text{kg} \cdot \text{m} \cdot \text{s}^{-1}$	$\text{kg} \cdot \text{m}/\text{s}$	$\text{kg} \cdot \text{m}/\text{s}$

N 2021-02-24

\momentumflux $\{\langle magnitude \rangle\}$
\momentumfluxvector $\{\langle c_1, \dots, c_n \rangle\}$
\vectormomentumflux $\{\langle c_1, \dots, c_n \rangle\}$

command	\momentumflux	
base	derived	alternate
$\text{kg} \cdot \text{m}^{-1} \cdot \text{s}^{-2}$	N/m^2	N/m^2

\numberdensity $\{\langle magnitude \rangle\}$

command	\numberdensity	
base	derived	alternate
m^{-3}	$/\text{m}^3$	$/\text{m}^3$

\permeability $\{\langle magnitude \rangle\}$

command	\permeability	
base	derived	alternate
$\text{kg} \cdot \text{m} \cdot \text{A}^{-2} \cdot \text{s}^{-2}$	H/m	$\text{T} \cdot \text{m}/\text{A}$

\permittivity $\{\langle magnitude \rangle\}$

command	\permittivity	
base	derived	alternate
$\text{A}^2 \cdot \text{s}^4 \cdot \text{kg}^{-1} \cdot \text{m}^{-3}$	F/m	$\text{C}^2/\text{N} \cdot \text{m}^2$

\planeangle $\{\langle magnitude \rangle\}$

command	\planeangle	
base	derived	alternate
$\text{m} \cdot \text{m}^{-1}$	rad	rad

\polarizability $\{\langle magnitude \rangle\}$

command	\polarizability	
base	derived	alternate
$\text{A}^2 \cdot \text{s}^4 \cdot \text{kg}^{-1}$	$\text{C} \cdot \text{m}^2/\text{V}$	$\text{C}^2 \cdot \text{m}/\text{N}$

\power $\{\langle magnitude \rangle\}$

command	\power	
base	derived	alternate
$\text{kg} \cdot \text{m}^2 \cdot \text{s}^{-3}$	W	J/s

\poynting{ $\langle magnitude \rangle$ }

\poyntingvector{ $\langle c_1, \dots, c_n \rangle$ }

\vectorpoynting{ $\langle c_1, \dots, c_n \rangle$ }

command	\poynting	
base	derived	alternate
$\text{kg} \cdot \text{s}^{-3}$	W/m^2	W/m^2

\pressure{ $\langle magnitude \rangle$ }

command	\pressure	
base	derived	alternate
$\text{kg} \cdot \text{m}^{-1} \cdot \text{s}^{-2}$	Pa	N/m^2

\relativepermeability{ $\langle magnitude \rangle$ }

command	\relativepermeability	
base	derived	alternate

\relativepermittivity{ $\langle magnitude \rangle$ }

command	\relativepermittivity	
base	derived	alternate

\resistance{ $\langle magnitude \rangle$ }

command	\resistance	
base	derived	alternate
$\text{kg} \cdot \text{m}^2 \cdot \text{A}^{-2} \cdot \text{s}^{-3}$	Ω	Ω

\resistivity{ $\langle magnitude \rangle$ }

command	\resistivity	
base	derived	alternate
$\text{kg} \cdot \text{m}^3 \cdot \text{A}^{-2} \cdot \text{s}^{-3}$	$\Omega \cdot \text{m}$	$\text{V} \cdot \text{m}/\text{A}$

\solidangle{ $\langle magnitude \rangle$ }

command	\solidangle	
base	derived	alternate
$\text{m}^2 \cdot \text{m}^{-2}$	sr	sr

\specificheatcapacity{ $\langle magnitude \rangle$ }

command	\specificheatcapacity	
base	derived	alternate
$\text{m}^2 \cdot \text{s}^{-2} \cdot \text{K}^{-1}$	$\text{J/K} \cdot \text{kg}$	$\text{J/K} \cdot \text{kg}$

\springstiffness{ $\langle magnitude \rangle$ }

command	\springstiffness	
base	derived	alternate
$\text{kg} \cdot \text{s}^{-2}$	N/m	N/m

\springstretch{ $\langle magnitude \rangle$ }

command	\springstretch	
base	derived	alternate
m	m	m

\stress{ $\langle magnitude \rangle$ }

command	\stress	
base	derived	alternate
$\text{kg} \cdot \text{m}^{-1} \cdot \text{s}^{-2}$	Pa	N/m^2

\strain{ $\langle magnitude \rangle$ }

command	\strain	
base	derived	alternate

\temperature{ $\langle magnitude \rangle$ }

command	\temperature	
base	derived	alternate
K	K	K

\torque{ $\langle magnitude \rangle$ }

\torquevector{ $\langle c_1, \dots, c_n \rangle$ }

\vectortorque{ $\langle c_1, \dots, c_n \rangle$ }

command	\torque	
base	derived	alternate
$\text{kg} \cdot \text{m}^2 \cdot \text{s}^{-2}$	$\text{N} \cdot \text{m}$	$\text{N} \cdot \text{m}$

\velocity{ $\langle magnitude \rangle$ }

\velocityvector{ $\langle c_1, \dots, c_n \rangle$ }

\vectorvelocity{ $\langle c_1, \dots, c_n \rangle$ }

N 2021-02-24

N 2021-02-24

N 2021-02-24

command	\velocity	
base	derived	alternate
$\text{m} \cdot \text{s}^{-1}$	m/s	m/s

\velocityc $\{\langle magnitude \rangle\}$
\velocitycvector $\{\langle c_1, \dots, c_n \rangle\}$
\vectorvelocityc $\{\langle c_1, \dots, c_n \rangle\}$

command	\velocityc	
base	derived	alternate
c	c	c

\volume $\{\langle magnitude \rangle\}$

command	\volume	
base	derived	alternate
m^3	m^3	m^3

\volumechargedensity $\{\langle magnitude \rangle\}$

command	\volumechargedensity	
base	derived	alternate
$\text{A} \cdot \text{s}/\text{m}^{-3}$	C/m^3	C/m^3

\volumemassdensity $\{\langle magnitude \rangle\}$

command	\volumemassdensity	
base	derived	alternate
$\text{kg} \cdot \text{m}^{-3}$	kg/m^3	kg/m^3

\wavelength $\{\langle magnitude \rangle\}$

command	\wavelength	
base	derived	alternate
m	m	m

N 2021-02-24

\wavenumber $\{\langle magnitude \rangle\}$
\wavenumbervector $\{\langle c_1, \dots, c_n \rangle\}$
\vectorwavenumber $\{\langle c_1, \dots, c_n \rangle\}$

command	\wavenumber	
base	derived	alternate
m^{-1}	$/\text{m}$	$/\text{m}$

\work $\{\langle magnitude \rangle\}$

command	\work	
base	derived	alternate
$\text{kg} \cdot \text{m}^2 \cdot \text{s}^{-2}$	J	J

\youngsm modulus{*<magnitude>*}

command	\youngsm modulus	
base	derived	alternate
$\text{kg} \cdot \text{m}^{-1} \cdot \text{s}^{-2}$	Pa	N/m ²

6.4.4 Defining and Redefining Physical Quantities

N 2021-02-16

\NewScalarQuantity{*<name>*}{*<base units>*}[*<derived units>*][*<alternate units>*]

N 2021-02-21

\RenewScalarQuantity{*<name>*}{*<base units>*}[*<derived units>*][*<alternate units>*]

Command to (re)define a new/existing scalar quantity. If the derived or alternate units are omitted, they are defined to be the same as the base units. Do not use both this command and **\NewVectorQuantity** or **\RenewVectorQuantity** to (re)define a quantity.

N 2021-02-16

\NewVectorQuantity{*<name>*}{*<base units>*}[*<derived units>*][*<alternate units>*]

N 2021-02-21

\RenewVectorQuantity{*<name>*}{*<base units>*}[*<derived units>*][*<alternate units>*]

Command to (re)define a new/existing vector quantity. If the derived or alternate units are omitted, they are defined to be the same as the base units. Do not use both this command and **\NewScalarQuantity** or **\RenewScalarQuantity** to (re)define a quantity.

6.4.5 Changing Units

Units are set when **mandi** is loaded, but the default setting can be easily overridden in four ways: command variants that are defined when a **physical quantity**^{P.9} or **physical constant**^{P.23} is defined, a global modal command (switch), a command that sets units for a single instance, and an environment that sets units for its duration. All of these methods work for both physical quantities and physical constants.

U 2021-02-26

\alwaysusebaseunits

U 2021-02-26

\alwaysusederivedunits

U 2021-02-26

\alwaysusealternateunits

Modal commands (switches) for setting the default unit form for the entire document. When **mandi** is loaded, one of these three commands is executed depending on whether the optional **units** key is provided. See the section on loading the package for details. Alternate units are the default because they are the most likely ones to be seen in introductory physics textbooks.

U 2021-02-26

\hereusebaseunits{*<content>*}

U 2021-02-26

\hereusederivedunits{*<content>*}

U 2021-02-26

\hereusealternateunits{*<content>*}

Commands for setting the unit form on the fly for a single instance. The example uses momentum and the Coulomb constant, but they work for any defined quantity and constant.

<code>\(\hereusebaseunits{\momentum{5}} \)</code>	<code>\)</code>	$5 \text{ kg} \cdot \text{m} \cdot \text{s}^{-1}$
<code>\(\hereusederivedunits{\momentum{5}} \)</code>	<code>\)</code>	$5 \text{ kg} \cdot \text{m/s}$
<code>\(\hereusealternateunits{\momentum{5}} \)</code>	<code>\)</code>	$5 \text{ kg} \cdot \text{m/s}$
<code>\(\hereusebaseunits{\oofpez} \)</code>	<code>\)</code>	$9 \times 10^9 \text{ kg} \cdot \text{m}^3 \cdot \text{A}^{-2} \cdot \text{s}^{-4}$
<code>\(\hereusederivedunits{\oofpez} \)</code>	<code>\)</code>	$9 \times 10^9 \text{ m/F}$
<code>\(\hereusealternateunits{\oofpez} \)</code>		$9 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2$

U 2021-02-26

`\begin{usebaseunits}` (use base units)
`\(environment content\)`

U 2021-02-26

`\begin{usederivedunits}` (use derived units)
`\(environment content\)`

U 2021-02-26

`\begin{usealternateunits}` (use alternate units)
`\(environment content\)`
`\end{usealternateunits}`

Inside these environments units are changed for the duration of the environment regardless of the global default setting.

<code>\(\momentum{5} \)</code>	<code>\)</code>	
<code>\(\oofpez \)</code>	<code>\)</code>	
<code>\begin{usebaseunits}</code>		$5 \text{ kg} \cdot \text{m/s}$
<code>\(\momentum{5} \)</code>	<code>\)</code>	$9 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2$
<code>\(\oofpez \)</code>	<code>\)</code>	$5 \text{ kg} \cdot \text{m} \cdot \text{s}^{-1}$
<code>\end{usebaseunits}</code>		$9 \times 10^9 \text{ kg} \cdot \text{m}^3 \cdot \text{A}^{-2} \cdot \text{s}^{-4}$
<code>\begin{usederivedunits}</code>		$5 \text{ kg} \cdot \text{m/s}$
<code>\(\momentum{5} \)</code>	<code>\)</code>	$9 \times 10^9 \text{ m/F}$
<code>\(\oofpez \)</code>	<code>\)</code>	$5 \text{ kg} \cdot \text{m/s}$
<code>\end{usederivedunits}</code>		$9 \times 10^9 \text{ m/F}$
<code>\begin{usealternateunits}</code>		$5 \text{ kg} \cdot \text{m/s}$
<code>\(\momentum{5} \)</code>	<code>\)</code>	$9 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2$
<code>\(\oofpez \)</code>	<code>\)</code>	
<code>\end{usealternateunits}</code>		

6.5 Physical Constants

6.5.1 Typesetting Physical Constants

Take the quantity $\frac{1}{4\pi\epsilon_0}$, sometimes called the **Coulomb constant**, as the prototypical **physical constant** in an introductory physics course. Here are all the ways to access this quantity in **mandi**. As you can see, these commands are almost identical to the corresponding commands for physical quantities.

`\oofpez`

Command for the Coulomb constant. The constant's numerical precision and default units will depend on the options passed to **mandi** at load time. Alternate units and approximate numerical values are the defaults. Other units can be forced as demonstrated.

<code>\(\oofpez \)</code>	<code>\\</code>	$9 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2$
<code>\(\oofpezapproximatevalue \)</code>	<code>\\</code>	9×10^9
<code>\(\oofpezprecisevalue \)</code>	<code>\\</code>	8.9875517923×10^9
<code>\(\oofpezmathsymbol \)</code>	<code>\\</code>	$\frac{1}{4\pi\epsilon_0}$
<code>\(\oofpezbaseunits \)</code>	<code>\\</code>	$9 \times 10^9 \text{ kg} \cdot \text{m}^3 \cdot \text{A}^{-2} \cdot \text{s}^{-4}$
<code>\(\oofpezderivedunits \)</code>	<code>\\</code>	$9 \times 10^9 \text{ m/F}$
<code>\(\oofpezalternateunits \)</code>	<code>\\</code>	$9 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2$
<code>\(\oofpezonlybaseunits \)</code>	<code>\\</code>	$\text{kg} \cdot \text{m}^3 \cdot \text{A}^{-2} \cdot \text{s}^{-4}$
<code>\(\oofpezonlyderivedunits \)</code>	<code>\\</code>	m/F
<code>\(\oofpezonlyalternateunits \)</code>	<code>\\</code>	$\text{N} \cdot \text{m}^2/\text{C}^2$

6.5.2 Checking Physical Constants

U 2022-01-27

`\CheckConstant{ $\langle name \rangle$ }`

Command to check and typeset the constant's name, mathematical symbol, approximate value, precise value, base units, derived units, and alternate units.

6.5.3 Predefined Physical Constants

Every other defined physical constant can be treated similarly to `\oofpez`^{P.27}. Just replace `oofpez` with the constant's name. Unfortunately, there is no universal agreement on the names of every constant so don't fret if the names used here vary from other sources. Here are all the physical constants, with all their units, defined in `mandi`. The constants `\coulombconstant`^{P.25} and `\biotsavartconstant` are defined as semantic aliases for, respectively, `\oofpez`^{P.27} and `\mzofp`^{P.27}.

`\avogadro` (exact)

command	<code>\avogadro</code>	
symbol	approximate	precise
N _A	6×10^{23}	$6.02214076 \times 10^{23}$
base	derived	alternate
mol ⁻¹	/mol	/mol

N 2021-02-02

\biotsavartconstant

command	<code>\biotsavartconstant</code>	
symbol	approximate	precise
$\frac{\mu_0}{4\pi}$	10^{-7}	10^{-7}
base	derived	alternate
$\text{kg} \cdot \text{m} \cdot \text{A}^{-2} \cdot \text{s}^{-2}$	H/m	T · m/A

\bohrradius

command	<code>\bohrradius</code>	
symbol	approximate	precise
a_o	5.3×10^{-11}	$5.29177210903 \times 10^{-11}$
base	derived	alternate
m	m	m

`\boltzmann` (exact)

command	<code>\boltzmann</code>	
symbol	approximate	precise
k_B	1.4×10^{-23}	1.380649×10^{-23}
base	derived	alternate
$\text{kg} \cdot \text{m}^2 \cdot \text{s}^{-2} \cdot \text{K}^{-1}$	J/K	J/K

N 2021-02-02

`\coulombconstant`

command	<code>\coulombconstant</code>	
symbol	approximate	precise
$\frac{1}{4\pi\epsilon_o}$	9×10^9	8.9875517923×10^9
base	derived	alternate
$\text{kg} \cdot \text{m}^3 \cdot \text{A}^{-2} \cdot \text{s}^{-4}$	m/F	$\text{N} \cdot \text{m}^2/\text{C}^2$

`\earthmass`

command	<code>\earthmass</code>	
symbol	approximate	precise
M_{Earth}	6.0×10^{24}	5.9722×10^{24}
base	derived	alternate
kg	kg	kg

`\earthmoondistance`

command	<code>\earthmoondistance</code>	
symbol	approximate	precise
d_{EM}	3.8×10^8	3.81550×10^8
base	derived	alternate
m	m	m

`\earthradius`

command	<code>\earthradius</code>	
symbol	approximate	precise
R_{Earth}	6.4×10^6	6.3781×10^6
base	derived	alternate
m	m	m

`\earthsundistance`

command	<code>\earthsundistance</code>	
symbol	approximate	precise
d_{ES}	1.5×10^{11}	1.496×10^{11}
base	derived	alternate
m	m	m

`\electroncharge`

command	<code>\electroncharge</code>	
symbol	approximate	precise
q_e	-1.6×10^{-19}	$-1.602176634 \times 10^{-19}$
base	derived	alternate
A · s	C	C

`\electronCharge`

command	<code>\electronCharge</code>	
symbol	approximate	precise
Q_e	-1.6×10^{-19}	$-1.602176634 \times 10^{-19}$
base	derived	alternate
A · s	C	C

`\electronmass`

command	<code>\electronmass</code>	
symbol	approximate	precise
m_e	9.1×10^{-31}	$9.1093837015 \times 10^{-31}$
base	derived	alternate
kg	kg	kg

`\elementarycharge` (exact)

command	<code>\elementarycharge</code>	
symbol	approximate	precise
e	1.6×10^{-19}	$1.602176634 \times 10^{-19}$
base	derived	alternate
A · s	C	C

`\finestructure`

command	<code>\finestructure</code>	
symbol	approximate	precise
α	$\frac{1}{137}$	$7.2973525693 \times 10^{-3}$
base	derived	alternate

`\hydrogenmass`

command	<code>\hydrogenmass</code>	
symbol	approximate	precise
m_{H}	1.7×10^{-27}	$1.6737236 \times 10^{-27}$
base	derived	alternate
kg	kg	kg

`\moonearthdistance`

command	<code>\moonearthdistance</code>	
symbol	approximate	precise
d_{ME}	3.8×10^8	3.81550×10^8
base	derived	alternate
m	m	m

`\moonmass`

command	<code>\moonmass</code>	
symbol	approximate	precise
M_{Moon}	7.3×10^{22}	7.342×10^{22}
base	derived	alternate
kg	kg	kg

`\moonradius`

command	<code>\moonradius</code>	
symbol	approximate	precise
R_{Moon}	1.7×10^6	1.7371×10^6
base	derived	alternate
m	m	m

`\mzofp`

command	<code>\mzofp</code>	
symbol	approximate	precise
$\frac{\mu_0}{4\pi}$	10^{-7}	10^{-7}
base	derived	alternate
$\text{kg} \cdot \text{m} \cdot \text{A}^{-2} \cdot \text{s}^{-2}$	H/m	T · m/A

`\neutronmass`

command	<code>\neutronmass</code>	
symbol	approximate	precise
m_{n}	1.7×10^{-27}	$1.67492749804 \times 10^{-27}$
base	derived	alternate
kg	kg	kg

`\oofpez`

command	\oofpez	precise
symbol	approximate	
$\frac{1}{4\pi\epsilon_0}$	9×10^9	8.9875517923×10^9
base	derived	alternate
$\text{kg} \cdot \text{m}^3 \cdot \text{A}^{-2} \cdot \text{s}^{-4}$	m/F	$\text{N} \cdot \text{m}^2/\text{C}^2$

\oofpezcs

command	\oofpezcs	precise
symbol	approximate	
$\frac{1}{4\pi\epsilon_0 c^2}$	10^{-7}	10^{-7}
base	derived	alternate
$\text{kg} \cdot \text{m} \cdot \text{A}^{-2} \cdot \text{s}^{-2}$	$\text{T} \cdot \text{m}^2$	$\text{N} \cdot \text{s}^2/\text{C}^2$

\planck (exact)

command	\planck	precise
symbol	approximate	
h	6.6×10^{-34}	$6.62607015 \times 10^{-34}$
base	derived	alternate
$\text{kg} \cdot \text{m}^2 \cdot \text{s}^{-1}$	$\text{J} \cdot \text{s}$	$\text{J} \cdot \text{s}$

\planckbar

command	\planckbar	precise
symbol	approximate	
\hbar	1.1×10^{-34}	$1.054571817 \times 10^{-34}$
base	derived	alternate
$\text{kg} \cdot \text{m}^2 \cdot \text{s}^{-1}$	$\text{J} \cdot \text{s}$	$\text{J} \cdot \text{s}$

\planckc

command	\planckc	precise
symbol	approximate	
hc	2.0×10^{-25}	$1.98644586 \times 10^{-25}$
base	derived	alternate
$\text{kg} \cdot \text{m}^3 \cdot \text{s}^{-2}$	$\text{J} \cdot \text{m}$	$\text{J} \cdot \text{m}$

\protoncharge

command	\protoncharge	precise
symbol	approximate	
q_p	$+1.6 \times 10^{-19}$	$+1.602176634 \times 10^{-19}$
base	derived	alternate
$\text{A} \cdot \text{s}$	C	C

\protonCharge

command	<code>\protonCharge</code>	
symbol	approximate	precise
Q_p	$+1.6 \times 10^{-19}$	$+1.602176634 \times 10^{-19}$
base	derived	alternate
A · s	C	C

`\protonmass`

command	<code>\protonmass</code>	
symbol	approximate	precise
m_p	1.7×10^{-27}	$1.672621898 \times 10^{-27}$
base	derived	alternate
kg	kg	kg

`\rydberg`

command	<code>\rydberg</code>	
symbol	approximate	precise
R_∞	1.1×10^7	$1.0973731568160 \times 10^7$
base	derived	alternate
m^{-1}	m^{-1}	m^{-1}

`\speedoflight` (exact)

command	<code>\speedoflight</code>	
symbol	approximate	precise
c	3×10^8	2.99792458×10^8
base	derived	alternate
$m \cdot s^{-1}$	m/s	m/s

`\stefanboltzmann`

command	<code>\stefanboltzmann</code>	
symbol	approximate	precise
σ	5.7×10^{-8}	5.670374×10^{-8}
base	derived	alternate
$kg \cdot s^{-3} \cdot K^{-4}$	$W/m^2 \cdot K^4$	$W/m^2 \cdot K^4$

`\sunearthdistance`

command	<code>\sunearthdistance</code>	
symbol	approximate	precise
d_{SE}	1.5×10^{11}	1.496×10^{11}
base	derived	alternate
m	m	m

`\sunradius`

command	<code>\sunradius</code>	
symbol	approximate	precise
R_{Sun}	7.0×10^8	6.957×10^8
base	derived	alternate
m	m	m

`\surfacegravfield`

command	<code>\surfacegravfield</code>	
symbol	approximate	precise
g	9.8	9.807
base	derived	alternate
$\text{m} \cdot \text{s}^{-2}$	N/kg	N/kg

`\universalgrav`

command	<code>\universalgrav</code>	
symbol	approximate	precise
G	6.7×10^{-11}	6.67430×10^{-11}
base	derived	alternate
$\text{m}^3 \cdot \text{kg}^{-1} \cdot \text{s}^{-2}$	$\text{N} \cdot \text{m}^2/\text{kg}^2$	$\text{N} \cdot \text{m}^2/\text{kg}^2$

`\vacuumpermeability`

command	<code>\vacuumpermeability</code>	
symbol	approximate	precise
μ_0	$4\pi \times 10^{-7}$	$4\pi \times 10^{-7}$
base	derived	alternate
$\text{kg} \cdot \text{m} \cdot \text{A}^{-2} \cdot \text{s}^{-2}$	H/m	$\text{T} \cdot \text{m}/\text{A}$

`\vacuumpermittivity`

command	<code>\vacuumpermittivity</code>	
symbol	approximate	precise
ϵ_0	9×10^{-12}	$8.854187817 \times 10^{-12}$
base	derived	alternate
$\text{A}^2 \cdot \text{s}^4 \cdot \text{kg}^{-1} \cdot \text{m}^{-3}$	F/m	$\text{C}^2/\text{N} \cdot \text{m}^2$

6.5.4 Defining and Redefining Physical Constants

N 2021-02-16

`\NewPhysicalConstant` $\{\langle name \rangle\}\{\langle symbol \rangle\}\{\langle approximate value \rangle\}\{\langle precise value \rangle\}\{\langle base units \rangle\}$
 $[\langle derived units \rangle] [\langle alternate units \rangle]$

N 2021-02-21

`\RenewPhysicalConstant` $\{\langle name \rangle\}\{\langle symbol \rangle\}\{\langle approximate value \rangle\}\{\langle precise value \rangle\}\{\langle base units \rangle\}$
 $[\langle derived units \rangle] [\langle alternate units \rangle]$

Command to define/redefine a new/existing physical constant. If the derived or alternate units are omitted, they are defined to be the same as the base units.

6.5.5 Changing Precision

[Changing units](#)^{→ P.22} works for physical constants just as it does for physical quantities. A similar mechanism is provided for changing the precision of physical constants' numerical values.

N 2021-02-16
N 2021-02-16

`\alwaysuseapproximateconstants`
`\alwaysusepreciseconstants`

Modal commands (switches) for setting the default precision for the entire document. The default when the package is loaded is set by the presence or absence of the [preciseconstants](#)^{→P.8} key.

N 2021-02-16
N 2021-02-16

`\hereuseapproximateconstants{<content>}`
`\hereusepreciseconstants{<content>}`

Commands for setting the precision on the fly for a single instance.

<code>\(\hereuseapproximateconstants{\oofpez} \) \\\</code>	$9 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2$
<code>\(\hereusepreciseconstants{\oofpez} \) \\\</code>	$8.9875517923 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2$

N 2021-02-16

`\begin{useapproximateconstants}` (use approximate constants)
<environment content>

`\end{useapproximateconstants}`

N 2021-02-16

`\begin{usepreciseconstants}` (use precise constants)
<environment content>

`\end{usepreciseconstants}`

Inside these environments precision is changed for the duration of the environment regardless of the global default setting.

<code>\(\oofpez \) \\\</code>	
<code>\begin{useapproximateconstants}</code>	
<code>\(\oofpez \) \\\</code>	$9 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2$
<code>\end{useapproximateconstants}</code>	$9 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2$
<code>\begin{usepreciseconstants}</code>	$8.9875517923 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2$
<code>\(\oofpez \) \\\</code>	$9 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2$
<code>\end{usepreciseconstants}</code>	
<code>\(\oofpez \) \\\</code>	

6.6 Predefined Units and Constructs

These commands should be used only in defining or redefining physical quantities or physical constants. One exception is [\emptyunit](#), which may be used for explanatory purposes.

U 2023-08-01

`\per`
`\usk`
`\units{<magnitude>}{<unit>}`
`\emptyunit`
`\ampere`
`\atomicmassunit`
`\candela`

	<code>\coulomb</code>	
	<code>\degree</code>	
	<code>\electronvolt</code>	(not SI but common in introductory physics)
<code>N 2021-04-15</code>	<code>\ev</code>	(alias)
	<code>\farad</code>	
	<code>\henry</code>	
	<code>\hertz</code>	
	<code>\joule</code>	
	<code>\kelvin</code>	
<code>N 2021-04-15</code>	<code>\kev</code>	(alias)
<code>N 2021-04-15</code>	<code>\kiloelectronvolt</code>	(not SI but common in introductory physics)
	<code>\kilogram</code>	
	<code>\lightspeed</code>	(not SI but common relativity)
<code>N 2021-04-15</code>	<code>\megaelectronvolt</code>	(not SI but common in introductory physics)
	<code>\meter</code>	
	<code>\metre</code>	(alias)
<code>N 2021-04-15</code>	<code>\mev</code>	(alias)
	<code>\mole</code>	
	<code>\newton</code>	
	<code>\ohm</code>	
	<code>\pascal</code>	
	<code>\radian</code>	
	<code>\second</code>	
	<code>\siemens</code>	
	<code>\steradian</code>	
	<code>\tesla</code>	
	<code>\volt</code>	
	<code>\watt</code>	
	<code>\weber</code>	
	<code>\tothetwo</code>	(postfix)
	<code>\tothethree</code>	(postfix)
	<code>\tothefour</code>	(postfix)
	<code>\inverse</code>	(postfix)
	<code>\totheinversetwo</code>	(postfix)
	<code>\totheinversethree</code>	(postfix)
	<code>\totheinversefour</code>	(postfix)

	/
	.
	3 m/s
	□
<code>\(\per \)</code>	A
<code>\(\usk \)</code>	u
<code>\(\units{3}{\meter\per\second} \)</code>	cd
<code>\(\emptyunit \)</code>	C
<code>\(\ampere \)</code>	°
<code>\(\atomicmassunit \)</code>	eV
<code>\(\candela \)</code>	F
<code>\(\coulomb \)</code>	H
<code>\(\degree \)</code>	Hz
<code>\(\electronvolt \)</code>	J
<code>\(\farad \)</code>	K
<code>\(\henry \)</code>	keV
<code>\(\hertz \)</code>	kg
<code>\(\joule \)</code>	c
<code>\(\kelvin \)</code>	m
<code>\(\kev \)</code>	m
<code>\(\kilogram \)</code>	MeV
<code>\(\lightspeed \)</code>	mol
<code>\(\meter \)</code>	N
<code>\(\metre \)</code>	Ω
<code>\(\mev \)</code>	Pa
<code>\(\mole \)</code>	rad
<code>\(\newton \)</code>	s
<code>\(\ohm \)</code>	S
<code>\(\pascal \)</code>	sr
<code>\(\radian \)</code>	T
<code>\(\second \)</code>	V
<code>\(\siemens \)</code>	W
<code>\(\steradian \)</code>	Wb
<code>\(\tesla \)</code>	□ ²
<code>\(\emptyunit\tothetwo \)</code>	□ ³
<code>\(\emptyunit\tothethree \)</code>	□ ⁴
<code>\(\emptyunit\tothefour \)</code>	□ ⁻¹
<code>\(\emptyunit\inverse \)</code>	□ ⁻²
<code>\(\emptyunit\totheinversetwo \)</code>	□ ⁻³
<code>\(\emptyunit\totheinversethree \)</code>	□ ⁻⁴
<code>\(\emptyunit\totheinversefour \)</code>	

`\(\hbar \)`

\hbar

`\tento{<number>}`

`\timestento{<number>}`

`\xtento{<number>}`

Commands for powers of ten and scientific notation.

`\(\tento{-4} \)` `\`

10^{-4}

`\(3\timestento{8} \)` `\`

3×10^8

`\(3\xtento{8} \)`

3×10^8

U 2022-01-27

`\mivector[<delimiter>]{<c1, ..., cn>}[<units>]`

Typesets a vector as either numeric or symbolic components with an optional unit (for numerical components only). There must be more than one component, and there can be more than three components. The delimiter used in the list of components can be specified; the default is a comma. The notation mirrors that of *Matter & Interactions*.

`\(\mivector{p_0,p_1,p_2,p_3} \)`

\rangle

`\(\mivector{\gamma mc,\gamma mv_x,\gamma mv_y,\gamma mv_z} \)`

$\langle p_0, p_1, p_2, p_3 \rangle$
 $\langle \gamma mc, \gamma mv_x, \gamma mv_y, \gamma mv_z \rangle$

`\(\mivector{\frac{Q_1Q_2}{x^2},0,0} \)`

\rangle

$\langle \frac{Q_1Q_2}{x^2}, 0, 0 \rangle$

`\(\mivector{-1,0,0} \)`

\rangle

$\langle -1, 0, 0 \rangle$

`\(\mivector{-1,0,0}[\text{velocityonlyderivedunits}] \)`

\rangle

$\langle -1, 0, 0 \rangle \text{ m/s}$

`\(\mivector{-1,0,0}[\text{meter\per\second}] \)`

\rangle

$\langle -1, 0, 0 \rangle \text{ m/s}$

`\(\velocity{\mivector{-1,0,0}} \)`

$\langle -1, 0, 0 \rangle \text{ m/s}$

6.7 mandi Source Code

Define the package version and date for global use, exploiting the fact that in a .sty file there is now no need for `\makeatletter` and `\makeatother`. This simplifies defining internal commands (with @ in the name) that are not for the user to know about.

```
1 \def\mandi@version{3.2.2a}
2 \def\mandi@date{2024-09-12}
3 \NeedsTeXFormat{LaTeX2e}[2020-02-02]
4 \DeclareRelease{v3.2.2a}{2024-09-12}{mandi.sty}
5 \DeclareCurrentRelease{v\mandi@version}{\mandi@date}
6 \ProvidesPackage{mandi}
7 [\mandi@date\space v\mandi@version\space Macros for physical quantities]
```

Define a convenient package version command.

```
8 \newcommand*{\mandiversion}{v\mandi@version\space dated \mandi@date}
```

Load third party packages, documenting why each one is needed.

We need `pgfopts` for a key-value interface.

```
9 \RequirePackage{pgfopts}
```

We need `array` for `\chkquantity` and `\chkconstant`.

```
10 \RequirePackage{array}
```

We need `iftex` so we can require `LuaATEX`.

```
11 \RequirePackage{iftex}
```

We need `mathtools` for intelligent delimiters.

```
12 \RequirePackage{mathtools}
```

We need `unicode-math` for Unicode support and changing fonts.

```
13 \RequirePackage{unicode-math}
```

Suppress some annoying warnings.

```
14 \unimathsetup{warnings-off={mathtools-colon,mathtools-overbracket}}
```

Load `xparse` if necessary.

```
15 \IfFormatAtLeastTF{2020-10-01}
```

```
16 {}%
```

```
17 {\RequirePackage{xparse}}%
```

We require the `LuaATEX` engine.

```
18 \RequireLuaTeX
```

Parts of the unit engine have been rewritten with `xparse` for both clarity and power. Note that `xparse` is now part of the `LATEX 2ε` kernel. Other parts have been rewritten in `expl` with a look to the future.

Define some generic internal selectors.

```
19 \newcommand*{\mandi@selectunits}{}
```

```
20 \newcommand*{\mandi@selectprecision}{}
```

Define some specific internal selectors. The first two are really just workalikes for `\@firstoftwo` and `\@secondoftwo`. The third, fourth, and fifth are `\@firstofthree`, `\@secondofthree`, and `\@thirdofthree` and apparently do not yet exist.

```
21 \newcommand*{\mandi@selectapproximate}[2]{#1}
```

```
22 \newcommand*{\mandi@selectprecise}[2]{#2}
```

```
23 \newcommand*{\mandi@selectbaseunits}[3]{#1}
```

```
24 \newcommand*{\mandi@selectderivedunits}[3]{#2}
```

```
25 \newcommand*{\mandi@selectalternateunits}[3]{#3}
```

Document level global switches.

```

26 \NewDocumentCommand{\alwaysusebaseunits}{-}{
27   {\renewcommand*{\mandi@selectunits}{\mandi@selectbaseunits}}}%
28 \NewDocumentCommand{\alwaysusederivedunits}{-}{
29   {\renewcommand*{\mandi@selectunits}{\mandi@selectderivedunits}}}%
30 \NewDocumentCommand{\alwaysusealternateunits}{-}{
31   {\renewcommand*{\mandi@selectunits}{\mandi@selectalternateunits}}}%
32 \NewDocumentCommand{\alwaysuseapproximateconstants}{-}{
33   {\renewcommand*{\mandi@selectprecision}{\mandi@selectapproximate}}}%
34 \NewDocumentCommand{\alwaysusepreciseconstants}{-}{
35   {\renewcommand*{\mandi@selectprecision}{\mandi@selectprecise}}}%

```

Document level localized variants.

```

36 \NewDocumentCommand{\hereusebaseunits}{ m }{\begingroup\alwaysusebaseunits#1\endgroup}%
37 \NewDocumentCommand{\hereusederivedunits}{ m }{\begingroup\alwaysusederivedunits#1\endgroup}%
38 \NewDocumentCommand{\hereusealternateunits}{ m }{\begingroup\alwaysusealternateunits#1\endgroup}%
39 \NewDocumentCommand{\hereuseapproximateconstants}{ m }{\begingroup\alwaysuseapproximateconstants#1\endgroup}%
40 \NewDocumentCommand{\hereusepreciseconstants}{ m }{\begingroup\alwaysusepreciseconstants#1\endgroup}%

```

Document level environments.

```

41 \NewDocumentEnvironment{usebaseunits}{-}{\alwaysusebaseunits}{-}%
42 \NewDocumentEnvironment{usederivedunits}{-}{\alwaysusederivedunits}{-}%
43 \NewDocumentEnvironment{usealternateunits}{-}{\alwaysusealternateunits}{-}%
44 \NewDocumentEnvironment{useapproximateconstants}{-}{\alwaysuseapproximateconstants}{-}%
45 \NewDocumentEnvironment{usepreciseconstants}{-}{\alwaysusepreciseconstants}{-}%

```

mandi now has a key-value interface, implemented with [pgfopts](#) and [pgfkeys](#). There are two options: [units](#)^{P.8}, with values **base**, **derived**, or **alternate** selects the default form of units [preciseconstants](#)^{P.8}, with values **true** and **false**, selects precise numerical values for constants rather than approximate values.

First, define the keys. The key handlers require certain commands defined by the unit engine.

```

46 \newif\ifusingpreciseconstants
47 \pgfkeys{
48   /mandi/options/.cd,
49   initial@setup/.style={%
50     /mandi/options/buffered@units/.initial=alternate,%
51   },%
52   initial@setup,%
53   preciseconstants/.is if=usingpreciseconstants,%
54   units/.is choice,%
55   units/.default=derived,%
56   units/alternate/.style={/mandi/options/buffered@units=alternate},%
57   units/base/.style={/mandi/options/buffered@units=base},%
58   units/derived/.style={/mandi/options/buffered@units=derived},%
59   .unknown/.code={%
60     \typeout{}%
61     \typeout{mandi: You used unknown option '\pgfkeyscurrentname'.}%
62   },%
63 }%

```

Process the options.

```

64 \ProcessPgfPackageOptions{/mandi/options}

```

Write a banner to the console showing the options in use.

```

65 \typeout{}%
66 \typeout{mandi: You are using mandi \mandiversion.}%
67 \typeout{mandi: This package requires LuaLaTeX.}%
68 \typeout{mandi: Loadtime options...}

```

Complete the banner by showing currently selected options. The value of the `units`^{P.8} key is used in situ to set the default units.

```

69 \newcommand*{\mandi@do@setup}{%
70   \csname alwaysuse\pgfkeysvalueof{/mandi/options/buffered@units}units\endcsname%
71   \typeout{mandi: You will get \pgfkeysvalueof{/mandi/options/buffered@units}\space units.}%
72   \ifusingpreciseconstants
73     \alwaysusepreciseconstants
74     \typeout{mandi: You will get precise constants.}%
75   \else
76     \alwaysuseapproximateconstants
77     \typeout{mandi: You will get approximate constants.}%
78   \fi
79   \typeout{}}%
80 }%
81 \mandi@do@setup

```

Define a setup command that overrides the loadtime options when called with new options. A new banner is written to the console.

```

82 \NewDocumentCommand{\mandisetup}{ m }%
83 {%
84   \IfValueT{#1}%
85     {%
86       \pgfqkeys{/mandi/options}{#1}
87       \typeout{}%
88       \typeout{mandi: mandisetup options...}
89       \mandi@do@setup
90     }%
91   }%

```

Define units and related constructs to be used with the unit engine. All single letter macros are now gone. We basically absorbed and adapted the now outdated `Slunits` package. We make use of `\symup{...}` from the `unicode-math` package.

```

92 \NewDocumentCommand{\per}{ }{ }{ }
93 \NewDocumentCommand{\usk}{ }{ }{ }
94 \NewDocumentCommand{\units}{ m m }%
95 {%
96   \IfValueTF{#2}
97     {%
98       {#1}{\,#2}
99     }%
100   {%
101     {#1}{ }
102   }%
103 }%
104 \NewDocumentCommand{\ampere}{ }{ }\symup{A}
105 \NewDocumentCommand{\atomicmassunit}{ }{ }\symup{u}
106 \NewDocumentCommand{\candela}{ }{ }\symup{cd}
107 \NewDocumentCommand{\coulomb}{ }{ }\symup{C}
108 \NewDocumentCommand{\degree}{ }{ }{ }
109 \NewDocumentCommand{\electronvolt}{ }{ }\symup{eV}
110 \NewDocumentCommand{\ev}{ }{ }\electronvolt
111 \NewDocumentCommand{\farad}{ }{ }\symup{F}
112 \NewDocumentCommand{\henry}{ }{ }\symup{H}
113 \NewDocumentCommand{\hertz}{ }{ }\symup{Hz}
114 \NewDocumentCommand{\joule}{ }{ }\symup{J}
115 \NewDocumentCommand{\kelvin}{ }{ }\symup{K}
116 \NewDocumentCommand{\kev}{ }{ }\kilolectronvolt
117 \NewDocumentCommand{\kilolectronvolt}{ }{ }\symup{keV}
118 \NewDocumentCommand{\kilogram}{ }{ }\symup{kg}

```

```

119 \NewDocumentCommand{\lightspeed}{-}{\!\symup{c}}
120 \NewDocumentCommand{\megaelectronvolt}{-}{\symup{MeV}}
121 \NewDocumentCommand{\meter}{-}{\symup{m}}
122 \NewDocumentCommand{\metre}{-}{\meter}
123 \NewDocumentCommand{\mev}{-}{\megaelectronvolt}
124 \NewDocumentCommand{\mole}{-}{\symup{mol}}
125 \NewDocumentCommand{\newton}{-}{\symup{N}}
126 \NewDocumentCommand{\ohm}{-}{\symup{\Omega}}
127 \NewDocumentCommand{\pascal}{-}{\symup{Pa}}
128 \NewDocumentCommand{\radian}{-}{\symup{rad}}
129 \NewDocumentCommand{\second}{-}{\symup{s}}
130 \NewDocumentCommand{\siemens}{-}{\symup{S}}
131 \NewDocumentCommand{\steradian}{-}{\symup{sr}}
132 \NewDocumentCommand{\tesla}{-}{\symup{T}}
133 \NewDocumentCommand{\volt}{-}{\symup{V}}
134 \NewDocumentCommand{\watt}{-}{\symup{W}}
135 \NewDocumentCommand{\weber}{-}{\symup{Wb}}
136 \NewDocumentCommand{\tothetwo}{-}{\sim{2}}
137 \NewDocumentCommand{\tothethree}{-}{\sim{3}}
138 \NewDocumentCommand{\tothefour}{-}{\sim{4}}
139 \NewDocumentCommand{\inverse}{-}{\sim{-1}}
140 \NewDocumentCommand{\totheinversetwo}{-}{\sim{-2}}
141 \NewDocumentCommand{\totheinversethree}{-}{\sim{-3}}
142 \NewDocumentCommand{\totheinversefour}{-}{\sim{-4}}
143 \NewDocumentCommand{\emptyunit}{-}{\mdlgwhtsquare}
144 \NewDocumentCommand{\tento}{ m }{10^{\#1}}
145 \NewDocumentCommand{\timestento}{ m }{\times\tento{\#1}}
146 \NewDocumentCommand{\xtento}{ m }{\times\tento{\#1}}

147 \ExplSyntaxOn
148 \cs_new:Npn \__mandi_newscalarquantity:nnnn #1#2#3#4
149 {
150   \cs_new:cpn {#1} ##1 {\units{##1}{\mandi@selectunits{#2}{#3}{#4}}}
151   \cs_new:cpn {#1value} ##1 {##1}
152   \cs_new:cpn {#1baseunits} ##1 {\units{##1}{\mandi@selectbaseunits{#2}{#3}{#4}}}
153   \cs_new:cpn {#1derivedunits} ##1 {\units{##1}{\mandi@selectderivedunits{#2}{#3}{#4}}}
154   \cs_new:cpn {#1alternateunits} ##1 {\units{##1}{\mandi@selectalternateunits{#2}{#3}{#4}}}
155   \cs_new:cpn {#1onlybaseunits} {\mandi@selectbaseunits{#2}{#3}{#4}}
156   \cs_new:cpn {#1onlyderivedunits} {\mandi@selectderivedunits{#2}{#3}{#4}}
157   \cs_new:cpn {#1onlyalternateunits} {\mandi@selectalternateunits{#2}{#3}{#4}}
158 }
159 \NewDocumentCommand{\NewScalarQuantity}{ m m O{#2} O{#2} }
160 {
161   \__mandi_newscalarquantity:nnnn { #1 }{ #2 }{ #3 }{ #4 }
162 }

```

Redefining an existing scalar quantity.

```

163 \cs_new:Npn \__mandi_renewscalarquantity:nnnn #1#2#3#4
164 {
165   \cs_set:cpn {#1} ##1 {\units{##1}{\mandi@selectunits{#2}{#3}{#4}}}
166   \cs_set:cpn {#1value} ##1 {##1}
167   \cs_set:cpn {#1baseunits} ##1 {\units{##1}{\mandi@selectbaseunits{#2}{#3}{#4}}}
168   \cs_set:cpn {#1derivedunits} ##1 {\units{##1}{\mandi@selectderivedunits{#2}{#3}{#4}}}
169   \cs_set:cpn {#1alternateunits} ##1 {\units{##1}{\mandi@selectalternateunits{#2}{#3}{#4}}}
170   \cs_set:cpn {#1onlybaseunits} {\mandi@selectbaseunits{#2}{#3}{#4}}
171   \cs_set:cpn {#1onlyderivedunits} {\mandi@selectderivedunits{#2}{#3}{#4}}
172   \cs_set:cpn {#1onlyalternateunits} {\mandi@selectalternateunits{#2}{#3}{#4}}
173 }
174 \NewDocumentCommand{\RenewScalarQuantity}{ m m O{#2} O{#2} }
175 {

```

```

176 \__mandi_renewscalarquantity:nnnn { #1 }{ #2 }{ #3 }{ #4 }
177 }

```

Defining a new vector quantity. Note that a corresponding scalar is also defined.

```

178 \cs_new:Npn \__mandi_newvectorquantity:nnnn #1#2#3#4
179 {
180   \__mandi_renewscalarquantity:nnnn { #1 }{ #2 }{ #3 }{ #4 }
181   \cs_new:cpn {vector#1} ##1 {\units{\mivector{##1}}{\mandi@selectunits{#2}{#3}{#4}}}
182   \cs_new:cpn {#1vector} ##1 {\units{\mivector{##1}}{\mandi@selectunits{#2}{#3}{#4}}}
183   \cs_new:cpn {vector#1value} ##1 {\mivector{##1}}
184   \cs_new:cpn {#1vectorvalue} ##1 {\mivector{##1}}
185   \cs_new:cpn {vector#1baseunits} ##1 {\units{\mivector{##1}}{\mandi@selectbaseunits{#2}{#3}{#4}}}
186   \cs_new:cpn {#1vectorbaseunits} ##1 {\units{\mivector{##1}}{\mandi@selectbaseunits{#2}{#3}{#4}}}
187   \cs_new:cpn {vector#1derivedunits} ##1 {\units{\mivector{##1}}{\mandi@selectderivedunits{#2}{#3}{#4}}}
188   \cs_new:cpn {#1vectorderivedunits} ##1 {\units{\mivector{##1}}{\mandi@selectderivedunits{#2}{#3}{#4}}}
189   \cs_new:cpn {vector#1alternateunits} ##1 {\units{\mivector{##1}}{\mandi@selectalternateunits{#2}{#3}{#4}}}
190   \cs_new:cpn {#1vectoralternateunits} ##1 {\units{\mivector{##1}}{\mandi@selectalternateunits{#2}{#3}{#4}}}
191   \cs_new:cpn {vector#1onlybaseunits} {\mandi@selectbaseunits{#2}{#3}{#4}}
192   \cs_new:cpn {#1vectoronlybaseunits} {\mandi@selectbaseunits{#2}{#3}{#4}}
193   \cs_new:cpn {vector#1onlyderivedunits} {\mandi@selectderivedunits{#2}{#3}{#4}}
194   \cs_new:cpn {#1vectoronlyderivedunits} {\mandi@selectderivedunits{#2}{#3}{#4}}
195   \cs_new:cpn {vector#1onlyalternateunits} {\mandi@selectalternateunits{#2}{#3}{#4}}
196   \cs_new:cpn {#1vectoronlyalternateunits} {\mandi@selectalternateunits{#2}{#3}{#4}}
197 }
198 \NewDocumentCommand{\NewVectorQuantity}{ m m O{#2} O{#2} }
199 {
200   \__mandi_newvectorquantity:nnnn { #1 }{ #2 }{ #3 }{ #4 }
201 }

```

Redefining an existing vector quantity. Note that a corresponding scalar is also redefined.

```

202 \cs_new:Npn \__mandi_renewvectorquantity:nnnn #1#2#3#4
203 {
204   \__mandi_renewscalarquantity:nnnn { #1 }{ #2 }{ #3 }{ #4 }
205   \cs_set:cpn {vector#1} ##1 {\units{\mivector{##1}}{\mandi@selectunits{#2}{#3}{#4}}}
206   \cs_set:cpn {#1vector} ##1 {\units{\mivector{##1}}{\mandi@selectunits{#2}{#3}{#4}}}
207   \cs_set:cpn {vector#1value} ##1 {\mivector{##1}}
208   \cs_set:cpn {#1vectorvalue} ##1 {\mivector{##1}}
209   \cs_set:cpn {vector#1baseunits} ##1 {\units{\mivector{##1}}{\mandi@selectbaseunits{#2}{#3}{#4}}}
210   \cs_set:cpn {#1vectorbaseunits} ##1 {\units{\mivector{##1}}{\mandi@selectbaseunits{#2}{#3}{#4}}}
211   \cs_set:cpn {vector#1derivedunits} ##1 {\units{\mivector{##1}}{\mandi@selectderivedunits{#2}{#3}{#4}}}
212   \cs_set:cpn {#1vectorderivedunits} ##1 {\units{\mivector{##1}}{\mandi@selectderivedunits{#2}{#3}{#4}}}
213   \cs_set:cpn {vector#1alternateunits} ##1 {\units{\mivector{##1}}{\mandi@selectalternateunits{#2}{#3}{#4}}}
214   \cs_set:cpn {#1vectoralternateunits} ##1 {\units{\mivector{##1}}{\mandi@selectalternateunits{#2}{#3}{#4}}}
215   \cs_set:cpn {vector#1onlybaseunits} {\mandi@selectbaseunits{#2}{#3}{#4}}
216   \cs_set:cpn {#1vectoronlybaseunits} {\mandi@selectbaseunits{#2}{#3}{#4}}
217   \cs_set:cpn {vector#1onlyderivedunits} {\mandi@selectderivedunits{#2}{#3}{#4}}
218   \cs_set:cpn {#1vectoronlyderivedunits} {\mandi@selectderivedunits{#2}{#3}{#4}}
219   \cs_set:cpn {vector#1onlyalternateunits} {\mandi@selectalternateunits{#2}{#3}{#4}}
220   \cs_set:cpn {#1vectoronlyalternateunits} {\mandi@selectalternateunits{#2}{#3}{#4}}
221 }
222 \NewDocumentCommand{\RenewVectorQuantity}{ m m O{#2} O{#2} }
223 {
224   \__mandi_renewvectorquantity:nnnn { #1 }{ #2 }{ #3 }{ #4 }
225 }

```

Defining a new physical constant.

```

226 \cs_new:Npn \__mandi_newphysicalconstant:nnnnnn #1#2#3#4#5#6#7
227 {
228   \cs_new:cpn {#1} {\units{\mandi@selectprecision{#3}{#4}}{\mandi@selectunits{#5}{#6}{#7}}}

```

```

229 \cs_new:cpn {#1mathsymbol} {#2}
230 \cs_new:cpn {#1approximatevalue} {#3}
231 \cs_new:cpn {#1precisevalue} {#4}
232 \cs_new:cpn {#1baseunits}
233   {\units{\mandi@selectprecision{#3}{#4}}{\mandi@selectbaseunits{#5}{#6}{#7}}}
234 \cs_new:cpn {#1derivedunits}
235   {\units{\mandi@selectprecision{#3}{#4}}{\mandi@selectderivedunits{#5}{#6}{#7}}}
236 \cs_new:cpn {#1alternateunits}
237   {\units{\mandi@selectprecision{#3}{#4}}{\mandi@selectalternateunits{#5}{#6}{#7}}}
238 \cs_new:cpn {#1onlybaseunits} {\mandi@selectbaseunits{#5}{#6}{#7}}
239 \cs_new:cpn {#1onlyderivedunits} {\mandi@selectderivedunits{#5}{#6}{#7}}
240 \cs_new:cpn {#1onlyalternateunits} {\mandi@selectalternateunits{#5}{#6}{#7}}
241 }
242 \NewDocumentCommand{\NewPhysicalConstant}{ m m m m m O{#5} O{#5} }
243 {
244   \__mandi_newphysicalconstant:nnnnnnn { #1 }{ #2 }{ #3 }{ #4 }{ #5 }{ #6 }{ #7 }
245 }

```

Redefining an existing physical constant.

```

246 \cs_new:Npn \__mandi_renewphysicalconstant:nnnnnnn #1#2#3#4#5#6#7
247 {
248   \cs_set:cpn {#1} {\units{\mandi@selectprecision{#3}{#4}}{\mandi@selectunits{#5}{#6}{#7}}}
249   \cs_set:cpn {#1mathsymbol} {#2}
250   \cs_set:cpn {#1approximatevalue} {#3}
251   \cs_set:cpn {#1precisevalue} {#4}
252   \cs_set:cpn {#1baseunits}
253     {\units{\mandi@selectprecision{#3}{#4}}{\mandi@selectbaseunits{#5}{#6}{#7}}}
254   \cs_set:cpn {#1derivedunits}
255     {\units{\mandi@selectprecision{#3}{#4}}{\mandi@selectderivedunits{#5}{#6}{#7}}}
256   \cs_set:cpn {#1alternateunits}
257     {\units{\mandi@selectprecision{#3}{#4}}{\mandi@selectalternateunits{#5}{#6}{#7}}}
258   \cs_set:cpn {#1onlybaseunits} {\mandi@selectbaseunits{#5}{#6}{#7}}
259   \cs_set:cpn {#1onlyderivedunits} {\mandi@selectderivedunits{#5}{#6}{#7}}
260   \cs_set:cpn {#1onlyalternateunits} {\mandi@selectalternateunits{#5}{#6}{#7}}
261 }
262 \NewDocumentCommand{\RenewPhysicalConstant}{ m m m m m O{#5} O{#5} }
263 {
264   \__mandi_renewphysicalconstant:nnnnnnn { #1 }{ #2 }{ #3 }{ #4 }{ #5 }{ #6 }{ #7 }
265 }
266 \ExplSyntaxOff

```

Define every quantity we need in introductory physics, alphabetically for convenience. This is really the core feature of **mandi** that no other package offers. There are commands for quantities that have no dimensions or units, and these quantities are defined for semantic completeness.

```

267 \NewVectorQuantity{acceleration}%
268   {\meter\usk\second\totheinversetwo}%
269   [\newton\per\kilogram]%
270   [\meter\per\second\tothetwo]%
271 \NewScalarQuantity{amount}%
272   {\mole}%
273 \NewVectorQuantity{angularacceleration}%
274   {\radian\usk\second\totheinversetwo}%
275   [\radian\per\second\tothetwo]%
276   [\radian\per\second\tothetwo]%
277 \NewScalarQuantity{angularfrequency}%
278   {\radian\usk\second\inverse}%
279   [\radian\per\second]%
280   [\radian\per\second]%

```


Angular impulse may also be $\text{newton}\cdot\text{usk}\cdot\text{meter}\cdot\text{usk}\cdot\text{second}$ or $\text{joule}\cdot\text{usk}\cdot\text{second}$.

```
281 \NewVectorQuantity{angularimpulse}%
282   {\kilogram\usk\meter\tothetwo\usk\second\inverse}%
283   [\kilogram\usk\meter\tothetwo\per\second]%
284   [\kilogram\usk\meter\tothetwo\per\second]%
```

Angular momentum may also be $\text{joule}\cdot\text{usk}\cdot\text{second}$ or $\text{newton}\cdot\text{usk}\cdot\text{meter}\cdot\text{usk}\cdot\text{second}$.

```
285 \NewVectorQuantity{angularmomentum}%
286   {\kilogram\usk\meter\tothetwo\usk\second\inverse}%
287   [\kilogram\usk\meter\tothetwo\per\second]%
288   [\kilogram\usk\meter\tothetwo\per\second]%

289 \NewVectorQuantity{angularvelocity}%
290   {\radian\usk\second\inverse}%
291   [\radian\per\second]%
292   [\radian\per\second]%

293 \NewScalarQuantity{area}%
294   {\meter\tothetwo}%
295 \NewScalarQuantity{areachargedensity}%
296   {\ampere\usk\second\usk\meter\totheinversetwo}%
297   [\coulomb\per\meter\tothetwo]%
298   [\coulomb\per\meter\tothetwo]%
299 \NewScalarQuantity{areamassdensity}%
300   {\kilogram\usk\meter\totheinversetwo}%
301   [\kilogram\per\meter\tothetwo]%
302   [\kilogram\per\meter\tothetwo]%
```

Capacitance may also be $\text{coulomb}\cdot\text{tothetwo}\cdot\text{per}\cdot\text{newton}\cdot\text{usk}\cdot\text{meter}$ or $\text{second}\cdot\text{per}\cdot\text{ohm}$.

```
303 \NewScalarQuantity{capacitance}%
304   {\ampere\tothetwo\usk\second\tothefour\usk\kilogram\inverse\usk\meter\totheinversetwo}%
305   [\farad]%
306   [\coulomb\per\volt]%
```

Charge may also be $\text{farad}\cdot\text{usk}\cdot\text{volt}$.

```
307 \NewScalarQuantity{charge}%
308   {\ampere\usk\second}%
309   [\coulomb]%
310   [\coulomb]%
```

Magnetic field times c may also be $\text{volt}\cdot\text{per}\cdot\text{meter}$.

```
311 \NewVectorQuantity{cmagneticfield}%
312   {\kilogram\usk\meter\usk\ampere\inverse\usk\second\totheinversethree}%
313   [\newton\per\coulomb]%
314   [\newton\per\coulomb]%

315 \NewScalarQuantity{conductance}%
316   {\ampere\tothetwo\usk\second\tothethree\usk\kilogram\inverse\usk\meter\totheinversetwo}%
317   [\siemens]%
318   [\ampere\per\volt]%
319 \NewScalarQuantity{conductivity}%
320   {\ampere\tothetwo\usk\second\tothethree\usk\kilogram\inverse\usk\meter\totheinversethree}%
321   [\siemens\per\meter]%
322   [\ampere\per\volt\usk\meter]%
323 \NewScalarQuantity{conventionalcurrent}%
324   {\ampere}%
325   [\coulomb\per\second]%
326   [\ampere]%
327 \NewScalarQuantity{current}%
328   {\ampere}%
329 \NewScalarQuantity{currentdensity}%
```

```

330 {\ampere\usk\meter\totheinversetwo}%
331 [\coulomb\per\second\usk\meter\tothetwo]%
332 [\ampere\per\meter\tothetwo]%
333 \NewScalarQuantity{dielectricconstant}%
334 {}%
335 \NewVectorQuantity{direction}%
336 {}%
337 \NewVectorQuantity{displacement}%
338 {\meter}%
339 \NewScalarQuantity{duration}%
340 {\second}%
341 \NewVectorQuantity{electricdipolemoment}%
342 {\ampere\usk\second\usk\meter}%
343 [\coulomb\usk\meter]%
344 [\coulomb\usk\meter]%
345 \NewVectorQuantity{electricfield}%
346 {\kilogram\usk\meter\usk\ampere\inverse\usk\second\totheinversethree}%
347 [\volt\per\meter]%
348 [\newton\per\coulomb]%
349 \NewScalarQuantity{electricflux}%
350 {\kilogram\usk\meter\tothethree\usk\ampere\inverse\usk\second\totheinversethree}%
351 [\volt\usk\meter]%
352 [\newton\usk\meter\tothetwo\per\coulomb]%

```

Electric potential, electric potential difference, and emf may also be \joule\per\coulomb.

```

353 \NewScalarQuantity{electricpotential}%
354 {\kilogram\usk\meter\tothetwo\usk\ampere\inverse\usk\second\totheinversethree}%
355 [\volt]%
356 [\volt]%
357 \NewScalarQuantity{electricpotentialdifference}%
358 {\kilogram\usk\meter\tothetwo\usk\ampere\inverse\usk\second\totheinversethree}%
359 [\volt]%
360 [\volt]%
361 \NewScalarQuantity{electroncurrent}%
362 {\second\inverse}%
363 [\ensuremath{\mathrm{e}}\per\second]%
364 [\ensuremath{\mathrm{e}}\per\second]%
365 \NewScalarQuantity{emf}%
366 {\kilogram\usk\meter\tothetwo\usk\ampere\inverse\usk\second\totheinversethree}%
367 [\volt]%
368 [\volt]%

```

Energy may also be \newton\usk\meter.

```

369 \NewScalarQuantity{energy}%
370 {\kilogram\usk\meter\tothetwo\usk\second\totheinversetwo}%
371 [\joule]%
372 [\joule]%
373 \NewScalarQuantity{energyinev}%
374 {\electronvolt}%
375 \NewScalarQuantity{energyinkev}%
376 {\kilolectronvolt}%
377 \NewScalarQuantity{energyinmev}%
378 {\megaelectronvolt}%
379 \NewScalarQuantity{energydensity}%
380 {\kilogram\usk\meter\inverse\usk\second\totheinversetwo}%
381 [\joule\per\meter\tothethree]%
382 [\joule\per\meter\tothethree]%
383 \NewScalarQuantity{energyflux}%

```

```

384  {\kilogram\usk\second\totheinversethree}%
385  [\watt\per\meter\tothetwo]%
386  [\watt\per\meter\tothetwo]%
387 \NewScalarQuantity{entropy}%
388  {\kilogram\usk\meter\tothetwo\usk\second\totheinversetwo\usk\kelvin\inverse}%
389  [\joule\per\kelvin]%
390  [\joule\per\kelvin]%

```

Force may also be \kilogram\usk\meter\per\second\tothetwo.

```

391 \NewVectorQuantity{force}%
392  {\kilogram\usk\meter\usk\second\totheinversetwo}%
393  [\newton]%
394  [\newton]%
395 \NewScalarQuantity{frequency}%
396  {\second\inverse}%
397  [\hertz]%
398  [\hertz]%
399 \NewVectorQuantity{gravitationalfield}%
400  {\meter\usk\second\totheinversetwo}%
401  [\newton\per\kilogram]%
402  [\newton\per\kilogram]%
403 \NewScalarQuantity{gravitationalpotential}%
404  {\meter\tothetwo\usk\second\totheinversetwo}%
405  [\joule\per\kilogram]%
406  [\joule\per\kilogram]%
407 \NewScalarQuantity{gravitationalpotentialdifference}%
408  {\meter\tothetwo\usk\second\totheinversetwo}%
409  [\joule\per\kilogram]%
410  [\joule\per\kilogram]%
411 \NewVectorQuantity{impulse}%
412  {\kilogram\usk\meter\usk\second\inverse}%
413  [\newton\usk\second]%
414  [\newton\usk\second]%
415 \NewScalarQuantity{indexofrefraction}%
416  {}%

```

Inductance may also be \square\meter\usk\kilogram\per\coulomb\tothetwo or \Wb\per\ampere.

```

417 \NewScalarQuantity{inductance}%
418  {\kilogram\usk\meter\tothetwo\usk\ampere\totheinversetwo\usk\second\totheinversetwo}%
419  [\henry]%
420  [\volt\usk\second\per\ampere]%
421 \NewScalarQuantity{linearchargedensity}%
422  {\ampere\usk\second\usk\meter\inverse}%
423  [\coulomb\per\meter]%
424  [\coulomb\per\meter]%
425 \NewScalarQuantity{linearmassdensity}%
426  {\kilogram\usk\meter\inverse}%
427  [\kilogram\per\meter]%
428  [\kilogram\per\meter]%
429 \NewScalarQuantity{lorentzfactor}%
430  {}%
431 \NewScalarQuantity{luminousintensity}%
432  {\candela}%

```

There is another convention for magnetic charge, so be careful.

```

433 \NewScalarQuantity{magneticcharge}%
434  {\ampere\usk\meter}%

```

```

435 \NewVectorQuantity{magneticdipolemoment}%
436   {\ampere\usk\meter\tothetwo}%
437   [\ampere\usk\meter\tothetwo]%
438   [\joule\per\tesla]%

Magnetic field may also be \Wb\per\meter\tothetwo.

439 \NewVectorQuantity{magneticfield}%
440   {\kilogram\usk\ampere\inverse\usk\second\totheinversetwo}%
441   [\newton\per\ampere\usk\meter]%
442   [\tesla]%

Magnetic flux may also be \Wb or \joule\per\ampere.

443 \NewScalarQuantity{magneticflux}%
444   {\kilogram\usk\meter\tothetwo\usk\ampere\inverse\usk\second\totheinversetwo}%
445   [\tesla\usk\meter\tothetwo]%
446   [\volt\usk\second]%

447 \NewScalarQuantity{mass}%
448   {\kilogram}%

449 \NewScalarQuantity{mobility}%
450   {\kilogram\usk\meter\tothetwo\usk\ampere\inverse\usk\second\totheinversefour}%
451   [\meter\tothetwo\per\volt\usk\second]%
452   [\coulomb\usk\meter\per\newton\usk\second]%

453 \NewScalarQuantity{momentofinertia}%
454   {\kilogram\usk\meter\tothetwo}%
455   [\joule\usk\second\tothetwo]%
456   [\kilogram\usk\meter\tothetwo]%

457 \NewVectorQuantity{momentum}%
458   {\kilogram\usk\meter\usk\second\inverse}%
459   [\kilogram\usk\meter\per\second]%
460   [\kilogram\usk\meter\per\second]%

461 \NewVectorQuantity{momentumflux}%
462   {\kilogram\usk\meter\inverse\usk\second\totheinversetwo}%
463   [\newton\per\meter\tothetwo]%
464   [\newton\per\meter\tothetwo]%

465 \NewScalarQuantity{numberdensity}%
466   {\meter\totheinversethree}%
467   [\per\meter\tothethree]%
468   [\per\meter\tothethree]%

469 \NewScalarQuantity{permeability}%
470   {\kilogram\usk\meter\usk\ampere\totheinversetwo\usk\second\totheinversetwo}%
471   [\henry\per\meter]%
472   [\tesla\usk\meter\per\ampere]%

473 \NewScalarQuantity{permittivity}%
474   {\ampere\tothetwo\usk\second\tothefour\usk\kilogram\inverse\usk\meter\totheinversethree}%
475   [\farad\per\meter]%
476   [\coulomb\tothetwo\per\newton\usk\meter\tothetwo]%

477 \NewScalarQuantity{planeangle}%
478   {\meter\usk\meter\inverse}%
479   [\radian]%
480   [\radian]%

481 \NewScalarQuantity{polarizability}%
482   {\ampere\tothetwo\usk\second\tothefour\usk\kilogram\inverse}%
483   [\coulomb\usk\meter\tothetwo\per\volt]%
484   [\coulomb\tothetwo\usk\meter\per\newton]%

485 \NewScalarQuantity{power}%
486   {\kilogram\usk\meter\tothetwo\usk\second\totheinversethree}%
487   [\watt]%
488   [\joule\per\second]%

489 \NewVectorQuantity{poynting}%

```

```

490  {\kilogram\usk\second\totheinversethree}%
491  [\watt\per\meter\tothetwo]%
492  [\watt\per\meter\tothetwo]%
493 \NewScalarQuantity{pressure}%
494  {\kilogram\usk\meter\inverse\usk\second\totheinversetwo}%
495  [\pascal]%
496  [\newton\per\meter\tothetwo]%
497 \NewScalarQuantity{relativepermeability}%
498  {}%
499 \NewScalarQuantity{relativepermittivity}%
500  {}%

Resistance may also be \volt\per\ampere.

501 \NewScalarQuantity{resistance}%
502  {\kilogram\usk\meter\tothetwo\usk\ampere\totheinversetwo\usk\second\totheinversethree}%
503  [\ohm]%
504  [\ohm]%

505 \NewScalarQuantity{resistivity}%
506  {\kilogram\usk\meter\tothethree\usk\ampere\totheinversetwo\usk\second\totheinversethree}%
507  [\ohm\usk\meter]%
508  [\volt\usk\meter\per\ampere]%
509 \NewScalarQuantity{solidangle}%
510  {\meter\tothetwo\usk\meter\totheinversetwo}%
511  [\steradian]%
512  [\steradian]%
513 \NewScalarQuantity{specificheatcapacity}%
514  {\meter\tothetwo\usk\second\totheinversetwo\usk\kelvin\inverse}%
515  [\joule\per\kelvin\usk\kilogram]%
516  [\joule\per\kelvin\usk\kilogram]
517 \NewScalarQuantity{springstiffness}%
518  {\kilogram\usk\second\totheinversetwo}%
519  [\newton\per\meter]%
520  [\newton\per\meter]%

Spring stretch is really just a displacement.

521 \NewScalarQuantity{springstretch}%
522  {\meter}%

523 \NewScalarQuantity{stress}%
524  {\kilogram\usk\meter\inverse\usk\second\totheinversetwo}%
525  [\pascal]%
526  [\newton\per\meter\tothetwo]%
527 \NewScalarQuantity{strain}%
528  {}%
529 \NewScalarQuantity{temperature}%
530  {\kelvin}%
531 \NewVectorQuantity{torque}%
532  {\kilogram\usk\meter\tothetwo\usk\second\totheinversetwo}%
533  [\newton\usk\meter]%
534  [\newton\usk\meter]%
535 \NewVectorQuantity{velocity}%
536  {\meter\usk\second\inverse}%
537  [\meter\per\second]%
538  [\meter\per\second]%
539 \NewVectorQuantity{velocityc}%
540  {\lightspeed}%
541  [\lightspeed]%
542  [\lightspeed]%
543 \NewScalarQuantity{volume}%

```

```

544 {\meter\tothethree}%
545 \NewScalarQuantity{volumechargedensity}%
546 {\ampere\usk\second\per\meter\totheinversethree}%
547 [\coulomb\per\meter\tothethree]%
548 [\coulomb\per\meter\tothethree]%
549 \NewScalarQuantity{volumemassdensity}%
550 {\kilogram\usk\meter\totheinversethree}%
551 [\kilogram\per\meter\tothethree]%
552 [\kilogram\per\meter\tothethree]%

Wavelength is really just a displacement.

553 \NewScalarQuantity{wavelength}%
554 {\meter}%

555 \NewVectorQuantity{wavenumber}%
556 {\meter\inverse}%
557 [\per\meter]%
558 [\per\meter]%

Work may also be \newton\usk\meter but this is discouraged to avoid confusion with torque.

559 \NewScalarQuantity{work}%
560 {\kilogram\usk\meter\tothetwo\usk\second\totheinversetwo}%
561 [\joule]%
562 [\joule]%

Young's modulus is really just a stress.

563 \NewScalarQuantity{youngsmodulus}%
564 {\kilogram\usk\meter\inverse\usk\second\totheinversetwo}%
565 [\pascal]%
566 [\newton\per\meter\tothetwo]%

We need a better glyph for Planck's constant over  $2\pi$ .

567 \AtBeginDocument{%
568 \DeclareRobustCommand{\hbar}{\mathpalette\hbar@\relax\symup{h}}%
569 }%
570 \newcommand*{\hbar@}[2]{%
571 \makebox[0pt][l]{\raisebox{-0.07\height}{\(\m@th#1\mkern-2mu\mathchar"AF\)}}}%

Optional line to make the bar thicker; must use -0.11 in \raisebox.

572 \makebox[0pt][l]{\raisebox{-0.11\height}{\(\m@th#1\mkern-2mu\mathchar"AF\)}}%
573 }%

Define physical constants for introductory physics, again alphabetically for convenience.

574 \NewPhysicalConstant{avogadro}%
575 {\symup{N}_{A}}}%
576 {6\times10^{23}}{6.02214076\times10^{23}}}%
577 {\mole\inverse}%
578 [\per\mole]%
579 [\per\mole]%

\biotsavartconstant is an alias for \mzofp.

580 \NewPhysicalConstant{biotsavartconstant}%
581 {\symup{\frac{\mu_0}{4\pi}}}%
582 {10^{-7}}{10^{-7}}}%
583 {\kilogram\usk\meter\usk\ampere\totheinversetwo\usk\second\totheinversetwo}%
584 [\henry\per\meter]%
585 [\tesla\usk\meter\per\ampere]%
586 \NewPhysicalConstant{bohrradius}%
587 {\symup{a_0}}}%
588 {5.3\times10^{-11}}{5.29177210903\times10^{-11}}}%

```

```

589 {\meter}%
590 \NewPhysicalConstant{boltzmann}%
591 {\symup{k_{B}}}%
592 {1.4\times10^{-23}}{1.380649\times10^{-23}}%
593 {\kilogram\usk\meter\tothetwo\usk\second\totheinversetwo\usk\kelvin\inverse}%
594 [\joule\per\kelvin]%
595 [\joule\per\kelvin]%

\coulombconstant is an alias for \oofpez.

596 \NewPhysicalConstant{coulombconstant}%
597 {\symup{\frac{1}{4\pi\epsilon_0}}}%
598 {9\times10^9}{8.9875517923\times10^9}%
599 {\kilogram\usk\meter\tothethree\usk\ampere\totheinversetwo\usk\second\totheinversefour}%
600 [\meter\per\farad]%
601 [\newton\usk\meter\tothetwo\per\coulomb\tothetwo]%
602 \NewPhysicalConstant{earthmass}%
603 {\symup{M_{Earth}}}%
604 {6.0\times10^{24}}{5.9722\times10^{24}}%
605 {\kilogram}%
606 \NewPhysicalConstant{earthmoondistance}%
607 {\symup{d_{EM}}}%
608 {3.8\times10^8}{3.81550\times10^8}%
609 {\meter}%
610 \NewPhysicalConstant{earthradius}%
611 {\symup{R_{Earth}}}%
612 {6.4\times10^6}{6.3781\times10^6}%
613 {\meter}%
614 \NewPhysicalConstant{earthsundistance}%
615 {\symup{d_{ES}}}%
616 {1.5\times10^{11}}{1.496\times10^{11}}%
617 {\meter}%
618 \NewPhysicalConstant{electroncharge}%
619 {\symup{q_e}}%
620 {-\elementarychargeapproximatevalue}{-\elementarychargeprecisevalue}%
621 {\ampere\usk\second}%
622 [\coulomb]%
623 [\coulomb]%
624 \NewPhysicalConstant{electronCharge}%
625 {\symup{Q_e}}%
626 {-\elementarychargeapproximatevalue}{-\elementarychargeprecisevalue}%
627 {\ampere\usk\second}%
628 [\coulomb]%
629 [\coulomb]%
630 \NewPhysicalConstant{electronmass}%
631 {\symup{m_e}}%
632 {9.1\times10^{-31}}{9.1093837015\times10^{-31}}%
633 {\kilogram}%
634 \NewPhysicalConstant{elementarycharge}%
635 {\symup{e}}%
636 {1.6\times10^{-19}}{1.602176634\times10^{-19}}%
637 {\ampere\usk\second}%
638 [\coulomb]%
639 [\coulomb]%
640 \NewPhysicalConstant{finestructure}%
641 {\symup{\alpha}}%
642 {\frac{1}{137}}{7.2973525693\times10^{-3}}%
643 {}%
644 \NewPhysicalConstant{hydrogenmass}%
645 {\symup{m_H}}%

```

```

646 {1.7\times10^{-27}}{1.6737236\times10^{-27}}%
647 {\kilogram}%
648 \NewPhysicalConstant{moonearthdistance}%
649 {\symup{d_{ME}}}%
650 {3.8\times10^8}{3.81550\times10^8}%
651 {\meter}%
652 \NewPhysicalConstant{moonmass}%
653 {\symup{M_{Moon}}}%
654 {7.3\times10^{22}}{7.342\times10^{22}}%
655 {\kilogram}%
656 \NewPhysicalConstant{moonradius}%
657 {\symup{R_{Moon}}}%
658 {1.7\times10^6}{1.7371\times10^6}%
659 {\meter}%
660 \NewPhysicalConstant{mzofp}%
661 {\symup{\frac{\mu_o}{4\pi}}}%
662 {10^{-7}}{10^{-7}}%
663 {\kilogram\usk\meter\usk\ampere\totheinversetwo\usk\second\totheinversetwo}%
664 [\henry\per\meter]%
665 [\tesla\usk\meter\per\ampere]%
666 \NewPhysicalConstant{neutronmass}%
667 {\symup{m_n}}%
668 {1.7\times10^{-27}}{1.67492749804\times10^{-27}}%
669 {\kilogram}%
670 \NewPhysicalConstant{oofpez}%
671 {\symup{\frac{1}{4\pi\epsilon_o}}}%
672 {9\times10^9}{8.9875517923\times10^9}%
673 {\kilogram\usk\meter\tothethree\usk\ampere\totheinversetwo\usk\second\totheinversefour}%
674 [\meter\per\farad]%
675 [\newton\usk\meter\tothetwo\per\coulomb\tothetwo]%
676 \NewPhysicalConstant{oofpezcs}%
677 {\symup{\frac{1}{4\pi\epsilon_o c^2}}}%
678 {10^{-7}}{10^{-7}}%
679 {\kilogram\usk\meter\usk\ampere\totheinversetwo\usk\second\totheinversetwo}%
680 [\tesla\usk\meter\tothetwo]%
681 [\newton\usk\second\tothetwo\per\coulomb\tothetwo]%
682 \NewPhysicalConstant{planck}%
683 {\symup{h}}%
684 {6.6\times10^{-34}}{6.62607015\times10^{-34}}%
685 {\kilogram\usk\meter\tothetwo\usk\second\inverse}%
686 [\joule\usk\second]%
687 [\joule\usk\second]

```

See <https://tex.stackexchange.com/a/448565/218142>.

```

688 \NewPhysicalConstant{planckbar}%
689 {\hbar}%
690 {1.1\times10^{-34}}{1.054571817\times10^{-34}}%
691 {\kilogram\usk\meter\tothetwo\usk\second\inverse}%
692 [\joule\usk\second]%
693 [\joule\usk\second]
694 \NewPhysicalConstant{planckc}%
695 {\symup{hc}}%
696 {2.0\times10^{-25}}{1.98644586\times10^{-25}}%
697 {\kilogram\usk\meter\tothethree\usk\second\totheinversetwo}%
698 [\joule\usk\meter]%
699 [\joule\usk\meter]
700 \NewPhysicalConstant{protoncharge}%
701 {\symup{q_p}}%
702 {+\elementarychargeapproximatevalue}{+\elementarychargeprecisevalue}%

```



```

703 {\ampere\usk\second}%
704 [\coulomb]%
705 [\coulomb]%
706 \NewPhysicalConstant{protonCharge}%
707 {\symup{Q_p}}%
708 {+\elementarychargeapproximatevalue}{+\elementarychargeprecisevalue}%
709 {\ampere\usk\second}%
710 [\coulomb]%
711 [\coulomb]%
712 \NewPhysicalConstant{protonmass}%
713 {\symup{m_p}}%
714 {1.7\times10^{-27}}{1.672621898\times10^{-27}}%
715 {\kilogram}%
716 \NewPhysicalConstant{rydberg}%
717 {\symup{R_{\infty}}}%
718 {1.1\times10^7}{1.0973731568160\times10^7}%
719 {\meter\inverse}%
720 \NewPhysicalConstant{speedoflight}%
721 {\symup{c}}%
722 {3\times10^8}{2.99792458\times10^8}%
723 {\meter\usk\second\inverse}%
724 [\meter\per\second]%
725 [\meter\per\second]
726 \NewPhysicalConstant{stefanboltzmann}%
727 {\symup{\sigma}}%
728 {5.7\times10^{-8}}{5.670374\times10^{-8}}%
729 {\kilogram\usk\second\totheinversethree\usk\kelvin\totheinversefour}%
730 [\watt\per\meter\tothetwo\usk\kelvin\tothefour]%
731 [\watt\per\meter\tothetwo\usk\kelvin\tothefour]
732 \NewPhysicalConstant{sunearthdistance}%
733 {\symup{d_{SE}}}%
734 {1.5\times10^{11}}{1.496\times10^{11}}%
735 {\meter}%
736 \NewPhysicalConstant{sunmass}%
737 {\symup{M_{Sun}}}%
738 {2.0\times10^{30}}{1.98855\times10^{30}}%
739 {\kilogram}%
740 \NewPhysicalConstant{sunradius}%
741 {\symup{R_{Sun}}}%
742 {7.0\times10^8}{6.957\times10^8}%
743 {\meter}%
744 \NewPhysicalConstant{surfacegravfield}%
745 {\symup{g}}%
746 {9.8}{9.807}%
747 {\meter\usk\second\totheinversetwo}%
748 [\newton\per\kilogram]%
749 [\newton\per\kilogram]%

```

The gravitational constant may also have units of $\text{joule}\,\text{usk}\,\text{meter}\,\text{per}\,\text{kilogram}\,\text{tothetwo}$.

```

750 \NewPhysicalConstant{universalgrav}%
751 {\symup{G}}%
752 {6.7\times10^{-11}}{6.67430\times10^{-11}}%
753 {\meter\tothethree\usk\kilogram\inverse\usk\second\totheinversetwo}%
754 [\newton\usk\meter\tothetwo\per\kilogram\tothetwo]%
755 [\newton\usk\meter\tothetwo\per\kilogram\tothetwo]%

```

As of 2018 the vacuum permeability is no longer defined as $4\pi \times 10^{-7}$.

```

756 \NewPhysicalConstant{vacuumpermeability}%
757 {\symup{\mu_o}}%

```

```

758 {4\pi\times10^{-7}}{4\pi\times10^{-7}}%
759 {\kilogram\usk\meter\usk\ampere\totheinversetwo\usk\second\totheinversetwo}%
760 [\henry\per\meter]%
761 [\tesla\usk\meter\per\ampere]%
762 \NewPhysicalConstant{vacuumpermittivity}%
763 {\symup{\epsilon_o}}%
764 {9\times10^{-12}}{8.854187817\times10^{-12}}%
765 {\ampere\tothetwo\usk\second\tothefour\usk\kilogram\inverse\usk\meter\totheinversethree}%
766 [\farad\per\meter]%
767 [\coulomb\tothetwo\per\newton\usk\meter\tothetwo]%

```

Diagnostic commands to provide sanity checks on commands that represent physical quantities and constants.

```

768 \ExplSyntaxOn
769 \NewDocumentCommand{\@aux}{ m }
770 {
771   \use:c { #1 }
772 }
773 \NewDocumentCommand{\@auy}{ m }
774 {
775   \normalfont\ttfamily\token_to_str:c { #1 }
776 }
777 \ExplSyntaxOff
778 \newcolumnntype{M}{>{\(}p{0.25\linewidth}<{\)}}
779 \NewDocumentCommand{\CheckQuantity}{ m }
780 {%
781   \begin{center}
782     \begin{tabular}{MMM}
783       \textbf{command} & & \multicolumn{2}{l}{\@auy{#1}} & \tabularnewline
784       \text{\textbf{base}} & & \text{\textbf{derived}} & & \text{\textbf{alternate}} & \tabularnewline
785       \@aux{#1onlybaseunits} & & \@aux{#1onlyderivedunits} & & \@aux{#1onlyalternateunits} & \tabularnewline
786     \end{tabular}
787   \end{center}
788 }%
789 \NewDocumentCommand{\CheckConstant}{ m }
790 {%
791   \begin{center}
792     \begin{tabular}{MMM}
793       \textbf{command} & & \multicolumn{2}{l}{\@auy{#1}} & \tabularnewline
794       \text{\textbf{symbol}} & & \text{\textbf{approximate}} & & \text{\textbf{precise}} & \tabularnewline
795       \@aux{#1mathsymbol} & & \@aux{#1approximatevalue} & & \@aux{#1precisevalue} & \tabularnewline
796       \text{\textbf{base}} & & \text{\textbf{derived}} & & \text{\textbf{alternate}} & \tabularnewline
797       \@aux{#1onlybaseunits} & & \@aux{#1onlyderivedunits} & & \@aux{#1onlyalternateunits} & \tabularnewline
798     \end{tabular}
799   \end{center}
800 }%

```

`\mivector`^{P.34} is a workhorse command.

See <https://tex.stackexchange.com/a/39054/218142>.

```

801 \ExplSyntaxOn
802 \NewDocumentCommand{\mivector}{ O{,} m o }
803 {
804   \__mandi_vector:nn { #1 } { #2 }
805   \IfValueT{#3}{\,{#3}}
806 }
807 \seq_new:N \l__mandi_list_seq
808 \cs_new_protected:Npn \__mandi_vector:nn #1#2
809 {
810   \seq_set_split:Nnn \l__mandi_list_seq { , } { #2 }
811   \int_compare:nT { \seq_count:N \l__mandi_list_seq = 1 }

```

```

812 {
813   \msg_new:nnnn { mandi } { onecomponent }
814   {
815     More~than~one~component~expected.      \iow_newline:
816     You~provided~one~component~to~a~command \iow_newline:
817     that~expects~a~vector.~Either~you~don't \iow_newline:
818     need~a~vector~here~or~you~didn't~supply \iow_newline:
819     all~the~components.
820   }
821   {
822     Decide~whether~or~not~you~really~need~a~vector~command~here. \iow_newline:
823     \msg_see_documentation_text:n { mandi }
824   }
825   \msg_fatal:nn { mandi } { onecomponent }
826
827 }
828 \left\langle
829   \seq_use:Nnnn \l__mandi_list_seq { #1 } { #1 } { #1 }
830 \right\rangle
831 }
832 \ExplSyntaxOff

```

7 The mandistudent Package

mandi comes with an accessory package mandistudent, which extends mandi with a collection of commands physics students can use for writing problem solutions. This package focuses on the most frequently needed tools. These commands should always be used in math mode. mandistudent requires, and loads, mandi but mandi doesn't require, and doesn't load, mandistudent.

Load mandistudent as you would any package in your preamble. There are no package options.

```
\usepackage{mandistudent}
```

```
\mandistudentversion
```

Typesets the current version and build date.

```
The version is \mandistudentversion{} and is a stable build.
```

```
The version is v3.2.2a dated 2024-09-12 and is a stable build.
```

7.1 Traditional Vector Notation

U 2021-09-18

```
\vec{\langle symbol \rangle} [\langle labels \rangle]
```

(use this variant for boldface notation)

U 2021-09-18

```
\vec*{\langle symbol \rangle} [\langle labels \rangle]
```

(use this variant for arrow notation)

Powerful and intelligent command for symbolic vector notation. The mandatory argument is the symbol for the vector quantity. The optional label(s) consists of superscripts and/or subscripts and can be mathematical or textual in nature. If textual, be sure to wrap them in `\symup{...}` for proper typesetting. The starred variant gives arrow notation whereas without the star you get boldface notation. Subscript and superscript labels can be arbitrarily mixed, and order doesn't matter. This command redefines the default L^AT_EX 2_ε `\vec` command.

```
\( \vec{p} ) \qquad \qquad \qquad \\
\(\ \vec{p}_{2} ) \qquad \qquad \qquad \\
\(\ \vec{p}^{\symup{ball}} ) \qquad \qquad \qquad \\
\(\ \vec{p}_{\symup{final}} ) \qquad \qquad \qquad \\
\(\ \vec{p}^{\symup{ball}}_{\symup{final}} ) \qquad \qquad \qquad \\
\(\ \vec{p}^{\symup{final}}_{\symup{ball}} ) \qquad \qquad \qquad \\
\(\ \vec*{p} ) \qquad \qquad \qquad \\
\(\ \vec*{p}_{2} ) \qquad \qquad \qquad \\
\(\ \vec*{p}^{\symup{ball}} ) \qquad \qquad \qquad \\
\(\ \vec*{p}_{\symup{final}} ) \qquad \qquad \qquad \\
\(\ \vec*{p}^{\symup{ball}}_{\symup{final}} ) \qquad \qquad \qquad \\
\(\ \vec*{p}^{\symup{final}}_{\symup{ball}} ) \qquad \qquad \qquad
```

p
*p*₂
p^{ball}
p^{final}_{ball}
p^{final}_{final}
p^{ball}_{ball}
 \vec{p}
 \vec{p}_2
 \vec{p}^{ball}
 \vec{p}^{final}
 \vec{p}^{final} _{ball}
 \vec{p}^{final} _{final}

U 2021-09-18

```
\dirvec{\langle symbol \rangle} [\langle labels \rangle]
```

(use this variant for boldface notation)

\dirvec*{*symbol*}[*labels*]

(use this variant for arrow notation)

Powerful and intelligent command for typesetting the direction of a vector. The options are the same as those for `\vec`.

```
\( \dirvec{p} \)          \\\
\(\ \dirvec{p}_{2} \)      \\\
\(\ \dirvec{p}^{\symup{ball}} \) \\\
\(\ \dirvec{p}_{\symup{final}} \) \\\
\(\ \dirvec{p}^{\symup{ball}}_{\symup{final}} \) \\\
\(\ \dirvec{p}_{\symup{final}}_{\symup{ball}} \) \\\
\(\ \dirvec*{p} \)          \\\
\(\ \dirvec*{p}_{2} \)      \\\
\(\ \dirvec*{p}^{\symup{ball}} \) \\\
\(\ \dirvec*{p}_{\symup{final}} \) \\\
\(\ \dirvec*{p}^{\symup{ball}}_{\symup{final}} \) \\\
\(\ \dirvec*{p}_{\symup{final}}_{\symup{ball}} \) \\\
```

$$\begin{array}{c} \widehat{p} \\ \widehat{p}_2 \\ \widehat{p}_{\text{ball}} \\ \widehat{p}_{\text{final}} \\ \widehat{p}_{\text{final}} \\ \widehat{p}_{\text{ball}} \\ \widehat{p} \\ \widehat{p}_2 \\ \widehat{p}_{\text{ball}} \\ \widehat{p} \\ \widehat{p}_{\text{final}} \\ \widehat{p}_{\text{ball}} \\ \widehat{p}_{\text{final}} \\ \widehat{p}_{\text{final}} \\ \widehat{p}_{\text{ball}} \end{array}$$
\zerovec

(use this variant for boldface notation)

\zerovec*

(use this variant for arrow notation)

Command for typesetting the zero vector. The starred variant gives arrow notation. Without the star you get boldface notation.

```
\( \zerovec \) \\\
\(\ \zerovec* \) \\\
```

$$\begin{array}{c} \mathbf{0} \\ \vec{0} \end{array}$$
\changein

Semantic alias for `\Delta`.

```
\( \changein t \) \\\
\(\ \changein \vec{p} \) \\\
```

$$\begin{array}{c} \Delta t \\ \Delta p \end{array}$$
\doublebars[*size*]{*quantity*}

(double bars)

\doublebars*[*size*]{*quantity*}

(double bars for fractions)

\singlebars[*size*]{*quantity*}

(single bars)

\singlebars*[*size*]{*quantity*}

(single bars for fractions)

\anglebrackets[*size*]{*quantity*}

(angle brackets)

\anglebrackets*[*size*]{*quantity*}

(angle brackets for fractions)

\parentheses[*size*]{*quantity*}

(parentheses)

\parentheses*[*size*]{*quantity*}

(parentheses for fractions)

\squarebrackets[*size*]{*quantity*}

(square brackets)

\squarebrackets*[*size*]{*quantity*}

(square brackets for fractions)

`\curlybraces` [*size*] {*quantity*} (curly braces)
`\curlybraces*` [*size*] {*quantity*} (curly braces for fractions)

If no argument is given, a placeholder is provided. Sizers like `\big`, `\Big`, `\bigg`, and `\Bigg` can be optionally specified. Beginners are encouraged not to use them. See the [mathtools](#) package documentation for details.

```
\[ \doublebars{} \]
\[ \doublebars{\vec{a}} \]
\[ \doublebars*{\frac{\vec{a}}{3}} \]
\[ \doublebars[\Bigg]{\frac{\vec{a}}{3}} \]
```

$$\| \cdot \|$$

$$\| \boldsymbol{a} \|$$

$$\left\| \frac{\boldsymbol{a}}{3} \right\|$$

$$\left\| \frac{\boldsymbol{a}}{3} \right\|$$

```
\[ \singlebars{} \]
\[ \singlebars{x} \]
\[ \singlebars*{\frac{x}{3}} \]
\[ \singlebars[\Bigg]{\frac{x}{3}} \]
```

$$| \cdot |$$

$$| x |$$

$$\left| \frac{x}{3} \right|$$

$$\left| \frac{x}{3} \right|$$

```
\[ \anglebrackets{} \]
\[ \anglebrackets{\vec{a}} \]
\[ \anglebrackets*{\frac{\vec{a}}{3}} \]
\[ \anglebrackets[\Bigg]{\frac{\vec{a}}{3}} \]
```

$$\langle \cdot \rangle$$

$$\langle \boldsymbol{a} \rangle$$

$$\left\langle \frac{\boldsymbol{a}}{3} \right\rangle$$

$$\left\langle \frac{\boldsymbol{a}}{3} \right\rangle$$

```
\[ \parentheses{} \]
\[ \parentheses{x} \]
\[ \parentheses*{\frac{x}{3}} \]
\[ \parentheses[\Bigg]{\frac{x}{3}} \]
```

$$(\cdot)$$

$$(x)$$

$$\left(\frac{x}{3} \right)$$

$$\left(\frac{x}{3} \right)$$

$\backslash[\ \squarebrackets{} \ \backslash]$	$[\cdot]$
$\backslash[\ \squarebrackets{x} \ \backslash]$	$[x]$
$\backslash[\ \squarebrackets*{\frac{x}{3}} \ \backslash]$	$\left[\frac{x}{3}\right]$
$\backslash[\ \squarebrackets[\Big]{\frac{x}{3}} \ \backslash]$	$\left[\frac{x}{3}\right]$

$\backslash[\ \curlybraces{} \ \backslash]$	$\{\cdot\}$
$\backslash[\ \curlybraces{x} \ \backslash]$	$\{x\}$
$\backslash[\ \curlybraces*{\frac{x}{3}} \ \backslash]$	$\left\{\frac{x}{3}\right\}$
$\backslash[\ \curlybraces[\Big]{\frac{x}{3}} \ \backslash]$	$\left\{\frac{x}{3}\right\}$

N 2021-02-21
N 2021-02-21
N 2021-02-21
N 2021-02-21
N 2021-02-21
N 2021-02-21

$\backslash\text{magnitude}$ $[\langle size \rangle]{\langle quantity \rangle}$ (alias for double bars)
 $\backslash\text{magnitude*}$ $[\langle size \rangle]{\langle quantity \rangle}$ (alias for double bars for fractions)
 $\backslash\text{norm}$ $[\langle size \rangle]{\langle quantity \rangle}$ (alias for double bars)
 $\backslash\text{norm*}$ $[\langle size \rangle]{\langle quantity \rangle}$ (alias for double bars for fractions)
 $\backslash\text{absolutevalue}$ $[\langle size \rangle]{\langle quantity \rangle}$ (alias for single bars)
 $\backslash\text{absolutevalue*}$ $[\langle size \rangle]{\langle quantity \rangle}$ (alias for single bars for fractions)

Semantic aliases. Use $\backslash\text{magnitude}$ or $\backslash\text{magnitude*}$ to typeset the magnitude of a vector.

$\backslash[\ \text{magnitude}\{\text{vec}\{p\}\} \ \backslash]$	$\ \boldsymbol{p}\ $
$\backslash[\ \text{magnitude}\{\text{vec*}\{p\}\} \ \backslash]$	$\ \vec{p}\ $
$\backslash[\ \text{magnitude*}\{\text{vec}\{p\}_{\text{symup}\{final\}}\} \ \backslash]$	$\ \boldsymbol{p}_{\text{final}}\ $
$\backslash[\ \text{magnitude*}\{\text{vec*}\{p\}_{\text{symup}\{final\}}\} \ \backslash]$	$\ \vec{p}_{\text{final}}\ $

N 2021-04-06
N 2021-04-06

$\backslash\text{parallelto}$
 $\backslash\text{perpendicularto}$

Commands for geometric relationships, mainly intended for subscripts.

$\backslash(\ \text{vec}\{F\}_{\text{parallelto}} + \text{vec}\{F\}_{\text{perpendicularto}} \ \backslash)$	$\boldsymbol{F}_{\parallel} + \boldsymbol{F}_{\perp}$
--	---

7.2 Problems and Annotated Problem Solutions

U 2023-08-01 `\begin{physicsproblem}{\langle title \rangle}` `\end{physicsproblem}` `\langle problem \rangle` (use this variant for vertical lists)

U 2023-08-01 `\begin{physicsproblem*}{\langle title \rangle}` `\end{physicsproblem*}` `\langle problem \rangle` (use this variant for in-line lists)

U 2023-08-01 `\begin{parts}{\langle title \rangle}` `\end{parts}` `\langle problem \rangle` (provides problem parts)

Provides an environment for stating physics problems. Each problem will begin on a new page. See the examples for how to handle single and multiple part problems. Due to incompatibilities with BEAMER and the `enumitem` package, these environments are not defined if BEAMER is loaded.

U 2023-08-01 `\problempart`
Denotes a part of a problem within a `parts` environment. This command is not defined if BEAMER is loaded.

```
\begin{physicsproblem}{Problem 1}
  This is a physics problem with no parts.
\end{physicsproblem}
```

Problem 1

This is a physics problem with no parts.

```
\begin{physicsproblem}{Problem 2}
  This is a physics problem with multiple parts.
  The list is vertical.
  \begin{parts}
    \problempart This is the first part.
    \problempart This is the second part.
    \problempart This is the third part.
  \end{parts}
\end{physicsproblem}
```

Problem 2

This is a physics problem with multiple parts. The list is vertical.

- (a) This is the first part.
- (b) This is the second part.
- (c) This is the third part.


```

\begin{physicsproblem*}{Problem 3}
  This is a physics problem with multiple parts.
  The list is in-line.
  \begin{parts}
    \problempart This is the first part.
    \problempart This is the second part.
    \problempart This is the third part.
  \end{parts}
\end{physicsproblem*}

```

Problem 3

This is a physics problem with multiple parts. The list is in-line. **(a)** This is the first part. **(b)** This is the second part. **(c)** This is the third part.

U 2021-02-26

```

\begin{physicssolution}
  \langle solution steps \rangle

```

(use this variant for numbered steps)

U 2021-02-26

```

\begin{physicssolution*}
  \langle solution steps \rangle
\end{physicssolution*}

```

(use this variant for unnumbered steps)

This environment is only for mathematical solutions. The starred variant omits numbering of steps. See the examples.

```

\begin{physicssolution}
  x &= y + z \\
  z &= x - y \\
  y &= x - z
\end{physicssolution}
\begin{physicssolution*}
  x &= y + z \\
  z &= x - y \\
  y &= x - z
\end{physicssolution*}

```

$$x = y + z \quad (1)$$

$$z = x - y \quad (2)$$

$$y = x - z \quad (3)$$

$$x = y + z$$

$$z = x - y$$

$$y = x - z$$

U 2021-02-26

```

\reason{\langle reason \rangle}

```

Provides an annotation in a step-by-step solution. Keep reasons short and to the point. Wrap mathematical content in math mode.

```

\begin{physicssolution}
  x &= y + z \reason{This is a reason.} \\
  z &= x - y \reason{This is a reason too.} \\
  y &= x - z \reason{final answer}
\end{physicssolution}
\begin{physicssolution*}
  x &= y + z \reason{This is a reason.} \\
  z &= x - y \reason{This is a reason too.} \\
  y &= x - z \reason{final answer}
\end{physicssolution*}

```

$$x = y + z \quad \text{This is a reason.} \quad (4)$$

$$z = x - y \quad \text{This is a reason too.} \quad (5)$$

$$y = x - z \quad \text{final answer} \quad (6)$$

$$x = y + z \quad \text{This is a reason.}$$

$$z = x - y \quad \text{This is a reason too.}$$

$$y = x - z \quad \text{final answer}$$

When writing solutions, remember that the `physicssolution`^{P.57} environment is *only* for mathematical content, not textual content or explanations.

```

\begin{physicsproblem}{Combined Problem and Solution}
  This is an interesting physics problem.
  \begin{physicssolution}
    The solution goes here.
  \end{physicssolution}
\end{physicsproblem}

```

```

\begin{physicsproblem}{Combined Multipart Problem with Solutions}
  This is a physics problem with multiple parts.
  \begin{parts}
    \problempart This is the first part.
    \begin{physicssolution}
      The solution goes here.
    \end{physicssolution}
    \problempart This is the second part.
    \begin{physicssolution}
      The solution goes here.
    \end{physicssolution}
    \problempart This is the third part.
    \begin{physicssolution}
      The solution goes here.
    \end{physicssolution}
  \end{parts}
\end{physicsproblem}

```

Hilites the desired target, which can be an entire mathematical expression or a part thereof. The default color is magenta and the default shape is a rectangle. This command is not defined if BEAMER is loaded.

```

\begin{align*}
(\Delta s)^2 &= -(\Delta t)^2 + (\Delta x)^2 + (\Delta y)^2 + \\
&\quad (\Delta z)^2 \\\
(\Delta s)^2 &= \hilite{-(\Delta t)^2 + (\Delta x)^2}[rounded rectangle] + \\
&\quad (\Delta y)^2 + (\Delta z)^2 \\\
(\Delta s)^2 &= \hilite{-(\Delta t)^2 + (\Delta x)^2}[rectangle] + \\
&\quad (\Delta y)^2 + (\Delta z)^2 \\\
(\Delta s)^2 &= \hilite{-(\Delta t)^2 + (\Delta x)^2}[ellipse] + \\
&\quad (\Delta y)^2 + (\Delta z)^2 \\\
(\Delta s)^{\textcolor{black}{2}} &= \hilite[green]{-}[circle] \\
&\quad (\Delta t)^{\textcolor{cyan}{2}}[circle] + \\
&\quad (\Delta x)^{\textcolor{orange}{2}}[circle] + \\
&\quad (\Delta y)^{\textcolor{blue!50}{2}}[circle] + \\
&\quad (\Delta z)^{\textcolor{violet!45}{2}}[circle]
\end{align*}

```

$$\begin{aligned}
 (\Delta s)^2 &= -(\Delta t)^2 + (\Delta x)^2 + (\Delta y)^2 + (\Delta z)^2 \\
 (\Delta s)^2 &= \textcolor{pink}{-(\Delta t)^2 + (\Delta x)^2} + (\Delta y)^2 + (\Delta z)^2 \\
 (\Delta s)^2 &= \textcolor{pink}{-(\Delta t)^2 + (\Delta x)^2} + (\Delta y)^2 + (\Delta z)^2 \\
 (\Delta s)^2 &= \textcolor{pink}{-(\Delta t)^2 + (\Delta x)^2} + \textcolor{pink}{(\Delta y)^2} + (\Delta z)^2 \\
 (\Delta s)^{\textcolor{pink}{2}} &= \textcolor{green}{-}(\Delta t)^{\textcolor{cyan}{2}} + (\Delta x)^{\textcolor{orange}{2}} + (\Delta y)^{\textcolor{blue}{2}} + (\Delta z)^{\textcolor{violet}{2}}
 \end{aligned}$$

```

\begin{align*}
\Delta\vec{p} &= \vec{F}_{\text{sumup}\{net\}}\Delta t \\\
\textcolor{orange}{\Delta\vec{p}} &= \vec{F}_{\text{symup}\{net\}}\Delta t \\\
\Delta\vec{p} &= \hilite[yellow!50]{\vec{F}_{\text{symup}\{net\}}} \\
&\quad [rounded rectangle]\Delta t \\\
\Delta\vec{p} &= \vec{F}_{\text{symup}\{net\}}\hilite[olive!50] \\
&\quad \{\Delta t\}[rectangle] \\\
\Delta\vec{p} &= \hilite[cyan!50]{\vec{F}_{\text{symup}\{net\}}}\Delta t \\
&\quad [ellipse] \\\
\hilite{\Delta\vec{p}}[rectangle] &= \vec{F}_{\text{symup}\{net\}}\Delta t
\end{align*}

```

$$\begin{aligned}
 \Delta p &= F_{\text{net}} \Delta t \\
 \textcolor{orange}{\Delta p} &= F_{\text{net}} \Delta t \\
 \Delta p &= \textcolor{yellow}{F_{\text{net}}} \Delta t \\
 \Delta p &= F_{\text{net}} \textcolor{olive}{\Delta t} \\
 \Delta p &= \textcolor{cyan}{F_{\text{net}}} \Delta t \\
 \textcolor{pink}{\Delta p} &= F_{\text{net}} \Delta t
 \end{aligned}$$

`\image[<options>]{<caption>}{<label>}{<image>}`

Simplified interface for importing an image. The images are treated as floats, so they may not appear at the most logical intuitive place.

```
\image[scale=0.20]{example-image-1x1}
{Image shown 20 percent actual size.}{reffig1}
```



Figure 1: Image shown 20 percent actual size.

Figure \ref{reffig1} is nice.
It's captioned \nameref{reffig1} and is on page \pageref{reffig1}.

Figure 1 is nice. It's captioned Image shown 20 percent actual size and is on page 60.

```
\image[scale=0.20,angle=45]{example-image-1x1}
{Image shown 20 percent actual size and rotated.}{reffig1}
```




Figure 2: Image shown 20 percent actual size and rotated.

Figure \ref{reffig2} is nice.
It's captioned \nameref{reffig2} and is on page \pageref{reffig2}.

Figure 2 is nice. It's captioned Image shown 20 percent actual size and rotated and is on page 60.

7.3 Coordinate-Free and Index Notation

Beyond the current level of introductory physics, we need intelligent commands for typesetting vector and tensor symbols and components suitable for both coordinate-free and index notations.

```
\colvec[⟨delimiter⟩]{⟨c1, ..., cn⟩}
\rowvec[⟨delimiter⟩]{⟨c1, ..., cn⟩}
```

Typesets column vectors and row vectors as numeric or symbolic components. There can be more than three components. The delimiter used in the list of components can be specified; the default is a comma. Units are not supported, so these are mainly for symbolic work.

```
\[ \colvec{1,2,3} \]
\[ \rowvec{1,2,3} \]
\[ \colvec{x^0,x^1,x^2,x^3} \]
\[ \rowvec{x_0,x_1,x_2,x_3} \]
```

$$\begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}$$

$$(1 \ 2 \ 3)$$

$$\begin{pmatrix} x^0 \\ x^1 \\ x^2 \\ x^3 \end{pmatrix}$$

$$(x_0 \ x_1 \ x_2 \ x_3)$$

`\veccomp{symbol}`

(use this variant for coordinate-free vector notation)

`\veccomp*{symbol}`

(use this variant for index vector notation)

`\tencomp{symbol}`

(use this variant for coordinate-free tensor notation)

`\tencomp*{symbol}`

(use this variant for index tensor notation)

Conforms to ISO 80000-2 notation.

```
\( \veccomp{r} \) \\\
\(\ \veccomp*{r} \) \\\
\(\ \tencomp{r} \) \\\
\(\ \tencomp*{r} \) \\\
```

\boldsymbol{r}

r

\boldsymbol{r}

r

`\valence{index}{index}`

`\valence*{index}{index}`

Typesets tensor valence. The starred variant typesets it horizontally.

```
A vector is a \( \valence{1}{0} \) tensor. \\\
A vector is a \( \valence*{1}{0} \) tensor.
```

A vector is a $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ tensor.

A vector is a $(1,0)$ tensor.

`\contraction{slot,slot}`

`\contraction*{slot,slot}`

Typesets tensor contraction in coordinate-free notation. There is no standard for this so we assert one here.

```
\( \contraction{1,2} \) \\\
\(\ \contraction*{1,2} \)
```

$\mathbb{C}_{1,2}$

$C_{1,2}$

`\slot[vector]`

`\slot*[vector]`

An intelligent slot command for coordinate-free vector and tensor notation. The starred variants suppress the underscore.

<code>\(\slot) \)</code>	<code>\)</code>	(\quad)
<code>\(\slot[\vec{a}]) \)</code>	<code>\)</code>	(\underline{a})
<code>\(\slot*) \)</code>	<code>\)</code>	(\quad)
<code>\(\slot*[\vec{a}]) \)</code>	<code>\)</code>	(\underline{a})

U 2022-01-27

\df

Intelligent differential and exterior derivative operator.

<code>\[</code>		$\int x \, dx$
<code>\int x \, dx</code>		
<code>\]</code>		
<code>\[</code>		$\int x \, dx$
<code>\int x \, \df{x}</code>		
<code>\]</code>		
<code>\[</code>		$\int x \, \mathbf{d}x$
<code>\int x \, \df{*x}</code>		
<code>\]</code>		

7.4 Web VPython and VPython Program Listings

Web VPython³ and VPython⁴ are programming environments (both use Python) frequently used in introductory physics to introduce students for modeling physical systems. **mandi** makes including code listings very simple for students.

7.5 The webvpythonblock Environment

U 2023-08-01

`\begin{webvpythonblock}[\textit{options}](\textit{link}){\textit{caption}}` (now includes a QR code)
`\end{webvpythonblock}`

U 2023-08-01

`\begin{webvpythonblock*}[\textit{options}](\textit{link}){\textit{caption}}` (use this variant to omit QR code)
`\end{webvpythonblock*}`

Code placed here is nicely formatted and optionally linked to its source on [WebVPython.org](https://webvpython.org), which must be in a public (not private) folder. Clicking anywhere in the code window (between the black horizontal bars) or on the URL will open the link in the default browser. A caption is mandatory, and a label is internally generated. The listing always begins on a new page. A URL shortening utility is recommended to keep the URL from getting unruly especially if it generates an overfull box error. For convenience, `https://` is automatically prepended to the URL and can be omitted. The `#` character in a URL should not cause problems. The default URL is that of the Web VPython home page.

³On November 9, 2021 GlowScript was renamed to Web VPython. The website was changed to <https://webvpython.org>.

⁴<https://vpython.org>

```

\begin{webvpythonblock}
(glowscript.org/#/user/heafnerj/folder/mandidemo/program/mandidemo)
{Example With QR Code}
Web VPython 3.2

scene.width = 400
scene.height = 760
# constants and data
g = 9.8          # m/s^2
mball = 0.03     # kg
Lo = 0.26        # m
ks = 1.8         # N/m
deltat = 0.01    # s

# objects (origin is at ceiling)
ceiling = box(pos=vector(0,0,0), length=0.2, height=0.01,
              width=0.2)
ball = sphere(pos=vector(0,-0.3,0),radius=0.025,
              color=color.orange)
spring = helix(pos=ceiling.pos, axis=ball.pos-ceiling.pos,
               color=color.cyan,thickness=0.003,coils=40,
               radius=0.010)

# initial values
pball = mball * vector(0,0,0)      # kg m/s
Fgrav = mball * g * vector(0,-1,0) # N
t = 0

# improve the display
scene.autoscale = False           # turn off automatic camera zoom
scene.center = vector(0,-Lo,0)    # move camera down
scene.waitfor('click')           # wait for a mouse click

# initial calculation loop
# calculation loop
while t < 10:
    rate(100)
    # we need the stretch
    s = mag(ball.pos) - Lo
    # we need the spring force
    Fspring = ks * s * -norm(spring.axis)
    Fnet = Fgrav + Fspring
    pball = pball + Fnet * deltat
    ball.pos = ball.pos + (pball / mball) * deltat
    spring.axis = ball.pos - ceiling.pos
    t = t + deltat
\end{webvpythonblock}

```



Web VPython Program 1: Example With QR Code

<https://glowsript.org/#/user/heafnerj/folder/mandidemo/program/mandidemo>

```
1 Web VPython 3.2
2
3 scene.width = 400
4 scene.height = 760
5 # constants and data
6 g = 9.8 # m/s^2
7 mball = 0.03 # kg
8 Lo = 0.26 # m
9 ks = 1.8 # N/m
10 deltat = 0.01 # s
11
12 # objects (origin is at ceiling)
13 ceiling = box(pos=vector(0,0,0), length=0.2, height=0.01,
14               width=0.2)
15 ball = sphere(pos=vector(0,-0.3,0), radius=0.025,
16               color=color.orange)
17 spring = helix(pos=ceiling.pos, axis=ball.pos-ceiling.pos,
18                color=color.cyan, thickness=0.003, coils=40,
19                radius=0.010)
20
21 # initial values
22 pball = mball * vector(0,0,0) # kg m/s
23 Fgrav = mball * g * vector(0,-1,0) # N
24 t = 0
25
26 # improve the display
27 scene.autoscale = False # turn off automatic camera zoom
28 scene.center = vector(0,-Lo,0) # move camera down
29 scene.waitfor('click') # wait for a mouse click
30
31 # initial calculation loop
32 # calculation loop
33 while t < 10:
34     rate(100)
35     # we need the stretch
36     s = mag(ball.pos) - Lo
37     # we need the spring force
38     Fspring = ks * s * -norm(spring.axis)
39     Fnet = Fgrav + Fspring
40     pball = pball + Fnet * deltat
41     ball.pos = ball.pos + (pball / mball) * deltat
42     spring.axis = ball.pos - ceiling.pos
43     t = t + deltat
```


Here is how one would reference this program elsewhere. Notice the references are numbered sequentially within the document.

```
\WebVPython{} program \ref{gs:1} is nice.  
It's called \nameref{gs:1} and is on page \pageref{gs:1}.
```

Web VPython program [1](#) is nice. It's called [Example With QR Code](#) and is on page [64](#).

```

\begin{webvpythonblock*}
(glowscript.org/#/user/heafnerj/folder/mandidemo/program/mandidemo)
{Example Without QR Code}
Web VPython 3.2

scene.width = 400
scene.height = 760
# constants and data
g = 9.8          # m/s^2
mball = 0.03     # kg
Lo = 0.26        # m
ks = 1.8         # N/m
deltat = 0.01   # s

# objects (origin is at ceiling)
ceiling = box(pos=vector(0,0,0), length=0.2, height=0.01,
              width=0.2)
ball = sphere(pos=vector(0,-0.3,0),radius=0.025,
              color=color.orange)
spring = helix(pos=ceiling.pos, axis=ball.pos-ceiling.pos,
               color=color.cyan,thickness=0.003,coils=40,
               radius=0.010)

# initial values
pball = mball * vector(0,0,0)      # kg m/s
Fgrav = mball * g * vector(0,-1,0) # N
t = 0

# improve the display
scene.autoscale = False           # turn off automatic camera zoom
scene.center = vector(0,-Lo,0)    # move camera down
scene.waitfor('click')           # wait for a mouse click

# initial calculation loop
# calculation loop
while t < 10:
    rate(100)
    # we need the stretch
    s = mag(ball.pos) - Lo
    # we need the spring force
    Fspring = ks * s * -norm(spring.axis)
    Fnet = Fgrav + Fspring
    pball = pball + Fnet * deltat
    ball.pos = ball.pos + (pball / mball) * deltat
    spring.axis = ball.pos - ceiling.pos
    t = t + deltat
\end{webvpythonblock*}

```

Web VPython Program 2: Example Without QR Code

<https://glowsript.org/#/user/heafnerj/folder/mandidemo/program/mandidemo>

```
1 Web VPython 3.2
2
3 scene.width = 400
4 scene.height = 760
5 # constants and data
6 g = 9.8 # m/s^2
7 mball = 0.03 # kg
8 Lo = 0.26 # m
9 ks = 1.8 # N/m
10 deltat = 0.01 # s
11
12 # objects (origin is at ceiling)
13 ceiling = box(pos=vector(0,0,0), length=0.2, height=0.01,
14               width=0.2)
15 ball = sphere(pos=vector(0,-0.3,0), radius=0.025,
16               color=color.orange)
17 spring = helix(pos=ceiling.pos, axis=ball.pos-ceiling.pos,
18                color=color.cyan, thickness=0.003, coils=40,
19                radius=0.010)
20
21 # initial values
22 pball = mball * vector(0,0,0) # kg m/s
23 Fgrav = mball * g * vector(0,-1,0) # N
24 t = 0
25
26 # improve the display
27 scene.autoscale = False # turn off automatic camera zoom
28 scene.center = vector(0,-Lo,0) # move camera down
29 scene.waitfor('click') # wait for a mouse click
30
31 # initial calculation loop
32 # calculation loop
33 while t < 10:
34     rate(100)
35     # we need the stretch
36     s = mag(ball.pos) - Lo
37     # we need the spring force
38     Fspring = ks * s * -norm(spring.axis)
39     Fnet = Fgrav + Fspring
40     pball = pball + Fnet * deltat
41     ball.pos = ball.pos + (pball / mball) * deltat
42     spring.axis = ball.pos - ceiling.pos
43     t = t + deltat
```

\WebVPython{} program \ref{gs:2} is nice.
It's called \nameref{gs:2} and is on page \pageref{gs:2}.

Web VPython program 2 is nice. It's called [Example Without QR Code](#) and is on page 67.

7.6 The `vpythonfile` Command

U 2023-08-01

`\vpythonfile` [*<options>*] (<link>){<file>}{<caption>}

Command to load and typeset a VPython program, read from local file {<file>}. Clicking anywhere in the code window (between the black horizontal bars) or on the URL will open the link in the default browser. A caption is mandatory, and a label is internally generated. The listing always begins on a new page. A URL shortening utility is recommended to keep the URL from getting unruly especially if it generates an overfull box error. For convenience, `https://` is automatically prepended to the URL and can be omitted. The default URL is that of the VPython home page.

```
\vpythonfile{vdemo.py}{A \VPython{} Program}
```

VPython Program 1: A VPython Program

```

1  from vpython import *
2
3  scene.width = 400
4  scene.height = 760
5  # constants and data
6  g = 9.8      # m/s^2
7  mball = 0.03 # kg
8  Lo = 0.26    # m
9  ks = 1.8     # N/m
10 deltatt = 0.01 # s
11
12 # objects (origin is at ceiling)
13 ceiling = box(pos=vector(0,0,0), length=0.2, height=0.01,
14               width=0.2)
15 ball = sphere(pos=vector(0,-0.3,0), radius=0.025,
16               color=color.orange)
17 spring = helix(pos=ceiling.pos, axis=ball.pos-ceiling.pos,
18               color=color.cyan, thickness=0.003, coils=40,
19               radius=0.010)
20
21 # initial values
22 pball = mball * vector(0,0,0) # kg m/s
23 Fgrav = mball * g * vector(0,-1,0) # N
24 t = 0
25
26 # improve the display
27 scene.autoscale = False # turn off automatic camera zoom
28 scene.center = vector(0,-Lo,0) # move camera down
29 scene.waitfor('click') # wait for a mouse click
30
31 # initial calculation loop
32 # calculation loop
33 while t < 10:
34     rate(100)
35     # we need the stretch
36     s = mag(ball.pos) - Lo
37     # we need the spring force
38     Fspring = ks * s * -norm(spring.axis)
39     Fnet = Fgrav + Fspring
40     pball = pball + Fnet * deltatt
41     ball.pos = ball.pos + (pball / mball) * deltatt
42     spring.axis = ball.pos - ceiling.pos
43     t = t + deltatt

```

\VPython{} program \ref{vp:1} is nice.
 It's called \nameref{vp:1} and is on page \pageref{vp:1}.

VPython program 1 is nice. It's called [A VPython Program](#) and is on page 69.

7.7 The `webvpythoninline` and `vpythoninline` Commands

U 2021-02-26

`\webvpythoninline{<Web VPython code>}`

U 2021-02-26

`\vpythoninline{<VPython code>}`

Typesets a small, in-line snippet of code. The snippet should be less than one line long.

`\WebVPython{}` programs begin with `\webvpythoninline{Web VPython 3.2}`.

Web VPython programs begin with `Web VPython 3.2`.

`\VPython{}` programs begin with `\vpythoninline{from vpython import *}`.

VPython programs begin with `from vpython import *`.

7.8 mandistudent Source Code

Define the package version and date for global use, exploiting the fact that in a .sty file there is now no need for `\makeatletter` and `\makeatother`. This simplifies defining internal commands, with @ in the name, that are not for the user to know about.

```
1 \def\mandistudent@version{3.2.2a}
2 \def\mandistudent@date{2024-09-12}
3 \NeedsTeXFormat{LaTeX2e}[2020-02-02]
4 \DeclareRelease{v3.2.2a}{2024-09-12}{mandistudent.sty}
5 \DeclareCurrentRelease{v\mandistudent@version}{\mandistudent@date}
6 \ProvidesPackage{mandistudent}
7 [\mandistudent@date\space v\mandistudent@version\space Macros for introductory physics]

Define a convenient package version command.

8 \newcommand*{\mandistudentversion}{v\mandistudent@version\space dated \mandistudent@date}

Load third party packages, documenting why each one is needed.
AMS goodness. Don't load amssymb or amsfonts.

9 \RequirePackage{amsmath}

We need enumitem for the physicsproblem environment.
BEAMER is not compatible with enumitem so if BEAMER is loaded certain commands are not defined.

10 \IfClassLoadedTF{beamer}%
11 {}%
12 {%
13     \RequirePackage[inline]{enumitem}%
14 }%

We need eso-pic for \hilite.

15 \RequirePackage{eso-pic}

We need esvect for nice vector arrows, style g.

16 \RequirePackage[g]{esvect}

We need pgfopts for a key-value interface.

17 \RequirePackage{pgfopts}

We need iftex so we can require LuaLATEX.

18 \RequirePackage{iftex}

We need makebox for consistent \dirvect notation.

19 \RequirePackage{makebox}

We need mandi to load mathtools and unicode-math.

20 \IfPackageLoadedTF{mandi}%
21 {}%
22 {%
23     \RequirePackage{mandi}%
24 }%

We need nicematrix for column and row vectors.

25 \RequirePackage{nicematrix}

We need qrcode for QR codes in webvpythonblock.

26 \RequirePackage{qrcode}

Set the default size for QR codes.

27 \qrcode{height=1.5cm}

We need tcolorbox for program listings.

28 \RequirePackage[most]{tcolorbox}
```

We need tensor for index notation.

```
29 \RequirePackage{tensor}
30 %
31 % We need \pkg{tikz} for |\hilight|..
32 %
33 \RequirePackage{tikz}
34 \usetikzlibrary{shapes,fit,tikzmark}
```

Load xparse if necessary.

```
35 \IfFormatAtLeastTF{2020-10-01}
36 {}%
37 {\RequirePackage{xparse}}%
```

Always load hyperref last if possible.

```
38 \RequirePackage{hyperref}
```

We require the Lua \LaTeX engine.

```
39 \RequireLuaTeX
```

Set up the fonts to be consistent with ISO 80000-2 notation. The `unicode-math` package loads the `fontspec` and `xparse` packages. Note that `xparse` is now part of the \LaTeX 2_ε kernel. Because `unicode-math` is required, all documents using `mandi` must be compiled with an engine that supports Unicode, and I recommend Lua \LaTeX .

```
40 \unimathsetup{math-style=ISO}
```

Use normal math letters from Latin Modern Math for familiarity with textbooks. This gives a better J.

```
41 \setmathfont[Scale=MatchLowercase]
42 {Latin Modern Math}
```

Borrow from TeX Gyre DejaVu Math for vectors and tensors to get single-storey lowercase g.

```
43 \setmathfont[Scale=MatchLowercase,range={sfit/{latin},bfsfit/{latin}}]
44 {TeX Gyre DejaVu Math}
```

Borrow from TeX Gyre DejaVu Math to get single-storey lowercase g.

```
45 \setmathfont[Scale=MatchLowercase,range={sfup/{latin},bfsfup/{latin}}]
46 {TeX Gyre DejaVu Math}
```

Borrow `mathscr` and `mathbfscr` from XITS Math.

See <https://tex.stackexchange.com/a/120073/218142>.

```
47 \setmathfont[Scale=MatchLowercase,range={\mathscr,\mathbfscr}]{XITS Math}
```

Get original and bold `mathcal` fonts.

See <https://tex.stackexchange.com/a/21742/218142>.

```
48 \setmathfont[Scale=MatchLowercase,range={\mathcal,\mathbfcal},StylisticSet=1]{XITS Math}
```

Borrow Greek `sfup` and `sfit` letters from STIX Two Math. Since this isn't officially supported in `unicode-math` we have to manually set this up.

```
49 \setmathfont[Scale=MatchLowercase,range={"E17C-"E1F6}]{STIX Two Math}
50 \newfontfamily{\symsfgreek}{STIX Two Math}
```

I don't understand why `\text{...}` is necessary.

```
51 \newcommand{\symsfupalpha} {\text{\symsfgreek{~~~~e196}}}
52 \newcommand{\symsfupbeta} {\text{\symsfgreek{~~~~e197}}}
53 \newcommand{\symsfupgamma} {\text{\symsfgreek{~~~~e198}}}
54 \newcommand{\symsfupdelta} {\text{\symsfgreek{~~~~e199}}}
55 \newcommand{\symsfupepsilon} {\text{\symsfgreek{~~~~e1af}}}
56 \newcommand{\symsfupvarepsilon} {\text{\symsfgreek{~~~~e19a}}}
57 \newcommand{\symsfupzeta} {\text{\symsfgreek{~~~~e19b}}}
58 \newcommand{\symsfupeta} {\text{\symsfgreek{~~~~e19c}}}
```


59 \newcommand{\symsfuptheta}	{\text{\symsfgreek{~~~~e19d}}}
60 \newcommand{\symsfupvartheta}	{\text{\symsfgreek{~~~~e1b0}}}
61 \newcommand{\symsfupiota}	{\text{\symsfgreek{~~~~e19e}}}
62 \newcommand{\symsfupkappa}	{\text{\symsfgreek{~~~~e19f}}}
63 \newcommand{\symsfuplambda}	{\text{\symsfgreek{~~~~e1a0}}}
64 \newcommand{\symsfupmu}	{\text{\symsfgreek{~~~~e1a1}}}
65 \newcommand{\symsfupnu}	{\text{\symsfgreek{~~~~e1a2}}}
66 \newcommand{\symsfupxi}	{\text{\symsfgreek{~~~~e1a3}}}
67 \newcommand{\symsfupomicron}	{\text{\symsfgreek{~~~~e1a4}}}
68 \newcommand{\symsfuppi}	{\text{\symsfgreek{~~~~e1a5}}}
69 \newcommand{\symsfupvarpi}	{\text{\symsfgreek{~~~~e1b3}}}
70 \newcommand{\symsfuprho}	{\text{\symsfgreek{~~~~e1a6}}}
71 \newcommand{\symsfupvarrho}	{\text{\symsfgreek{~~~~e1b2}}}
72 \newcommand{\symsfupsigma}	{\text{\symsfgreek{~~~~e1a8}}}
73 \newcommand{\symsfupvarsigma}	{\text{\symsfgreek{~~~~e1a7}}}
74 \newcommand{\symsfuptau}	{\text{\symsfgreek{~~~~e1a9}}}
75 \newcommand{\symsfupupsilon}	{\text{\symsfgreek{~~~~e1aa}}}
76 \newcommand{\symsfupphi}	{\text{\symsfgreek{~~~~e1b1}}}
77 \newcommand{\symsfupvarphi}	{\text{\symsfgreek{~~~~e1ab}}}
78 \newcommand{\symsfupchi}	{\text{\symsfgreek{~~~~e1ac}}}
79 \newcommand{\symsfuppsi}	{\text{\symsfgreek{~~~~e1ad}}}
80 \newcommand{\symsfupomega}	{\text{\symsfgreek{~~~~e1ae}}}
81 \newcommand{\symsfupDelta}	{\text{\symsfgreek{~~~~e180}}}
82 \newcommand{\symsfupGamma}	{\text{\symsfgreek{~~~~e17f}}}
83 \newcommand{\symsfupTheta}	{\text{\symsfgreek{~~~~e18e}}}
84 \newcommand{\symsfupLambda}	{\text{\symsfgreek{~~~~e187}}}
85 \newcommand{\symsfupXi}	{\text{\symsfgreek{~~~~e18a}}}
86 \newcommand{\symsfupPi}	{\text{\symsfgreek{~~~~e18c}}}
87 \newcommand{\symsfupSigma}	{\text{\symsfgreek{~~~~e18f}}}
88 \newcommand{\symsfupUpsilon}	{\text{\symsfgreek{~~~~e191}}}
89 \newcommand{\symsfupPhi}	{\text{\symsfgreek{~~~~e192}}}
90 \newcommand{\symsfupPsi}	{\text{\symsfgreek{~~~~e194}}}
91 \newcommand{\symsfupOmega}	{\text{\symsfgreek{~~~~e195}}}
92 \newcommand{\symsfitalpha}	{\text{\symsfgreek{~~~~e1d8}}}
93 \newcommand{\symsfitbeta}	{\text{\symsfgreek{~~~~e1d9}}}
94 \newcommand{\symsfitgamma}	{\text{\symsfgreek{~~~~e1da}}}
95 \newcommand{\symsfitdelta}	{\text{\symsfgreek{~~~~e1db}}}
96 \newcommand{\symsfitepsilon}	{\text{\symsfgreek{~~~~e1f1}}}
97 \newcommand{\symsfitvarepsilon}	{\text{\symsfgreek{~~~~e1dc}}}
98 \newcommand{\symsfitzeta}	{\text{\symsfgreek{~~~~e1dd}}}
99 \newcommand{\symsfiteta}	{\text{\symsfgreek{~~~~e1de}}}
100 \newcommand{\symsfittheta}	{\text{\symsfgreek{~~~~e1df}}}
101 \newcommand{\symsfitvartheta}	{\text{\symsfgreek{~~~~e1f2}}}
102 \newcommand{\symsfitiota}	{\text{\symsfgreek{~~~~e1e0}}}
103 \newcommand{\symsfitkappa}	{\text{\symsfgreek{~~~~e1e1}}}
104 \newcommand{\symsfitlambda}	{\text{\symsfgreek{~~~~e1e2}}}
105 \newcommand{\symsfitmu}	{\text{\symsfgreek{~~~~e1e3}}}
106 \newcommand{\symsfitnu}	{\text{\symsfgreek{~~~~e1e4}}}
107 \newcommand{\symsfitxi}	{\text{\symsfgreek{~~~~e1e5}}}
108 \newcommand{\symsfitomicron}	{\text{\symsfgreek{~~~~e1e6}}}
109 \newcommand{\symsfitpi}	{\text{\symsfgreek{~~~~e1e7}}}
110 \newcommand{\symsfitvarpi}	{\text{\symsfgreek{~~~~e1f5}}}
111 \newcommand{\symsfitrho}	{\text{\symsfgreek{~~~~e1e8}}}
112 \newcommand{\symsfitvarrho}	{\text{\symsfgreek{~~~~e1f4}}}
113 \newcommand{\symsfitsigma}	{\text{\symsfgreek{~~~~e1ea}}}
114 \newcommand{\symsfitvarsigma}	{\text{\symsfgreek{~~~~e1e9}}}
115 \newcommand{\symsfittau}	{\text{\symsfgreek{~~~~e1eb}}}
116 \newcommand{\symsfitupsilon}	{\text{\symsfgreek{~~~~e1ec}}}
117 \newcommand{\symsfitphi}	{\text{\symsfgreek{~~~~e1f3}}}

```

118 \newcommand{\symsfitvarphi} {\text{\symsfgreek{~~~~e1ed}}}
119 \newcommand{\symsfitchi} {\text{\symsfgreek{~~~~e1ee}}}
120 \newcommand{\symsfitpsi} {\text{\symsfgreek{~~~~e1ef}}}
121 \newcommand{\symsfitomega} {\text{\symsfgreek{~~~~e1f0}}}
122 \newcommand{\symsfitDelta} {\text{\symsfgreek{~~~~e1c2}}}
123 \newcommand{\symsfitGamma} {\text{\symsfgreek{~~~~e1c1}}}
124 \newcommand{\symsfitTheta} {\text{\symsfgreek{~~~~e1d0}}}
125 \newcommand{\symsfitLambda} {\text{\symsfgreek{~~~~e1c9}}}
126 \newcommand{\symsfitXi} {\text{\symsfgreek{~~~~e1cc}}}
127 \newcommand{\symsfitPi} {\text{\symsfgreek{~~~~e1ce}}}
128 \newcommand{\symsfitSigma} {\text{\symsfgreek{~~~~e1d1}}}
129 \newcommand{\symsfitUpsilon} {\text{\symsfgreek{~~~~e1d3}}}
130 \newcommand{\symsfitPhi} {\text{\symsfgreek{~~~~e1d4}}}
131 \newcommand{\symsfitPsi} {\text{\symsfgreek{~~~~e1d6}}}
132 \newcommand{\symsfitOmega} {\text{\symsfgreek{~~~~e1d7}}}

```

Tweak the `esvect` package fonts to get the correct font size.

See <https://tex.stackexchange.com/a/566676>.

```

133 \DeclareFontFamily{U}{esvect}{}
134 \DeclareFontShape{U}{esvect}{m}{n}{%
135   <-5.5> vect5
136   <5.5-6.5> vect6
137   <6.5-7.5> vect7
138   <7.5-8.5> vect8
139   <8.5-9.5> vect9
140   <9.5-> vect10
141 }{}%

```

Write a banner to the console showing the options in use.

```

142 \typeout{}%
143 \typeout{mandistudent: You are using mandistudent \mandistudentversion.}%
144 \typeout{mandistudent: This package requires LuaLaTeX.}%
145 \typeout{mandistudent: This package changes the default math font(s).}%
146 \typeout{mandistudent: This package redefines the \protect\vec\space command.}%
147 \IfClassLoadedTF{beamer}%
148   {%
149     \typeout{mandistudent: BEAMER detected. Certain commands will not be defined.}%
150   }%
151 {}%
152 \typeout{}%

```

A better, intelligent coordinate-free $\vec{^{52}}$ command. Note the use of the `e_{~}` type of optional argument. This accounts for much of the flexibility and power of this command. Also note the use of the \TeX primitives `\sb{}` and `\sp{}`. Why doesn't it work when I put spaces around #3 or #4? Because outside of `\ExplSyntaxOn... \ExplSyntaxOff`, the `_` character has a different catcode and is treated as a mathematical entity.

See <https://tex.stackexchange.com/q/554706/218142>.

See also <https://tex.stackexchange.com/a/531037/218142>.

```

153 \RenewDocumentCommand{\vec}{ s m e_{~} }%
154   {%
155     \IfBooleanTF{#1}
156       {%
157         \vv{#2}%
158         \IfValueT{#4}%
159           {\sp{~, #4\vpantom{\smash[t]{\big|}}}}
160       }%
161     {%
162       \ymbfit{#2}
163       \IfValueT{#4}%
164         {\sp{#4\vpantom{\smash[t]{\big|}}}}

```

```

165     }%
166     \IfValueT{#3}%
167     {\sb{#3\vphantom{\smash[b]{|}}}}
168 }%

```

A command for the direction of a vector. We use a slight tweak to get uniform hats that requires the [makebox](https://tex.stackexchange.com/a/391204/218142) package. See <https://tex.stackexchange.com/a/391204/218142>.

```

169 \NewDocumentCommand{\dirvec}{ s m e_{~} }%
170 {%
171     \widehat%
172     {%
173         \makebox*{\(w\)}%
174         {%
175             \ensuremath{%
176                 \IfBooleanTF {#1}%
177                 {%
178                     #2%
179                 }%
180                 {%
181                     \symbf{#2}%
182                 }%
183             }%
184         }%
185     }%
186     \IfValueT{#3}%
187     {\sb{#3\vphantom{\smash[b]{|}}}}%
188     \IfValueT{#4}%
189     {\sp{\,#4\vphantom{\smash[t]{\big|}}}}%
190 }%

```

The zero vector.

```

191 \NewDocumentCommand{\zerovec}{ s }%
192 {%
193     \IfBooleanTF {#1}
194     {\vv{0}}%
195     {\symbfup{0}}%
196 }%

```

Notation for column and row vectors.

See <https://tex.stackexchange.com/a/39054/218142>.

```

197 \ExplSyntaxOn
198 \NewDocumentCommand{\colvec}{ O{,} m }
199 {
200     \__mandi_vectormain:nnnn { p } { \ } { #1 } { #2 }
201 }
202 \NewDocumentCommand{\rowvec}{ O{,} m }
203 {
204     \__mandi_vectormain:nnnn { p } { & } { #1 } { #2 }
205 }
206 \seq_new:N \l__mandi_vectorarg_seq
207 \cs_new_protected:Npn \__mandi_vectormain:nnnn #1#2#3#4
208 {
209     \seq_set_split:Nnn \l__mandi_vectorarg_seq { #3 } { #4 }
210     \begin{#1NiceMatrix}[r]
211         \seq_use:Nnnn \l__mandi_vectorarg_seq { #2 } { #2 } { #2 }
212     \end{#1NiceMatrix}
213 }
214 \ExplSyntaxOff

```

Students always need this symbol.

```
215 \NewDocumentCommand{\changein}{-}{\Delta}
```

Intelligent delimiters provided via the [mathtools](#) package. Use the starred variants for fractions. You can supply optional sizes. Note that default placeholders are used when the argument is empty.

```
216 \DeclarePairedDelimiterX{\doublebars}[1]{\lVert}{\rVert}{\ifblank{#1}{\:\cdot\:}{#1}}
217 \DeclarePairedDelimiterX{\singlebars}[1]{\lvert}{\rvert}{\ifblank{#1}{\:\cdot\:}{#1}}
218 \DeclarePairedDelimiterX{\anglebrackets}[1]{\langle}{\rangle}{\ifblank{#1}{\:\cdot\:}{#1}}
219 \DeclarePairedDelimiterX{\parentheses}[1]{\lparen}{\rparen}{\ifblank{#1}{\:\cdot\:}{#1}}
220 \DeclarePairedDelimiterX{\squarebrackets}[1]{\lbrack}{\rbrack}{\ifblank{#1}{\:\cdot\:}{#1}}
221 \DeclarePairedDelimiterX{\curlybraces}[1]{\lbrace}{\rbrace}{\ifblank{#1}{\:\cdot\:}{#1}}
```

Some semantic aliases. Because of the way $\vec{}$ ^{P.52} and $\vecdir{}$ ^{P.52} are defined, I reluctantly decided not to implement a \magvec command. It would require accounting for too many options. So $\mag{}$ ^{P.55} is the new solution.

```
222 \NewDocumentCommand{\magnitude}{-}{\doublebars}
223 \NewDocumentCommand{\norm}{-}{\doublebars}
224 \NewDocumentCommand{\absolutevalue}{-}{\singlebars}
```

Commands for two important geometric relationships. These are meant mainly to be subscripts.

```
225 \NewDocumentCommand{\parallelto}{-}%
226 {%
227   \mkern3mu\vphantom{\perp}\vrule depth 0pt\mkern2mu\vrule depth 0pt\mkern3mu%
228 }%
229 \NewDocumentCommand{\perpendicularto}{-}{\perp}
```

An environment for problem statements. The starred variant gives in-line lists. These are not defined if BEAMER is loaded.

```
230 \IfClassLoadedTF{beamer}
231 {}%
232 {}%
233 \NewDocumentEnvironment{physicsproblem}{ m }%
234 {%
235   \newpage%
236   \section*{#1}%
237   \newlist{parts}{enumerate}{2}%
238   \setlist[parts]{label=\bfseries(\alph*)}%
239 }%
240 {}%
241 \NewDocumentEnvironment{physicsproblem*}{ m }%
242 {%
243   \newpage%
244   \section*{#1}%
245   \newlist{parts}{enumerate*}{2}%
246   \setlist[parts]{label=\bfseries(\alph*)}%
247 }%
248 {}%
249 \NewDocumentCommand{\problempart}{-}{\item}%
250 }%
```

An environment for problem solutions. Equation numbering is consecutive through the document.

```
251 \NewDocumentEnvironment{physicssolution}{ +b }%
252 {%
253   \begin{align}
254     #1
255   \end{align}
256 }%
257 {}%
258 \NewDocumentEnvironment{physicssolution*}{ +b }%
259 {%
260   \begin{align*}
```

```

261      #1
262      \end{align*}
263    }%
264  }%

```

See <https://tex.stackexchange.com/q/570223/218142>.

```

265 \NewDocumentCommand{\reason}{ 0{4cm} m }%
266   {%
267     &&\begin{minipage}{#1}\raggedright\small #2\end{minipage}%
268   }%

```

Command for highlighting parts of, or entire, mathematical expressions.

This command is not defined if BEAMER is loaded.

See <https://texample.net/tikz/examples/beamer-arrows/>.

See also <https://tex.stackexchange.com/a/406084/218142>.

See also <https://tex.stackexchange.com/a/570858/218142>.

See also <https://tex.stackexchange.com/a/570789/218142>.

See also <https://tex.stackexchange.com/a/79659/218142>.

See also <https://tex.stackexchange.com/q/375032/218142>.

See also <https://tex.stackexchange.com/a/571744/218142>

```

269 \newcounter{tikzhighlightnode}
270 \NewDocumentCommand{\hilite}{ 0{magenta!60} m 0{rectangle} }%
271   {%
272     \stepcounter{tikzhighlightnode}%
273     \tikzmarknode{highlighted-node-\number\value{tikzhighlightnode}}{#2}%
274     \edef\temp{%
275       \noexpand\AddToShipoutPictureBG{%
276         \noexpand\begin{tikzpicture}[overlay,remember picture]%
277           \noexpand\iftikzmarkconcurrentpage{highlighted-node-\number\value{tikzhighlightnode}}%
278             \noexpand\node[inner sep=1.0pt,fill=#1,#3,fit=(highlighted-node-\number\value{tikzhighlightnode})]{};%
279           \noexpand\fi
280         \noexpand\end{tikzpicture}%
281       }%
282     }%
283     \temp%
284   }%

```

A simplified command for importing images.

See <https://tex.stackexchange.com/a/614478/218142>.

```

285 \NewDocumentCommand{\image}{ 0{scale=1} m m m }%
286   {%
287     \par
288     \begin{figure}[ht!]
289       \centering%
290       \includegraphics[#1]{#2}%
291       \caption{#3}%
292       \label{#4}%
293     \end{figure}%
294     \par
295   }%

```

Intelligent commands for typesetting vector and tensor symbols and components suitable for use with both coordinate-free and index notations. Use starred form for index notation, unstarred form for coordinate-free.

Consider renaming these to `\vectorsym` and `\tensorsym`.

```

296 \NewDocumentCommand{\veccomp}{ s m }%
297   {%
298     \IfBooleanTF{#1}
299     {%

```

```

300      \symnormal{#2}%
301    }%
302    {%
303      \symsfit{#2}%
304    }%
305  }%
306 \NewDocumentCommand{\tencomp}{ s m }%
307  {%
308    \IfBooleanTF{#1}%
309    {%
310      \symsfit{#2}%
311    }%
312    {%
313      \symsfsfit{#2}%
314    }%
315  }%

```

Command to typeset tensor valence.

```

316 \NewDocumentCommand{\valence}{ s m m }%
317  {%
318    \IfBooleanTF{#1}%
319    {%
320      (#2,#3)%
321    }%
322    {%
323      \binom{#2}{#3}%
324    }%
325  }%

```

Intelligent notation for contraction on pairs of slots.

```

326 \NewDocumentCommand{\contraction}{ s m }%
327  {%
328    \IfBooleanTF{#1}%
329    {%
330      \mathsf{C}%
331    }%
332    {%
333      \symsb{C}%
334    }%
335    _{#2}%
336  }%

```

Intelligent slot command for coordinate-free tensor notation. `d[]` must be used because of the way consecutive optional arguments are handled. See `xparse` docs for details.

```

337 \NewDocumentCommand{\slot}{ s d[] }%
338  {%
339    \IfBooleanTF{#1}%
340    {%
341      \IfValueTF{#2}%

```

Insert a vector, but don't show the slot.

```

342    {%
343      \smash{\makebox[1.5em]{\ensuremath{#2}}}
344    }%

```

No vector, no slot.

```

345    {%
346      \smash{\makebox[1.5em]{\ensuremath{}}}
347    }%

```

```

348 }%
349 {%
350 \IfValueTF{#2}

Insert a vector and show the slot.

351 {%
352 \underline{\smash{\makebox[1.5em]{\ensuremath{#2}}}}
353 }%

```

No vector; just show the slot.

```

354 {%
355 \underline{\smash{\makebox[1.5em]{\ensuremath{}}}}
356 }%
357 }%
358 }%

```

Intelligent differential (exterior derivative) operator.

```

359 \NewDocumentCommand{\df}{ s }%
360 {%
361 \mathop{}\!%
362 \IfBooleanTF{#1}%
363 {%
364 \sybfsfup{d}%
365 }%
366 {%
367 \symsfup{d}%
368 }%
369 }%

```

Here is a clever way to color digits in program listings thanks to Ulrike Fischer.

See <https://tex.stackexchange.com/a/570717/218142>.

```

370 \directlua{%
371 luaotfload.add_colorscheme("colordigits",
372 [{"8000FF"} = {"one","two","three","four","five","six","seven","eight","nine","zero"}])
373 }%
374 \newfontfamily\colordigits{DejaVuSansMono}[RawFeature={color=colordigits}]

```

Set up a color scheme and a new code environment for listings. The new colors are more restful on the eye. All listing commands now use `tclobox`.

See <https://tex.stackexchange.com/a/529421/218142>.

We set a new font for listings and some new colors (background gray, gray, green, orange, peach, pearl, and plum).

```

375 \newfontfamily\gsfontfamily{DejaVuSansMono}
376 \definecolor{gsbggray} {rgb}{0.90,0.90,0.90}
377 \definecolor{gsgray} {rgb}{0.30,0.30,0.30}
378 \definecolor{gsgreen} {rgb}{0.00,0.60,0.00}
379 \definecolor{gsorange} {rgb}{0.80,0.45,0.12}
380 \definecolor{gspeach} {rgb}{1.00,0.90,0.71}
381 \definecolor{gspearl} {rgb}{0.94,0.92,0.84}
382 \definecolor{gsplum} {rgb}{0.74,0.46,0.70}
383 \lstdefinestyle{vpython}%
384 {%
385 backgroundcolor=\color{gsbggray},% % background color
386 basicstyle=\colordigits\footnotesize,% % default style
387 breakatwhitespace=true% % break at whitespace
388 breaklines=true,% % break long lines
389 captionpos=b,% % position caption
390 classoffset=1,% % STILL DON'T UNDERSTAND THIS
391 commentstyle=\color{gsgray},% % font for comments
392 deletekeywords={print},% % delete keywords from the given language

```

```

393   emph={self,cls,@classmethod,@property},%      % words to emphasize
394   emphstyle=\color{gsorange}\itshape,%          % font for emphasis
395   escapeinside={(*@){@*}},%                    % add LaTeX within your code
396   frame=tb,%                                    % frame style
397   framerule=2.0pt,%                             % frame thickness
398   framexleftmargin=5pt,%                        % extra frame left margin
399   %identifierstyle=\sffamily,%                   % style for identifiers
400   keywordstyle=\gsfontfamily\color{gspium},%    % color for keywords
401   language=Python,%                             % select language
402   linewidth=\linewidth,%                       % width of listings
403   morekeywords={%                               % VPython/Web VPython specific keywords
404     __future__,abs,acos,align,ambient,angle,append,append_to_caption,%
405     append_to_title,arange,arrow,asin,astuple,atan,atan2,attach_arrow,%
406     attach_trail,autoscale,axis,background,billboard,bind,black,blue,border,%
407     bounding_box,box,bumpaxis,bumpmap,bumpmaps,camera,canvas,caption,capture,%
408     ceil,center,clear,clear_trail,click,clone,CoffeeScript,coils,color,combin,%
409     comp,compound,cone,convex,cos,cross,curve,cyan,cylinder,data,degrees,del,%
410     delete,depth,descender,diff_angle,digits,division,dot,draw_complete,%
411     ellipsoid,emissive,end_face_color,equal,explog,extrusion,faces,factorial,%
412     False,floor,follow,font,format,forward,fov,frame,gcurve,gdisplay,gdots,%
413     get_library,get_selected,ghbars,global,GlowScript,graph,graphs,green,gvbars,%
414     hat,headlength,headwidth,height,helix,hsv_to_rgb,index,interval,keydown,%
415     keyup,label,length,lights,line,linecolor,linewidth,logx,logy,lower_left,%
416     lower_right,mag,mag2,magenta,make_trail,marker_color,markers,material,%
417     max,min,mouse,mousedown,mousemove,mouseup,newball,norm,normal,objects,%
418     offset,one,opacity,orange,origin,path,pause,pi,pixel_to_world,pixels,plot,%
419     points,pos,pow,pps,print,print_function,print_options,proj,purple,pyramid,%
420     quad,radians,radius,random,rate,ray,read_local_file,readonly,red,redraw,%
421     retain,rgb_to_hsv,ring,rotate,round,scene,scroll,shaftwidth,shape,shapes,%
422     shininess,show_end_face,show_start_face,sign,sin,size,size_units,sleep,%
423     smooth,space,sphere,sqrt,start,start_face_color,stop,tan,text,textpos,%
424     texture,textures,thickness,title,trail_color,trail_object,trail_radius,%
425     trail_type,triangle,trigger,True,twist,unbind,up,upper_left,upper_right,%
426     userpan,userspin,userzoom,vec,vector,vertex,vertical_spacing,visible,%
427     visual,vpython,VPython,waitfor,Web,VPython,white,width,world,xtitle,%
428     yellow,yoffset,ytitle%
429   },%
430   morekeywords={print,None,TypeError},%         % additional keywords
431   morestring=[b]{"""},%                        % treat triple quotes as strings
432   numbers=left,%                               % where to put line numbers
433   numbersep=10pt,%                             % how far line numbers are from code
434   numberstyle=\bfseries\tiny,%                 % set to 'none' for no line numbers
435   showstringspaces=false,%                     % show spaces in strings
436   showtabs=false,%                             % show tabs within strings
437   stringstyle=\gsfontfamily\color{gsgreen},%   % color for strings
438   upquote=true,%                               % how to typeset quotes
439 }%

```

Introduce a new, more intelligent [webvpythonblock](https://tex.stackexchange.com/a/232208/218142)^{→P.62} environment.

See <https://tex.stackexchange.com/a/232208/218142>.

```

440 \AtBeginEnvironment{webvpythonblock}{\catcode\#12}
441 \AtEndEnvironment{webvpythonblock}{\catcode\#6}
442 \NewTCBListing[auto counter,list inside=gspgros]{webvpythonblock}{0}{D(){webvpython.org} m }%
443 {%
444   breakable,%
445   center,%
446   code = \newpage,%
447   %derivpeach,%
448   enhanced,%

```



```

449   hyperurl interior = https://#2,%
450   label = {gs:\thetcbcounter},%
451   left = 8mm,%
452   list entry = \thetcbcounter~~~~#3,%
453   listing only,%
454   listing style = vpython,%
455   nameref = {#3},%
456   title = \begin{minipage}{1.5cm}%
457           \protect\qr{code*{https://#2}}%
458           \end{minipage}\hspace{5mm}%
459           \begin{minipage}{0.8\textwidth}%
460             \texttt{Web VPython} Program \thetcbcounter: #3\\
461             \footnotesize{\href{https://#2}{\color{white}{https://#2}}}%
462           \end{minipage},%
463   width = 0.9\textwidth,%
464   {#1},
465 }%

```

Here is a variant that omits the QR code.

```

466 \AtBeginEnvironment{webvpythonblock*}{\catcode`\#=12}
467 \AtEndEnvironment{webvpythonblock*}{\catcode`\#=6}
468 \NewTCBListing[use counter from=webvpythonblock,list inside=gsprogs]
469   {webvpythonblock*}{ 0{ } D(){webvpython.org} m }%
470   {%
471     breakable,%
472     center,%
473     code = \newpage,%
474     %derivpeach,%
475     enhanced,%
476     hyperurl interior = https://#2,%
477     label = {gs:\thetcbcounter},%
478     left = 8mm,%
479     list entry = \thetcbcounter~~~~#3,%
480     listing only,%
481     listing style = vpython,%
482     nameref = {#3},%
483     title = \texttt{Web VPython} Program \thetcbcounter: #3\\%
484             \footnotesize{\href{https://#2}{\color{white}{https://#2}}},%
485     width = 0.9\textwidth,%
486     {#1},
487   }%

```

A new command for generating a list of Web VPython programs.

```

488 \NewDocumentCommand{\listofwebvpythonprograms}{}%
489   {%
490     \tblistof[\section*]{gsprogs}{List of \texttt{Web VPython} Programs}%
491   }%

```

Introduce a new, more intelligent `\vpythonfile`^{P.68} command.
See <https://tex.stackexchange.com/q/616205/218142>.

```

492 \newcommand*{\vpythonfile}{\catcode`\#=12 \vpythonfile@auxA}
493 \NewDocumentCommand{\vpythonfile@auxA}{ 0{ } D(){vpython.org} m m }%
494   {%
495     \vpythonfile@auxB[#1](#2){#3}{#4}%
496     \catcode`\#=6
497   }%
498 \NewTCBInputListing[auto counter,list inside=vpvprogs]
499   {\vpythonfile@auxB}{ 0{ } D(){vpython.org} m m }%
500   {%

```

```

501     breakable,%
502     center,%
503     code = \newpage,%
504     %derivgray,%
505     enhanced,%
506     hyperurl interior = https://#2,%
507     label = {vp:\thetcbcounter},%
508     left = 8mm,%
509     list entry = \thetcbcounter~~~~#4,%
510     listing file = {#3},%
511     listing only,%
512     listing style = vpython,%
513     nameref = {#4},%
514     title = \texttt{VPython} Program \thetcbcounter: #4,%
515     width = 0.9\textwidth,%
516     {#1},%
517 }%

```

A new command for generating a list of VPython programs.

```

518 \NewDocumentCommand{\listofvpythonprograms}{}%
519 {%
520     \tblistof[\section*]{vpprogs}{List of \texttt{VPython} Programs}%
521 }%

```

Introduce a new `\webvpythoninline`^{P. 70} command.

```

522 \DeclareTotalTCBox{\webvpythoninline}{ m }%
523 {%
524     bottom = 0pt,%
525     bottomrule = 0.0mm,%
526     boxsep = 1.0mm,%
527     colback = gsbggray,%
528     colframe = gsbggray,%
529     left = 0pt,%
530     leftrule = 0.0mm,%
531     nobeforeafter,%
532     right = 0pt,%
533     rightrule = 0.0mm,%
534     sharp corners,%
535     tcbox raise base,%
536     top = 0pt,%
537     toprule = 0.0mm,%
538 }%
539 {\lstinline[style = vpython]{#1}}%

```

Define `\vpythoninline`^{P. 70}, a semantic alias for VPython in-line listings.

```

540 \NewDocumentCommand{\vpythoninline}{}{\webvpythoninline}%

```

8 The `mandiexp` Package

`mandi` comes with an accessory package `mandiexp` which extends `mandi` with commands specific to *Matter & Interactions*.⁵ The commands are primarily for typesetting mathematical expressions used in that text. `mandiexp` requires, and loads, `mandi` but `mandi` doesn't require, and doesn't load, `mandiexp`. `mandiexp` requires the `\vec*` command and so loads `mandistudent` if it has not already been loaded.

Load `mandiexp` as you would any package in your preamble. There are no package options.

```
\usepackage{mandiexp}
```

`\mandiexpversion`

Typesets the current version and build date.

The version is `\mandiexpversion{}` and is a stable build.

The version is v3.2.2a dated 2024-09-12 and is a stable build.

8.1 The Fundamental Principles

<code>\lhsmomentumprinciple</code>	(LHS of delta form, bold vectors)
<code>\rhsmomentumprinciple</code>	(RHS of delta form, bold vectors)
<code>\lhsmomentumprincipleupdate</code>	(LHS of update form, bold vectors)
<code>\rhsmomentumprincipleupdate</code>	(RHS of update form, bold vectors)
<code>\momentumprinciple</code>	(delta form, bold vectors)
<code>\momentumprincipleupdate</code>	(update form, bold vectors)
<code>\lhsmomentumprinciple*</code>	(LHS of delta form, arrow vectors)
<code>\rhsmomentumprinciple*</code>	(RHS of delta form, arrow vectors)
<code>\lhsmomentumprincipleupdate*</code>	(LHS of update form, arrow vectors)
<code>\rhsmomentumprincipleupdate*</code>	(RHS of update form, arrow vectors)
<code>\momentumprinciple*</code>	(delta form, arrow vectors)
<code>\momentumprincipleupdate*</code>	(update form, arrow vectors)

Variants of command for typesetting the momentum principle. Use starred variants to get arrow notation for vectors.

⁵See *Matter & Interactions* and <https://matterandinteractions.org/> for details.

```

\l \lhsmomentumprinciple \)      \l
\l \rhsmomentumprinciple \)      \l
\l \lhsmomentumprincipleupdate \) \l
\l \rhsmomentumprincipleupdate \) \l
\l \momentumprinciple \)        \l
\l \momentumprincipleupdate \)   \l
\l \lhsmomentumprinciple* \)     \l
\l \rhsmomentumprinciple* \)     \l
\l \lhsmomentumprincipleupdate* \) \l
\l \rhsmomentumprincipleupdate* \) \l
\l \momentumprinciple* \)        \l
\l \momentumprincipleupdate* \)  \l

```

$$\begin{aligned}
&\Delta p_{\text{sys}} \\
&\mathbf{F}_{\text{sys,net}} \Delta t \\
&p_{\text{sys,final}} \\
&p_{\text{sys,initial}} + \mathbf{F}_{\text{sys,net}} \Delta t \\
&\Delta p_{\text{sys}} = \mathbf{F}_{\text{sys,net}} \Delta t \\
&p_{\text{sys,final}} = p_{\text{sys,initial}} + \mathbf{F}_{\text{sys,net}} \Delta t \\
&\Delta \vec{p}_{\text{sys}} \\
&\vec{F}_{\text{sys,net}} \Delta t \\
&\vec{p}_{\text{sys,final}} \\
&\vec{p}_{\text{sys,initial}} + \vec{F}_{\text{sys,net}} \Delta t \\
&\Delta \vec{p}_{\text{sys}} = \vec{F}_{\text{sys,net}} \Delta t \\
&\vec{p}_{\text{sys,final}} = \vec{p}_{\text{sys,initial}} + \vec{F}_{\text{sys,net}} \Delta t
\end{aligned}$$

<code>\lhsenergyprinciple</code>	(LHS of delta form)
<code>\rhsenergyprinciple[⟨+process...⟩]</code>	(RHS of delta form)
<code>\lhsenergyprincipleupdate</code>	(LHS of update form)
<code>\rhsenergyprincipleupdate[⟨+process...⟩]</code>	(RHS of update form)
<code>\energyprinciple[⟨+process...⟩]</code>	(delta form)
<code>\energyprincipleupdate[⟨+process...⟩]</code>	(update form)

Variants of command for typesetting the energy principle.

```

\l \lhsenergyprinciple \)      \l
\l \rhsenergyprinciple \)      \l
\l \rhsenergyprinciple[+Q] \)  \l
\l \energyprinciple \)        \l
\l \energyprinciple[+Q] \)    \l
\l \lhsenergyprincipleupdate \) \l
\l \rhsenergyprincipleupdate \) \l
\l \rhsenergyprincipleupdate[+Q] \) \l
\l \energyprincipleupdate \)   \l
\l \energyprincipleupdate[+Q] \) \l

```

$$\begin{aligned}
&\Delta E_{\text{sys}} \\
&W_{\text{ext}} \\
&W_{\text{ext}} + Q \\
&\Delta E_{\text{sys}} = W_{\text{ext}} \\
&\Delta E_{\text{sys}} = W_{\text{ext}} + Q \\
&E_{\text{sys,final}} \\
&E_{\text{sys,initial}} + W_{\text{ext}} \\
&E_{\text{sys,initial}} + W_{\text{ext}} + Q \\
&E_{\text{sys,final}} = E_{\text{sys,initial}} + W_{\text{ext}} \\
&E_{\text{sys,final}} = E_{\text{sys,initial}} + W_{\text{ext}} + Q
\end{aligned}$$

<code>\lhsangularmomentumprinciple</code>	(LHS of delta form, bold vectors)
<code>\rhsangularmomentumprinciple</code>	(RHS of delta form, bold vectors)
<code>\lhsangularmomentumprincipleupdate</code>	(LHS of update form, bold vectors)
<code>\rhsangularmomentumprincipleupdate</code>	(RHS of update form, bold vectors)
<code>\angularmomentumprinciple</code>	(delta form, bold vectors)
<code>\angularmomentumprincipleupdate</code>	(update form, bold vectors)
<code>\lhsangularmomentumprinciple*</code>	(LHS of delta form, arrow vectors)
<code>\rhsangularmomentumprinciple*</code>	(RHS of delta form, arrow vectors)
<code>\lhsangularmomentumprincipleupdate*</code>	(LHS of update form, arrow vectors)
<code>\rhsangularmomentumprincipleupdate*</code>	(RHS of update form, arrow vectors)
<code>\angularmomentumprinciple*</code>	(delta form, arrow vectors)
<code>\angularmomentumprincipleupdate*</code>	(update form, arrow vectors)

Variants of command for typesetting the angular momentum principle. Use starred variants to get arrow notation for vectors.

<code>\(\lhsangularmomentumprinciple \)</code>	<code>\)</code>	$\Delta \mathbf{L}_{A,\text{sys},\text{net}}$
<code>\(\rhsangularmomentumprinciple \)</code>	<code>\)</code>	$\boldsymbol{\tau}_{A,\text{sys},\text{net}} \Delta t$
<code>\(\lhsangularmomentumprincipleupdate \)</code>	<code>\)</code>	$\mathbf{L}_{A,\text{sys},\text{final}}$
<code>\(\rhsangularmomentumprincipleupdate \)</code>	<code>\)</code>	$\mathbf{L}_{A,\text{sys},\text{initial}} + \boldsymbol{\tau}_{A,\text{sys},\text{net}} \Delta t$
<code>\(\angularmomentumprinciple \)</code>	<code>\)</code>	$\Delta \mathbf{L}_{A,\text{sys},\text{net}} = \boldsymbol{\tau}_{A,\text{sys},\text{net}} \Delta t$
<code>\(\angularmomentumprincipleupdate \)</code>	<code>\)</code>	$\mathbf{L}_{A,\text{sys},\text{final}} = \mathbf{L}_{A,\text{sys},\text{initial}} + \boldsymbol{\tau}_{A,\text{sys},\text{net}} \Delta t$
<code>\(\lhsangularmomentumprinciple* \)</code>	<code>\)</code>	$\Delta \vec{\mathbf{L}}_{A,\text{sys},\text{net}}$
<code>\(\rhsangularmomentumprinciple* \)</code>	<code>\)</code>	$\vec{\boldsymbol{\tau}}_{A,\text{sys},\text{net}} \Delta t$
<code>\(\lhsangularmomentumprincipleupdate* \)</code>	<code>\)</code>	$\vec{\mathbf{L}}_{A,\text{sys},\text{final}}$
<code>\(\rhsangularmomentumprincipleupdate* \)</code>	<code>\)</code>	$\vec{\mathbf{L}}_{A,\text{sys},\text{initial}} + \vec{\boldsymbol{\tau}}_{A,\text{sys},\text{net}} \Delta t$
<code>\(\angularmomentumprinciple* \)</code>	<code>\)</code>	$\Delta \vec{\mathbf{L}}_{A,\text{sys},\text{net}} = \vec{\boldsymbol{\tau}}_{A,\text{sys},\text{net}} \Delta t$
<code>\(\angularmomentumprincipleupdate* \)</code>	<code>\)</code>	$\vec{\mathbf{L}}_{A,\text{sys},\text{final}} = \vec{\mathbf{L}}_{A,\text{sys},\text{initial}} + \vec{\boldsymbol{\tau}}_{A,\text{sys},\text{net}} \Delta t$

8.2 Other Symbols

N 2021-02-13

`\energyof{<label>}[<label>]`

Generic symbol for the energy of some entity.

<code>\(\energyof{\symup{electron}} \)</code>	<code>\)</code>	E_{electron}
<code>\(\energyof{\symup{electron}}[\symup{final}] \)</code>	<code>\)</code>	$E_{\text{electron},\text{final}}$

N 2021-02-13

`\systemenergy[<label>]`

Symbol for system energy.

<code>\(\systemenergy \)</code>	<code>\)</code>	E_{sys}
<code>\(\systemenergy[\symup{final}] \)</code>	<code>\)</code>	$E_{\text{sys},\text{final}}$

N 2021-02-13

`\particleenergy[<label>]`

Symbol for particle energy.

<code>\(\particleenergy \)</code>	<code>\)</code>	E_{particle}
<code>\(\particleenergy[\symup{final}] \)</code>	<code>\)</code>	$E_{\text{particle},\text{final}}$

N 2021-02-13

`\restenergy[<label>]`

Symbol for rest energy.

<code>\(\restenergy \)</code>	<code>\)</code>	E_{rest}
<code>\(\restenergy[\symup{final}] \)</code>	<code>\)</code>	$E_{\text{rest},\text{final}}$

N 2021-02-13

`\internalenergy[⟨label⟩]`

Symbol for internal energy.

`\(\internalenergy \) \\
\(\internalenergy[\symup{final}] \)`

E_{internal}
 $E_{\text{internal,final}}$

N 2021-02-13

`\chemicalenergy[⟨label⟩]`

Symbol for chemical energy.

`\(\chemicalenergy \) \\
\(\chemicalenergy[\symup{final}] \)`

E_{chem}
 $E_{\text{chem,final}}$

N 2021-02-13

`\thermalenergy[⟨label⟩]`

Symbol for thermal energy.

`\(\thermalenergy \) \\
\(\thermalenergy[\symup{final}] \)`

E_{therm}
 $E_{\text{therm,final}}$

N 2021-02-13

`\photonenergy[⟨label⟩]`

Symbol for photon energy.

`\(\photonenergy \) \\
\(\photonenergy[\symup{final}] \)`

E_{photon}
 $E_{\text{photon,final}}$

N 2021-02-13

`\translationalkineticenergy[⟨label⟩]`

N 2021-02-13

`\translationalkineticenergy*[⟨label⟩]`

Symbol for translational kinetic energy. The starred variant gives E notation.

`\(\translationalkineticenergy \) \\
\(\translationalkineticenergy[\symup{initial}] \) \\
\(\translationalkineticenergy* \) \\
\(\translationalkineticenergy*[\symup{initial}] \)`

K_{trans}
 $K_{\text{trans,initial}}$
 E_K
 $E_{K,\text{initial}}$

N 2021-02-13

`\rotationalkineticenergy[⟨label⟩]`

N 2021-02-13

`\rotationalkineticenergy*[⟨label⟩]`

Symbol for rotational kinetic energy. The starred variant gives E notation.

`\(\rotationalkineticenergy \) \\
\(\rotationalkineticenergy[\symup{initial}] \) \\
\(\rotationalkineticenergy* \) \\
\(\rotationalkineticenergy*[\symup{initial}] \)`

K_{rot}
 $K_{\text{rot,initial}}$
 E_{rot}
 $E_{\text{rot,initial}}$

N 2021-02-13

`\vibrationalkineticenergy[⟨label⟩]`

N 2021-02-13

`\vibrationalkineticenergy*[⟨label⟩]`

Symbol for vibrational kinetic energy. The starred variant gives E notation.

<code>\(\vibrationalkineticenergy \)</code>	<code>\)</code>	K_{vib}
<code>\(\vibrationalkineticenergy[\text{symup{initial}}] \)</code>	<code>\)</code>	$K_{\text{vib,initial}}$
<code>\(\vibrationalkineticenergy* \)</code>	<code>\)</code>	E_{vib}
<code>\(\vibrationalkineticenergy*[\text{symup{initial}}] \)</code>	<code>\)</code>	$E_{\text{vib,initial}}$

N 2021-02-13

`\gravitationalpotentialenergy[⟨label⟩]`

Symbol for gravitational potential energy.

<code>\(\gravitationalpotentialenergy \)</code>	<code>\)</code>	U_{g}
<code>\(\gravitationalpotentialenergy[\text{symup{final}}] \)</code>	<code>\)</code>	$U_{\text{g,final}}$

N 2021-02-13

`\electricpotentialenergy[⟨label⟩]`

Symbol for electric potential energy.

<code>\(\electricpotentialenergy \)</code>	<code>\)</code>	U_{e}
<code>\(\electricpotentialenergy[\text{symup{final}}] \)</code>	<code>\)</code>	$U_{\text{e,final}}$

N 2021-02-13

`\springpotentialenergy[⟨label⟩]`

Symbol for spring potential energy.

<code>\(\springpotentialenergy \)</code>	<code>\)</code>	U_{s}
<code>\(\springpotentialenergy[\text{symup{final}}] \)</code>	<code>\)</code>	$U_{\text{s,final}}$

8.3 mandiexp Source Code

Define the package version and date for global use, exploiting the fact that in a .sty file there is now no need for `\makeatletter` and `\makeatother`. This simplifies defining internal commands, with @ in the name, that are not for the user to know about.

```
1 \def\mandiexp@version{3.2.2a}
2 \def\mandiexp@date{2024-09-12}
3 \NeedsTeXFormat{LaTeX2e}[2020-02-02]
4 \DeclareRelease{v3.2.2a}{2024-09-12}{mandiexp.sty}
5 \DeclareCurrentRelease{v\mandiexp@version}{\mandiexp@date}
6 \ProvidesPackage{mandiexp}
7  [\mandiexp@date\space v\mandiexp@version\space Macros for Matter & Interactions]
```

Define a convenient package version command.

```
8 \newcommand*{\mandiexpversion}{v\mandiexp@version\space dated \mandiexp@date}
```

We need mandi to load mathtools and unicode-math.

```
9 \IfPackageLoadedTF{mandi}%
10 {}%
11 {%
12   \RequirePackage{mandi}%
13 }%
```

We need mandistudent for the new `\vec*` command.

```
14 \IfPackageLoadedTF{mandistudent}%
15 {}%
16 {%
17   \RequirePackage{mandistudent}%
18 }%
```

Load xparse if necessary.

```
19 \IfFormatAtLeastTF{2020-10-01}%
20 {}%
21 {\RequirePackage{xparse}}%
```

We require the Lua^AT_EX engine.

```
22 \RequireLuaTeX
23 \typeout{%
24 \typeout{mandiexp: You are using mandiexp \mandiexpversion.}
25 \typeout{mandiexp: This package requires LuaLaTeX.}%
26 \typeout{}}
```

The momentum principle's lefthand and righthand sides.

```
27 \NewDocumentCommand{\lhmomentumprinciple}{ s }%
28 {%
29   \Delta
30   \IfBooleanTF{#1}%
31     {%
32       \vec*{p}
33     }%
34     {%
35       \vec{p}%
36     }%
37   \sb{\symup{sys}}}%
38 }%
39 \NewDocumentCommand{\rhsmomentumprinciple}{ s }%
40 {%
41   \IfBooleanTF{#1}%
42   {%
```



```

43      \vec*{F}%
44    }%
45    {%
46      \vec{F}%
47    }%
48    \sb{\symup{sys,net}}\,\Delta t%
49  }%

```

The momentum principle in update form, lefthand and righthand sides.

```

50 \NewDocumentCommand{\lhsmomentumprincipleupdate}{ s }%
51  {%
52    \IfBooleanTF{#1}%
53    {%
54      \vec*{p}%
55    }%
56    {%
57      \vec{p}%
58    }%
59    \sb{\symup{sys,final}}%
60  }%
61 \NewDocumentCommand{\rhsmomentumprincipleupdate}{ s }%
62  {%
63    \IfBooleanTF{#1}%
64    {%
65      \vec*{p}%
66    }%
67    {%
68      \vec{p}%
69    }%
70    \sb{\symup{sys,initial}}+%
71    \IfBooleanTF{#1}%
72    {%
73      \vec*{F}%
74    }%
75    {%
76      \vec{F}%
77    }%
78    \sb{\symup{sys,net}}\,\Delta t%
79  }%

```

The full momentum principle as an expression.

```

80 \NewDocumentCommand{\momentumprinciple}{ s }%
81  {%
82    \IfBooleanTF{#1}%
83    {%
84      \lhsmomentumprinciple* = \rhsmomentumprinciple*%
85    }%
86    {%
87      \lhsmomentumprinciple = \rhsmomentumprinciple%
88    }%
89  }%

```

The full momentum principle in update form as an expression.

```

90 \NewDocumentCommand{\momentumprincipleupdate}{ s }%
91  {%
92    \IfBooleanTF{#1}%
93    {%
94      \lhsmomentumprincipleupdate* = \rhsmomentumprincipleupdate*%
95    }%

```

```

96      {%
97      \lhsmomentumprincipleupdate = \rhsmomentumprincipleupdate%
98      }%
99  }%

```

The energy principle's lefthand and righthand sides.

```

100 \NewDocumentCommand{\lhsenergyprinciple}{}%
101   {%
102   \Delta E_{\symup{sys}}%
103   }%
104 \NewDocumentCommand{\rhsenergyprinciple}{ 0{ } }%
105   {%
106   W_{\symup{ext}}#1%
107   }%

```

The energy principle in update form, lefthand and righthand sides.

```

108 \NewDocumentCommand{\lhsenergyprincipleupdate}{}%
109   {%
110   E_{\symup{sys,final}}%
111   }%
112 \NewDocumentCommand{\rhsenergyprincipleupdate}{ 0{ } }%
113   {%
114   E_{\symup{sys,initial}}+%
115   W_{\symup{ext}}#1%
116   }%

```

The full energy principle as an expression.

```

117 \NewDocumentCommand{\energyprinciple}{ 0{ } }%
118   {%
119   \lhsenergyprinciple = \rhsenergyprinciple[#1]%
120   }%

```

The full energy principle in update form as an expression.

```

121 \NewDocumentCommand{\energyprincipleupdate}{ 0{ } }%
122   {%
123   \lhsenergyprincipleupdate = \rhsenergyprincipleupdate[#1]%
124   }%

```

The angular momentum principle's lefthand and righthand sides.

```

125 \NewDocumentCommand{\lhsangularmomentumprinciple}{ s }%
126   {%
127   \Delta%
128   \IfBooleanTF{#1}%
129     {%
130     \vec*{L}%
131     }%
132     {%
133     \vec{L}%
134     }%
135   \sb{A\symup{,sys,net}}%
136   }%
137 \NewDocumentCommand{\rhsangularmomentumprinciple}{ s }%
138   {%
139   \IfBooleanTF{#1}%
140     {%
141     \vec*{\tau}%
142     }%
143     {%
144     \vec{\tau}%

```

```

145     }%
146     \sb{A\symup{,sys,net}}\,\Delta t%
147 }%

```

The energy principle in update form, lefthand and righthand sides.

```

148 \NewDocumentCommand{\lhsangularmomentumprincipleupdate}{ s }%
149 {%
150     \IfBooleanTF{#1}%
151     {%
152         \vec*{L}%
153     }%
154     {%
155         \vec{L}%
156     }%
157     \sb{A,\symup{sys,final}}%
158 }%
159 \NewDocumentCommand{\rhsangularmomentumprincipleupdate}{ s }%
160 {%
161     \IfBooleanTF{#1}%
162     {%
163         \vec*{L}%
164     }%
165     {%
166         \vec{L}%
167     }%
168     \sb{A\symup{,sys,initial}}+%
169     \IfBooleanTF{#1}%
170     {%
171         \vec*{\tau}%
172     }%
173     {%
174         \vec{\tau}%
175     }%
176     \sb{A\symup{,sys,net}}\,\Delta t%
177 }%

```

The full angular momentum principle as an expression.

```

178 \NewDocumentCommand{\angularmomentumprinciple}{ s }%
179 {%
180     \IfBooleanTF{#1}%
181     {%
182         \lhsangularmomentumprinciple* = \rhsangularmomentumprinciple*%
183     }%
184     {%
185         \lhsangularmomentumprinciple = \rhsangularmomentumprinciple%
186     }%
187 }%

```

The full angular momentum principle in update form as an expression.

```

188 \NewDocumentCommand{\angularmomentumprincipleupdate}{ s }%
189 {%
190     \IfBooleanTF{#1}%
191     {%
192         \lhsangularmomentumprincipleupdate* = \rhsangularmomentumprincipleupdate*%
193     }%
194     {%
195         \lhsangularmomentumprincipleupdate = \rhsangularmomentumprincipleupdate%
196     }%
197 }%

```

The symbol for an arbitrary entity.

```

198 \NewDocumentCommand{\energyof}{ m o }%
199 {%
200   E_{#1%
201     \IfValueT{#2}%
202       {,#2}%
203   }%
204 }%
```

The symbol for a system's energy.

```

205 \NewDocumentCommand{\systemenergy}{ o }%
206 {%
207   E_{\symup{sys}%
208     \IfValueT{#1}%
209       {,#1}%
210   }%
211 }%
212 \NewDocumentCommand{\particleenergy}{ o }%
213 {%
214   E_{\symup{particle}%
215     \IfValueT{#1}%
216       {,#1}%
217   }%
218 }%
```

The symbol for a particle's rest energy.

```

219 \NewDocumentCommand{\restenergy}{ o }%
220 {%
221   E_{\symup{rest}%
222     \IfValueT{#1}%
223       {,#1}%
224   }%
225 }%
```

The symbol for a system's internal energy.

```

226 \NewDocumentCommand{\internalenergy}{ o }%
227 {%
228   E_{\symup{internal}%
229     \IfValueT{#1}%
230       {,#1}%
231   }%
232 }%
```

The symbol for a system's chemical energy.

```

233 \NewDocumentCommand{\chemicalenergy}{ o }%
234 {%
235   E_{\symup{chem}%
236     \IfValueT{#1}%
237       {,#1}%
238   }%
239 }%
```

The symbol for a system's thermal energy.

```

240 \NewDocumentCommand{\thermalenergy}{ o }%
241 {%
242   E_{\symup{therm}%
243     \IfValueT{#1}%
244       {,#1}%
245   }%
246 }%
```

The symbol for a photon's energy.

```

247 \NewDocumentCommand{\photonenergy}{ o }%
248 {%
249   E_{\symup{photon}}%
250   \IfValueT{#1}%
251     {,#1}%
252   }%
253 }%

```

The symbol for translational kinetic energy. `d[]` must be used because of the way consecutive optional arguments are handled. See `xparse` docs for details.

See <https://tex.stackexchange.com/a/569011/218142>.

```

254 \NewDocumentCommand{\translationalkineticenergy}{ s d[] }%
255 {%
256   \IfBooleanTF{#1}%
257     {%
258       E_{\bgroup \symup{K}}%
259     }%
260     {%
261       K_{\bgroup \symup{trans}}%
262     }%
263     \IfValueT{#2}{,#2}%
264     \egroup%
265 }%

```

The symbol for rotational kinetic energy.

```

266 \NewDocumentCommand{\rotationalkineticenergy}{ s d[] }%
267 {%
268   \IfBooleanTF{#1}%
269     {%
270       E_{\bgroup%
271       }%
272       {%
273         K_{\bgroup%
274         }%
275         \symup{rot}\IfValueT{#2}{,#2}%
276       \egroup%
277 }%

```

The symbol for vibrational kinetic energy.

```

278 \NewDocumentCommand{\vibrationalkineticenergy}{ s d[] }%
279 {%
280   \IfBooleanTF{#1}%
281     {%
282       E_{\bgroup%
283       }%
284       {%
285         K_{\bgroup%
286         }%
287         \symup{vib}\IfValueT{#2}{,#2}%
288       \egroup%
289 }%

```

The symbol for a system's gravitational potential energy.

```

290 \NewDocumentCommand{\gravitationalpotentialenergy}{ o }%
291 {%
292   U_{\symup{g}}%
293   \IfValueT{#1}%

```

```

294      {,#1}%
295    }%
296  }%

```

The symbol for a system's electric potential energy.

```

297 \NewDocumentCommand{\electricpotentialenergy}{ o }%
298 {%
299   U_{\symp{e}%
300     \IfValueT{#1}%
301       {,#1}%
302     }%
303 }%

```

The symbol for a system's spring potential energy.

```

304 \NewDocumentCommand{\springpotentialenergy}{ o }%
305 {%
306   U_{\symp{s}%
307     \IfValueT{#1}%
308       {,#1}%
309     }%
310 }%

```

Index

Page numbers refer to page where the corresponding entry is documented and/or referenced.

A			
<code>\absolutevalue</code>	55	<code>\CheckConstant</code>	24
<code>\absolutevalue*</code>	55	<code>\CheckQuantity</code>	10
<code>\acceleration</code>	10	<code>\chemicalenergy</code>	86
<code>\accelerationvector</code>	10	<code>\cmagneticfield</code>	12
alternate value	8, 36	<code>\cmagneticfieldvector</code>	12
<code>\alwaysusealternateunits</code>	22	<code>\colvec</code>	60
<code>\alwaysuseapproximateconstants</code>	31	<code>\conductance</code>	12
<code>\alwaysusebaseunits</code>	22	<code>\conductivity</code>	12
<code>\alwaysusederivedunits</code>	22	<code>\contraction</code>	61
<code>\alwaysusepreciseconstants</code>	31	<code>\contraction*</code>	61
<code>\amount</code>	10	<code>\conventionalcurrent</code>	12
<code>\ampere</code>	31	<code>\coulomb</code>	32
<code>\anglebrackets</code>	53	<code>\coulombconstant</code>	25
<code>\anglebrackets*</code>	53	<code>\curlybraces</code>	54
<code>\angularacceleration</code>	10	<code>\curlybraces*</code>	54
<code>\angularaccelerationvector</code>	10	<code>\current</code>	12
<code>\angularfrequency</code>	10	<code>\currentdensity</code>	12
<code>\angularimpulse</code>	10	<code>\currentdensityvector</code>	12
<code>\angularimpulsevector</code>	10	D	
<code>\angularmomentum</code>	11	<code>\degree</code>	32
<code>\angularmomentumprinciple</code>	84	derived value	8, 36
<code>\angularmomentumprinciple*</code>	84	<code>\df</code>	62
<code>\angularmomentumprincipleupdate</code>	84	<code>\dielectricconstant</code>	12
<code>\angularmomentumprincipleupdate*</code>	84	<code>\direction</code>	13
<code>\angularmomentumvector</code>	11	<code>\directionvector</code>	13
<code>\angularvelocity</code>	11	<code>\dirvec</code>	52
<code>\angularvelocityvector</code>	11	<code>\dirvec*</code>	53
<code>\area</code>	11	<code>\displacement</code>	13
<code>\areachargedensity</code>	11	<code>\displacementvector</code>	13
<code>\areamassdensity</code>	11	<code>\doublebars</code>	53
<code>\atomicmassunit</code>	31	<code>\doublebars*</code>	53
<code>\avogadro</code>	24	<code>\duration</code>	13
B		E	
base value	8, 36	<code>\earthmass</code>	25
<code>\biotsavartconstant</code>	24	<code>\earthmoondistance</code>	25
<code>\bohrradius</code>	24	<code>\earthradius</code>	25
<code>\boltzmann</code>	25	<code>\earthsundistance</code>	25
C		<code>\electricdipolemoment</code>	13
<code>\candela</code>	31	<code>\electricdipolemomentvector</code>	13
<code>\capacitance</code>	11	<code>\electricfield</code>	13
<code>\changein</code>	53	<code>\electricfieldvector</code>	13
<code>\charge</code>	11	<code>\electricflux</code>	13
		<code>\electricpotential</code>	13

<code>\electricpotentialdifference</code>	14
<code>\electricpotentialenergy</code>	87
<code>\electronCharge</code>	26
<code>\electroncharge</code>	26
<code>\electroncurrent</code>	14
<code>\electronmass</code>	26
<code>\electronvolt</code>	32
<code>\elementarycharge</code>	26
<code>\emf</code>	14
<code>\emptyunit</code>	31
<code>\energy</code>	14
<code>\energydensity</code>	14
<code>\energyflux</code>	15
<code>\energyfluxvector</code>	15
<code>\energyinev</code>	14
<code>\energyinkev</code>	14
<code>\energyinmev</code>	14
<code>\energyof</code>	85
<code>\energyprinciple</code>	84
<code>\energyprincipleupdate</code>	84
<code>\entropy</code>	15
Environments	
<code>parts</code>	56
<code>physicsproblem</code>	56
<code>physicsproblem*</code>	56
<code>physicssolution</code>	57
<code>physicssolution*</code>	57
<code>usealternateunits</code>	23
<code>useapproximateconstants</code>	31
<code>usebaseunits</code>	23
<code>usederivedunits</code>	23
<code>usepreciseconstants</code>	31
<code>webvpythonblock</code>	62
<code>webvpythonblock*</code>	62
<code>\ev</code>	32
F	
<code>false value</code>	8, 36
<code>\farad</code>	32
<code>\finestructure</code>	26
<code>\force</code>	15
<code>\forcevector</code>	15
<code>\frequency</code>	15
G	
<code>\gravitationalfield</code>	15
<code>\gravitationalfieldvector</code>	15
<code>\gravitationalpotential</code>	15
<code>\gravitationalpotentialdifference</code>	15
<code>\gravitationalpotentialenergy</code>	87

H	
<code>\hbar</code>	33
<code>\henry</code>	32
<code>\hereusealternateunits</code>	22
<code>\hereuseapproximateconstants</code>	31
<code>\hereusebaseunits</code>	22
<code>\hereusederivedunits</code>	22
<code>\hereusepreciseconstants</code>	31
<code>\hertz</code>	32
<code>\hilite</code>	58
<code>\hydrogenmass</code>	26

I	
<code>\image</code>	59
<code>\impulse</code>	16
<code>\impulsevector</code>	16
<code>\indexofrefraction</code>	16
<code>\inductance</code>	16
<code>\internalenergy</code>	86
<code>\inverse</code>	32

J	
<code>\joule</code>	32

K	
<code>\kelvin</code>	32
<code>\kev</code>	32
Keys	
<code>preciseconstants</code>	8
<code>units</code>	8
<code>\kiloelectronvolt</code>	32
<code>\kilogram</code>	32

L	
<code>\lhsangularmomentumprinciple</code>	84
<code>\lhsangularmomentumprinciple*</code>	84
<code>\lhsangularmomentumprincipleupdate</code>	84
<code>\lhsangularmomentumprincipleupdate*</code>	84
<code>\lhsenergyprinciple</code>	84
<code>\lhsenergyprincipleupdate</code>	84
<code>\lhsmomentumprinciple</code>	83
<code>\lhsmomentumprinciple*</code>	83
<code>\lhsmomentumprincipleupdate</code>	83
<code>\lhsmomentumprincipleupdate*</code>	83
<code>\lightspeed</code>	32
<code>\linearchargedensity</code>	16
<code>\linearmassdensity</code>	16
<code>\lorentzfactor</code>	16
<code>\luminousintensity</code>	16

M	
\magneticcharge	17
\magneticdipolemoment	17
\magneticdipolemomentvector	17
\magneticfield	17
\magneticfieldvector	17
\magneticflux	17
\magnitude	55
\magnitude*	55
\mandiexpversion	83
\mandisetaup	8
\mandistudentversion	52
\mandiversion	8
\mass	17
\megaelectronvolt	32
\meter	32
\metre	32
\mev	32
\mivector	34
\mobility	17
\mole	32
\momentofinertia	17
\momentum	9, 18
\momentumflux	18
\momentumfluxvector	18
\momentumprinciple	83
\momentumprinciple*	83
\momentumprincipleupdate	83
\momentumprincipleupdate*	83
\momentumvector	9, 18
\moonearthdistance	27
\moonmass	27
\moonradius	27
\mzofp	27
N	
\neutronmass	27
\NewPhysicalConstant	30
\NewScalarQuantity	22
\newton	32
\NewVectorQuantity	22
\norm	55
\norm*	55
\numberdensity	18
O	
\ohm	32
\oofpez	23, 27
\oofpezcs	28

P	
\parallelto	55
\parentheses	53
\parentheses*	53
\particleenergy	85
parts environment	56
\pascal	32
\per	31
\permeability	18
\permittivity	18
\perpendicularto	55
\photonenergy	86
physicsproblem environment	56
physicsproblem* environment	56
physicssolution environment	57
physicssolution* environment	57
\planck	28
\planckbar	28
\planckc	28
\planeangle	18
\polarizability	18
\power	18
\poynting	19
\poyntingvector	19
preciseconstants key	8
\pressure	19
\problempart	56
\protonCharge	28
\protoncharge	28
\protonmass	29
R	
\radian	32
\reason	57
\relativepermeability	19
\relativepermittivity	19
\RenewPhysicalConstant	30
\RenewScalarQuantity	22
\RenewVectorQuantity	22
\resistance	19
\resistivity	19
\restenergy	85
\rhsangularmomentumprinciple	84
\rhsangularmomentumprinciple*	84
\rhsangularmomentumprincipleupdate	84
\rhsangularmomentumprincipleupdate*	84
\rhsenergyprinciple	84
\rhsenergyprincipleupdate	84
\rhsmomentumprinciple	83
\rhsmomentumprinciple*	83

<code>\rhsmomentumprincipleupdate</code>	83
<code>\rhsmomentumprincipleupdate*</code>	83
<code>\rotationalkineticenergy</code>	86
<code>\rotationalkineticenergy*</code>	86
<code>\rowvec</code>	60
<code>\rydberg</code>	29

S

<code>\second</code>	32
<code>\siemens</code>	32
<code>\singlebars</code>	53
<code>\singlebars*</code>	53
<code>\slot</code>	61
<code>\slot*</code>	61
<code>\solidangle</code>	19
<code>\specificeatcapacity</code>	20
<code>\speedoflight</code>	29
<code>\springpotentialenergy</code>	87
<code>\springstiffness</code>	20
<code>\springstretch</code>	20
<code>\squarebrackets</code>	53
<code>\squarebrackets*</code>	53
<code>\stefanboltzmann</code>	29
<code>\steradian</code>	32
<code>\strain</code>	20
<code>\stress</code>	20
<code>\sunearthdistance</code>	29
<code>\sunradius</code>	29
<code>\surfacegravfield</code>	30
<code>\systemenergy</code>	85

T

<code>\temperature</code>	20
<code>\tencomp</code>	61
<code>\tencomp*</code>	61
<code>\tento</code>	34
<code>\tesla</code>	32
<code>\thermalenergy</code>	86
<code>\timestento</code>	34
<code>\torque</code>	20
<code>\torquevector</code>	20
<code>\tothefour</code>	32
<code>\totheinversefour</code>	32
<code>\totheinversethree</code>	32
<code>\totheinversetwo</code>	32
<code>\tothethree</code>	32
<code>\tothetwo</code>	32
<code>\translationalkineticenergy</code>	86
<code>\translationalkineticenergy*</code>	86
true value	8, 36

U

<code>\units</code>	31
units key	8
<code>\universalgrav</code>	30
usealternateunits environment	23
useapproximateconstants environment	31
usebaseunits environment	23
usederivedunits environment	23
usepreciseconstants environment	31
<code>\usk</code>	31

V

<code>\vacuumpermeability</code>	30
<code>\vacuumpermittivity</code>	30
<code>\valence</code>	61
<code>\valence*</code>	61
Values	
alternate	8, 36
base	8, 36
derived	8, 36
false	8, 36
true	8, 36
<code>\vec</code>	52
<code>\vec*</code>	52
<code>\veccomp</code>	61
<code>\veccomp*</code>	61
<code>\vectoracceleration</code>	10
<code>\vectorangularacceleration</code>	10
<code>\vectorangularimpulse</code>	10
<code>\vectorangularmomentum</code>	11
<code>\vectorangularvelocity</code>	11
<code>\vectorcmagneticfield</code>	12
<code>\vectorcurrentdensity</code>	12
<code>\vectordirection</code>	13
<code>\vectordisplacement</code>	13
<code>\vectorelectricdipolemoment</code>	13
<code>\vectorelectricfield</code>	13
<code>\vectorenergyflux</code>	15
<code>\vectorforce</code>	15
<code>\vectorgravitationalfield</code>	15
<code>\vectorimpulse</code>	16
<code>\vectormagneticdipolemoment</code>	17
<code>\vectormagneticfield</code>	17
<code>\vectormomentum</code>	9, 18
<code>\vectormomentumflux</code>	18
<code>\vectorpoynting</code>	19
<code>\vectortorque</code>	20
<code>\vectorvelocity</code>	20
<code>\vectorvelocityc</code>	21
<code>\vectorwavenumber</code>	21

<code>\velocity</code>	20
<code>\velocityc</code>	21
<code>\velocitycvector</code>	21
<code>\velocityvector</code>	20
<code>\vibrationalkineticenergy</code>	87
<code>\vibrationalkineticenergy*</code>	87
<code>\volt</code>	32
<code>\volume</code>	21
<code>\volumechargedensity</code>	21
<code>\volumemassdensity</code>	21
<code>\vpythonfile</code>	68
<code>\vpythoninline</code>	70

W

<code>\watt</code>	32
<code>\wavelength</code>	21
<code>\wavenumber</code>	21
<code>\wavenumbervector</code>	21
<code>\weber</code>	32
<code>webvpythonblock</code> environment	62
<code>webvpythonblock*</code> environment	62
<code>\webvpythoninline</code>	70
<code>\work</code>	21

X

<code>\xtento</code>	34
----------------------------	----

Y

<code>\youngsmodulus</code>	22
-----------------------------------	----

Z

<code>\zerovec</code>	53
<code>\zerovec*</code>	53