

Lecture Note 3

Confidence Intervals

1 Confidence Intervals for Means

With Normally distributed data,

$$T_n = \frac{\bar{X} - \mu_X}{s_X/\sqrt{n}} \sim t(n-1),$$

where s_X is the sample standard deviation of X_i . And statistic T_n (called a “t-statistic”) is also asymptotically standard Normal regardless of the distribution of X_i .

Let c_α denote the critical value for a two-sided α -level test. For example, when $\alpha = .05$, $c_\alpha \approx 2$.

- A t-statistic is bracketed by $\pm c_n$ with probability $1 - \alpha$:

$$P[-c_\alpha \leq T_n \leq c_\alpha] = 1 - \alpha$$

Rearranging, we have:

$$P[\bar{X} - c_\alpha(s_X/\sqrt{n}) \leq \mu_X \leq \bar{X} + c_\alpha(s_X/\sqrt{n})] = 1 - \alpha$$

- s_X/\sqrt{n} is the *estimated standard error of the sample mean*, \bar{X} , in a sample of size n
- We say that

$$[\bar{X} - c_\alpha(s_X/\sqrt{n}), \bar{X} + c_\alpha(s_X/\sqrt{n})]$$

provides a $(1 - \alpha)$ confidence interval for μ_X .

- A 95% c.i. brackets μ_X 95% of the time. *Hey, what time is that?* When we draw repeated samples.
- But we don’t draw repeated samples, we draw only one! Still, this probabilistic statement tells us how likely the interval is to *cover* the pop mean were we to conduct such a sampling experiment. Like standard errors, confidence intervals quantify *sampling variance*.
- Confidence intervals are random; Parameters (like μ_X) are fixed.
- In practice, we often ballpark the c.i.
 - A rule of thumb 95% confidence interval for a pop mean is given by the sample mean plus/minus two standard errors (since, if $\alpha = .05$, $c_\alpha = 1.96$ for Normal)

2 Confidence Intervals for Differences in Means

- The t-statistic for a *difference* in means is

$$T_n = \frac{(\bar{X}_1 - \bar{X}_2) - (\mu_1 - \mu_2)}{([s_1^2/n_1] + [s_2^2/n_2])^{1/2}} \sim t(n_1 + n_2 - 2),$$

where $T_n \sim N(0, 1)$ for large n

- This comes from the fact that the *standard error of a difference in means* is

$$s^* = ([s_1^2/n_1] + [s_2^2/n_2])^{1/2}$$

so,

$$P[(\bar{X}_1 - \bar{X}_2) - c_\alpha s^* \leq \mu_1 - \mu_2 \leq (\bar{X}_1 - \bar{X}_2) + c_\alpha s^*] = 1 - \alpha,$$

where c_α is the $t(n_1 + n_2 - 2)$ critical value for a two-sided α -level test.

- A rule of thumb confidence interval for a difference in means is $(\bar{X}_1 - \bar{X}_2) \pm 2s^*$.

3 Confidence Intervals for Treatment Effects in ALO (2009)

TABLE 5—TREATMENT EFFECTS ON FIRST YEAR OUTCOMES IN THE SAMPLE WITH FALL GRADES

	SFP by type			Any SFP		
	All (1)	Men (2)	Women (3)	All (4)	Men (5)	Women (6)
<i>Panel A. Fall grade</i>						
Control mean	64.225 (11.902)	65.935 (11.340)	62.958 (12.160)	64.225 (11.902)	65.935 (11.340)	62.958 (12.160)
SSP	0.349 [0.917]	−0.027 [1.334]	0.737 [1.275]	0.344 [0.917]	−0.014 [1.332]	0.738 [1.274]
SFP	1.824 [0.847]**	0.331 [1.233]	2.602 [1.176]**			
SFSP	2.702 [1.124]**	−0.573 [2.010]	4.205 [1.325]***			
SFP (any)				2.125 [0.731]***	0.016 [1.164]	3.141 [0.972]***
Observations	1,255	526	729	1,255	526	729
<i>Panel B. First year GPA</i>						
Control mean	1.805 (0.902)	1.908 (0.908)	1.728 (0.891)	1.797 (0.904)	1.885 (0.910)	1.731 (0.894)
SSP	0.073 [0.066]	0.011 [0.107]	0.116 [0.082]	0.071 [0.066]	0.008 [0.107]	0.116 [0.082]
SFP	0.010 [0.064]	−0.110 [0.103]	0.086 [0.084]			
SFSP	0.210 [0.092]**	0.084 [0.162]	0.267 [0.117]**			
SFP (any)				0.079 [0.056]	−0.042 [0.095]	0.147 [0.073]**
Observations	1,255	526	729	1,255	526	729

Notes: The table reports regression estimates of treatment effects on full grades and first-year GPA computed using the full set of controls. Robust standard errors are reported in brackets. The sample is limited to students registered for at least two courses as of November 1 with data on the relevant set of controls and at least one fall grade. The last three columns report estimates from a model that combines the SFP and SFSP treatment groups into “SFP (any).”

* Significant at the 10 percent level.

** Significant at the 5 percent level.

*** Significant at the 1 percent level.

Ponder these questions, grasshopper:

- What determines c.i. width?
- Why are narrow intervals preferred to wide?
- Whats the connection between c.i. width and statistical power?

4 Monte Carlo Interval Coverage for Immigrant-Native Wage Diff

- Asymptotic intervals cover well in real (definitely not Normal) data, with $n = 100$ only

```

name: <unnamed>
log: /Users/joshangrist/Documents/teaching/14.32/SP2023/notes/LN3/LN3stata/LN3.log
log type: text
opened on: 16 Feb 2023, 12:25:16

```

```

.
.      /*
.      14.32 Lecture Note 3
.      c.i. coverage for immigrant-native wage gap
.      edited 02-16-23 to drop zeros and use unequal var ests
.      */
.
.      // Insheet data from 2016 ACS PUMS
.      import delimited "ss16pusa.csv", clear
(encoding automatically selected: UTF-8)
(284 vars, 1,623,216 obs)

.
.      // Sample of interest is men aged 40-49
.      keep if sex==1 & age>=40 & age <=49
(1,526,769 observations deleted)

.
.      // Restrict sample to men who worked 50-52 weeks
.      keep if wkwm==1
(24,305 observations deleted)

.
.      // Generate usual hourly earnings
.      lab var wagp "Raw annual earnings"

.      lab var wkhp "Usual hours per week"

.      gen uhe = wagp/(50*wkhp)

.      quietly sum uhe, d

.      replace uhe=. if uhe>`r(p99)'
(819 real changes made, 819 to missing)

.      lab var uhe "Usual hourly earnings (truncated at 99pct)"

.      gen loguhe=log(uhe)
(4,963 missing values generated)

.      lab var loguhe "Log hourly earnings"

.
.      // Code up immigrants
.      gen immig=(nativity==2)

.      lab var immig "Foreign born"

.
.      ttest loguhe, by(immig)

Two-sample t test with equal variances

+-----+-----+-----+-----+-----+-----+
| Group | Obs   | Mean   | Std. err. | Std. dev. | [95% conf. interval] |
+-----+-----+-----+-----+-----+-----+
| 0      | 52,300 | 3.300842 | .0031697 | .724887 | 3.294629 3.307055 |
| 1      | 14,879 | 3.137078 | .0064108 | .7819809 | 3.124512 3.149644 |
+-----+-----+-----+-----+-----+-----+
| Combined | 67,179 | 3.264571 | .002859 | .7410341 | 3.258967 3.270175 |
+-----+-----+-----+-----+-----+-----+
| diff   |        | .1637641 | .0068562 |          | .1503259 .1772022 |
+-----+-----+-----+-----+-----+-----+
| diff = mean(0) - mean(1) | t = 23.8855 |
| H0: diff = 0 | Degrees of freedom = 67177 |

      Ha: diff < 0      Ha: diff != 0      Ha: diff > 0
Pr(T < t) = 1.0000      Pr(|T| > |t|) = 0.0000      Pr(T > t) = 0.0000

.      gen popmu_native=r(mu_1)
.      gen popmu_immig=r(mu_2)
.      gen popdiff=popmu_native-popmu_immig

.
.      // Keep only key variables in a newdata set called "workingextract"
.
.      drop if missing(loguhe)
(4,963 observations deleted)

.
.      keep wagp loguhe uhe wkhp immig age popmu_native popmu_immig popdiff

.      save workingextract, replace
file workingextract.dta saved

.      summarize

```

Variable	Obs	Mean	Std. dev.	Min	Max
agep	67,179	44.59836	2.843473	40	49
wagp	67,179	77017	70468.34	4	665000
wkhp	67,179	44.73532	9.786132	1	99
uhe	67,179	33.90219	27.61486	.0016	201.5789
loguhe	67,179	3.264571	.7410341	-6.437752	5.306181
immig	67,179	.2214829	.4152479	0	1
popmu_native	67,179	3.300842	0	3.300842	3.300842
popmu_immig	67,179	3.137078	0	3.137078	3.137078

```

popdiff |      67,179      .163764      0      .163764      .163764
. keep if _n==1
(67,178 observations deleted)

. keep popdiff

. save expresresults, replace
file expresresults.dta saved

.
. /* draw samples of N=100 500 times compute means, diff, SEdiff */
.
.   forvalues s=1/500 {
2.       quietly use workingextract, clear
3.       quietly bsample 100
4.       quietly ttest loguhe, by(immig) unequal
5.       quietly gen mu_nat=r(mu_1)
6.       quietly gen mu_imm=r(mu_2)
7.       quietly gen diff=mu_nat-mu_imm
8.       quietly gen numNat=r(N_1)
9.       quietly gen numImm=r(N_2)
10.      quietly gen SEdiff=r(se)
11.      quietly gen hi95 = diff + (SEdiff*1.96)
12.      quietly gen lo95 = diff - (SEdiff*1.96)
13.      quietly gen cover95=0
14.      quietly replace cover95 = 1 if lo95<=popdiff & popdiff<=hi95
15.      quietly keep if _n==1
16.      quietly append using expresresults
17.      quietly save expresresults, replace
18.  }

.
.   keep popdiff mu_nat mu_imm diff numNat numImm SEdiff hi95 lo95 cover95

.   drop if missing(diff)
(1 observation deleted)

.
.   /* sampling experiment results */
.
.   list popdiff diff numNat numImm SEdiff lo95 hi95 cover95 if _n<=5

```

	popdiff	diff	numNat	numImm	SEdiff	lo95	hi95	cover95
1.	.163764	.0578763	76	24	.1725974	-.2804146	.3961673	1
2.	.163764	.2617667	78	22	.1882998	-.107301	.6308343	1
3.	.163764	.1345735	84	16	.1860601	-.2301043	.4992512	1
4.	.163764	.4340758	79	21	.1863383	.0688528	.7992989	1
5.	.163764	-.1437666	86	14	.1949807	-.5259289	.2383956	1

```

.   summarize diff lo95 hi95 cover95

Variable |      Obs      Mean   Std. dev.      Min      Max
-----+-----+-----+-----+-----+-----
diff     |      500   .1713004   .1923067   -.3448374   .8456693
lo95     |      500  -.1913058   .198277   -.7165971   .4191974
hi95     |      500   .5339065   .2044576   -.0338668   1.797269
cover95  |      500      .948   .2222494      0      1

```

```

.   log close
.   name: <unnamed>
.   log: /Users/joshangrist/Documents/teaching/14.32/SP2023/notes/LN3/LN3stata/LN3.log
.   log type: text
.   closed on: 16 Feb 2023, 12:26:34

```