




Ground Truth: Porphyria

MedGemma

The image shows a case of severe psoriasis, characterized by erythematous, scaly plaques with visible pustules, particularly on the hands. This indicates a significant inflammatory skin condition. The standard treatment for severe psoriasis includes a combination of topical corticosteroids, vitamin D analogs (calcipotriene), and phototherapy (UVB or PUVA). Adjunctive therapies include emollients for skin hydration and topical calcineurin inhibitors for mild to moderate psoriasis.

Primary Diagnosis: Psoriasis

Hulu-Med

The image shows a case of psoriasis vulgaris, plaque type, characterized by erythematous, scaly plaques with visible thickening, particularly on areas prone to plaque formation. This indicates a chronic inflammatory skin condition with well-demarcated lesions. The Standard treatment for psoriasis vulgaris includes a combination of topical corticosteroids, vitamin D analogsAdjunctive therapies include moisturizers for skin hydration and sun protection to prevent exacerbation.

Primary Diagnosis: Psoriasis Vulgaris

SkinGPT-X

Key Findings: The image shows multiple, discrete, raised, irregularly shaped nodules and papules on the dorsal hand and fingers. Lesions are erythematous to violaceous with hyperpigmentation, central ulceration/crusting, irregular/poorly defined borders, and heterogeneous surface texture (scaling, crusting, ulceration, induration). Marked asymmetry in distribution. No prominent telangiectasia or lichenification.

Case Reviewing: The visual findings of multiple, irregularly shaped, erythematous to violaceous nodules and papules with scaling, crusting, ulceration, and poorly defined borders are inconsistent with the standard description of erythema elevatum diutinum (smooth, glistening, well-circumscribed, no scaling/crusting/ulceration). **While the model predicts erythema elevatum diutinum (23.69%), the description aligns better with the potential presentations of porphyria (15.44%), which can include papules, nodules, plaques with scaling, crusting, and irregular borders.** Past cases of porphyria show varied morphologies, including scaly plaques and erosions, supporting the possibility of an atypical presentation. Factitial dermatitis (2.80%) is also a consideration given the ulceration and irregular morphology, but the model ranks it lower. Lupus erythematosus (4.87%) is less likely given the lack of typical features like atrophy or prominent telangiectasia. Milia (3.43%) is clearly excluded by the morphology (nodules, papules, plaques, ulceration vs. small, smooth papules).

Primary Diagnosis: Porphyria