

**Subject:** Panniculectomy and Abdominoplasty

**Guideline #:** CG-SURG-99

**Status:** Reviewed

**Publish Date:** 04/10/2024

**Last Review Date:** 02/15/2024

## Description

This document addresses the surgical procedures panniculectomy and abdominoplasty and when they are considered medically necessary, not medically necessary, and cosmetic.

**Medically Necessary:** In this document, procedures are considered medically necessary if there is a significant functional impairment AND the procedure can be reasonably expected to improve the functional impairment.

**Cosmetic:** In this document, procedures are considered cosmetic when intended to change a physical appearance that would be considered within normal human anatomic variation. Cosmetic services are often described as those that are primarily intended to preserve or improve appearance.

## Clinical Indications

### Medically Necessary:

- A. Panniculectomy is considered **medically necessary** for the individual who meets the following criteria:
  1. The panniculus hangs below the level of the pubis (which is documented in photographs); **and**
  2. **One** of the following:
    - a. there are documented recurrent or chronic rashes, infections, cellulitis, or non-healing ulcers, that do not respond to conventional treatment (for example, dressing changes; topical, oral or systemic antibiotics, corticosteroids or antifungals) for a period of 3 months; **or**
    - b. there is documented difficulty with ambulation and interference with the activities of daily living; **and**
  3. Symptoms or functional impairment persists despite significant\* weight loss which has been stable for at least 3 months or well-documented attempts at weight loss (medically supervised diet or bariatric surgery) have been unsuccessful; **and**
  4. If the individual has had bariatric surgery, he/she is at least 18 months post-operative or has documented stable weight for at least 3 months.

\*Significant weight loss varies based on the individual clinical circumstances and may be documented when the individual:

  - a. Reaches a body mass index (BMI) less than or equal to 30 kg/m<sup>2</sup>; **or**
  - b. Has documented at least a 100 pound weight loss; **or**
  - c. Has achieved a weight loss which is 40% or greater of the excess body weight that was present prior to the individual's weight loss program or surgical intervention.
- B. Panniculectomy is considered **medically necessary** as an adjunct to a medically necessary surgery when needed for exposure in extraordinary circumstances.

### Not Medically Necessary:

- A. Panniculectomy is considered **not medically necessary** when the criteria above are not met.
- B. Panniculectomy is considered **not medically necessary** as an adjunct to other medically necessary procedures, including, but not limited to, hysterectomy, or incisional or ventral hernia repair unless the criteria above are met.
- C. Panniculectomy or abdominoplasty, with or without diastasis recti repair, for the treatment of back pain is considered **not medically necessary**.

### Cosmetic and Not Medically Necessary:

- A. Liposuction is considered **cosmetic and not medically necessary** when used for the removal of excess abdominal fat.
- B. Abdominoplasty when done to remove excess skin or fat with or without tightening of the underlying muscles is considered **cosmetic and not medically necessary**.
- C. Repair of diastasis recti is considered **cosmetic and not medically necessary**.

## Coding

The following codes for treatments and procedures applicable to this guideline are included below for informational purposes. Inclusion or exclusion of a procedure, diagnosis or device code(s) does not constitute or imply member coverage or provider reimbursement policy. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

### Panniculectomy

#### When services may be Medically Necessary when criteria are met:

#### CPT

15830                      Excision, excessive skin and subcutaneous tissue (includes lipectomy); abdomen, infraumbilical panniculectomy

#### ICD-10 Procedure

0HB7XZZ                      For the following codes when described as panniculectomy:  
Excision of abdomen skin, external approach  
0J080ZZ                      Alteration of abdomen subcutaneous tissue and fascia, open approach

0WBF0ZZ      Excision of abdominal wall, open approach

#### ICD-10 Diagnosis

All diagnoses

#### When services are Not Medically Necessary:

For the procedure codes listed above when criteria are not met or for situations designated in the Clinical Indications section as not medically necessary.

#### Abdominoplasty, liposuction

#### When services are Not Medically Necessary or Cosmetic and Not Medically Necessary:

For the following procedure codes, or when the code describes a procedure designated in the Clinical Indications section as not medically necessary or cosmetic and not medically necessary.

#### CPT

15847	Excision, excessive skin and subcutaneous tissue (includes lipectomy); abdomen (eg, abdominoplasty) (includes umbilical transposition and fascial plication)
15877	Suction assisted lipectomy; trunk [when specified as abdominal liposuction]
17999	Unlisted procedure, skin, mucous membrane and subcutaneous tissue [when specified as other abdominoplasty, excision excessive skin and subcutaneous tissue, including lipectomy, of abdomen]

#### ICD-10 Procedure

0J080ZZ	Alteration of abdomen subcutaneous tissue and fascia, open approach [when specified as other abdominoplasty, excision excessive skin and subcutaneous tissue, including lipectomy]
0J083ZZ	Alteration of abdomen subcutaneous tissue and fascia, percutaneous approach
0W0F07Z-0W0F0ZZ	Alteration of abdominal wall with/without tissue substitute, open approach [includes codes 0W0F07Z, 0W0F0JZ, 0W0F0KZ, 0W0F0ZZ]
0W0F37Z-0W0F3ZZ	Alteration of abdominal wall with/without tissue substitute, percutaneous approach [includes codes 0W0F37Z, 0W0F3JZ, 0W0F3KZ, 0W0F3ZZ]
0W0F47Z-0W0F4ZZ	Alteration of abdominal wall with/without tissue substitute, percutaneous endoscopic approach [includes codes 0W0F47Z, 0W0F4JZ, 0W0F4KZ, 0W0F4ZZ]

#### ICD-10 Diagnosis

All diagnoses

#### Repair of diastasis recti

#### When services are Not Medically Necessary or Cosmetic and Not Medically Necessary:

For the following procedure codes, or when the code describes a procedure designated in the Clinical Indications section as not medically necessary or cosmetic and not medically necessary.

#### CPT

22999	Unlisted procedure, abdomen, musculoskeletal system [when specified as repair of diastasis recti]
-------	---

#### ICD-10 Procedure

0KQK0ZZ-0KQK4ZZ	Repair right abdomen muscle [by approach; includes codes 0KQK0ZZ, 0KQK3ZZ, 0KQK4ZZ]
0KQL0ZZ-0KQL4ZZ	Repair left abdomen muscle [by approach; includes codes 0KQL0ZZ, 0KQL3ZZ, 0KQL4ZZ]

#### ICD-10 Diagnosis

	For the following diagnoses when specified as diastasis recti:
M62.00	Separation of muscle (nontraumatic), unspecified site
M62.08	Separation of muscle (nontraumatic), other site
O71.89	Other specified obstetric trauma
Q79.59	Other congenital malformations of abdominal wall

## Discussion/General Information

#### Panniculectomy

The current medical evidence regarding panniculectomy consists mostly of individual case reports, review articles and a limited number of controlled trials. However, there is adequate clinical opinion to support the use of this procedure in some circumstances where an individual's health is compromised.

Early studies by Matory (1994) and Vastine (1999) demonstrated a direct relationship between BMI and operative risk with abdominal surgery and abdominoplasty in obese individuals. In a retrospective cohort series of individuals who underwent post-bariatric panniculectomy (n=126), the only factor that independently predicted postoperative complications after panniculectomy was pre-panniculectomy BMI (Arthurs, 2007). Those with a BMI greater than 25 kg/m<sup>2</sup> were at nearly three times the risk of postoperative wound complications. Although those who experienced a plateau in weight loss at a BMI of 30-35 kg/m<sup>2</sup> did have the largest functional improvement from a panniculectomy, they also experienced the highest risk postoperatively. The average weight loss following bariatric surgery prior to panniculectomy was 116 ± 35 lbs. A limitation of this study was its retrospective design and sample size.

Acarturk (2004) compared the surgical outcomes of panniculectomy following bariatric surgery in another retrospective series of 123 participants (mean age 44.5 years). The outcomes of 21 participants with panniculectomy performed at the time of bariatric surgery were compared with the surgical outcomes of 102 participants who waited 17 ± 11 months to undergo panniculectomy. Overall, individuals who had panniculectomy simultaneously with bariatric surgery experienced more complications. Wound infections were 48% versus 16%; wound dehiscence 33% versus 13%; and there was a higher incidence (24% versus 0 %) of postoperative respiratory distress in individuals with the combined procedures. There were 3 postoperative deaths in the combined procedure cohort and none in the group that delayed panniculectomy until an average weight loss of 126 ± 59 lbs was achieved. The authors concluded that an initial period of substantial weight loss prior to the procedure results in a safer and more effective panniculectomy procedure.

The American Society of Plastic Surgeons (ASPS) Practice Parameter for Surgical Treatment of Skin Redundancy for Obese and

Massive Weight Loss Patients (2007b) recommends that body contouring surgery, including panniculectomy, be performed only after an individual maintains a stable weight for 2 to 6 months. For individuals who are post-bariatric surgery, this is reported to occur 12-18 months after surgery when the BMI has reached the 25 kg/m<sup>2</sup> to 30 kg/m<sup>2</sup> range (Mechanick, 2013; Rubin, 2004). If performed prematurely, a potential exists for a second panniculus to develop once additional weight loss has occurred and the risks of postoperative complications are increased. Weight loss and BMI are important when considering panniculectomy and a significant amount of weight loss may not bring the BMI of an individual to less than 30 kg/m<sup>2</sup>; however, a panniculectomy may still be necessary (Arthurs, 2007). The American Society for Metabolic and Bariatric Surgery Consensus statement states weight loss can vary from about 25% to 70% of an individual's excess body weight depending on the type of bariatric surgery that is performed (Buchwald, 2005).

A study by Zemlyak and colleagues (2012) reported on a retrospective review of individuals who had panniculectomy alone versus individuals who had panniculectomy and simultaneous ventral hernia repair. There were 143 participants in the panniculectomy/ventral hernia repair group and 42 participants in the panniculectomy group. The rates for incisional complications and interventions between the two groups were not statistically significant. However, after controlling for age, gender, BMI, subcutaneous use of talc, and intraoperative pulse-a-vac irrigation in the multivariate regression analysis, the group that had both panniculectomy and ventral hernia repair was more likely to develop wound cellulitis. The authors note that while panniculectomy with ventral hernia repair reduces the stress on the hernia repair and potentially decreases the recurrence rate, this potential advantage remains to be proven in robust comparative studies.

Fischer and colleagues (2014) conducted a large retrospective database analysis to assess the additional risk of ventral hernia repair (VHR) and panniculectomy (PAN) compared with hernia repair alone (n=55,537) using the American College of Surgeons National Surgical Quality Improvement Program (ACS-NSQIP) data sets. To account for potential selection bias given the non-randomized assignment of concurrent panniculectomy and the retrospective study design, propensity scores were used which yielded two comparable groups, VHR (n=1250) and VHR+PAN (n=1250). The study authors found that individuals who underwent the combined procedure were at significantly higher risk for wound complications (p<0.001), venous thromboembolism (p=0.044), reoperation (p<0.001) and overall medical morbidity (p<0.001). Two notable limitations of this study include that the ACS-NSQIP dataset only includes 30-day outcomes, precluding analysis of long-term differences in the two study groups. Secondly, the dataset did not include details on the type of panniculectomy skin resection or wound closure techniques, therefore propensity matching, and exploratory analysis of these potentially confounding variables was not possible. Nonetheless, at 30-day follow-up in this large retrospective cohort, outcomes of panniculectomy performed with a concurrent ventral hernia repair appear to result in a significant increase in morbidity compared to VHR alone.

Giordano and colleagues (2017) published a retrospective study based on a prospectively maintained database of all consecutive midline abdominal wall reconstructions for an abdominal wall hernia or oncologic defect performed at a single site from 2005-2015. Of 548 consecutive surgeries, 305 individuals (56%) underwent abdominal wall reconstruction alone and 243 (44%) underwent abdominal wall reconstruction with concurrent panniculectomy. The mean follow-up period was 30 months. Prior to propensity matching, individuals with the combined procedure also had a higher number of previous abdominal surgeries and a larger mean abdominal wall defect size. After propensity matching, there were significantly higher incidences of fat necrosis, and surgical site abscess but no significant difference in hernia recurrence at follow-up. Abdominal wall reconstruction with concurrent panniculectomy was associated with higher wound morbidity with no difference in hernia recurrence rates in follow-up.

Derickson (2018) published results from retrospective review of all post-bariatric surgery cases who underwent panniculectomy over a 10-year period (n=706). The overall rate of complication was 56%: dehiscence (24%), surgical site infection (22%), seroma (18%), and post-operative bleeding (5%). A total of 12% of individuals necessitated a return to the operating room. The study demonstrated a high morbidity for post-bariatric panniculectomy and authors noted higher BMI, higher ASA class, and the use of fleur-de-lis incision were particularly associated with worse outcomes.

Nag and colleagues (2021) published results from another systematic review conducted by ACS-NSQIP to determine the benefit, if any, of adding panniculectomy to gynecologic surgery in obese and morbidly obese individuals. In total, 296 individuals were identified from the NSQIP database who fit the search criteria. A statistically significant association was found between the concomitant procedures and adverse outcomes, including superficial infection, wound infection, pulmonary embolism, sepsis, return to operating room, length of operation and length of stay. Furthermore, there was no significant benefit identified across the studies. Panniculectomy alone or with other abdominal surgical procedures, such as incisional or ventral hernia repair, or hysterectomy, is not clinically appropriate or an effective treatment of obesity. Recent meta-analyses have published mixed results of co-surgical procedures, but the studies lack documentation of a medical indication for removal of the pannus (Prodromidou, 2020; Rasmussen, 2017; Sosin, 2020). Although it has been suggested that the presence of a large overhanging panniculus may interfere with the surgery or compromise post-operative recovery, the presence of a pannus alone is not a medical condition which warrants surgical intervention. Removal of a pannus, for reasons other than those in the criteria for medical necessity is therefore considered cosmetic and not medically necessary.

#### *Abdominoplasty*

The literature addressing abdominoplasty and surgical repair of diastasis recti confirms the cosmetic benefits of these procedures. However, improvements in physical functioning, cessation of back pain, and other positive health outcomes have not been demonstrated. Carloni and colleagues conducted a systematic-review (2016) and confirmed that the quality of evidence surrounding abdominoplasty remains low and no standardization of surgical approaches has been established. Winocour (2015) reported results of a study which included 25,478 abdominoplasties and found high complication rates, compared to other cosmetic procedures, especially when abdominoplasty was combined with other procedures. Massenburg (2015) reported outcomes from 2946 abdominoplasties and found 8.5% of subjects were readmitted due to complications and 5% required revision surgery. Salari and colleagues (2021) conducted a systematic-review and meta-analysis to characterize the global prevalence of seroma following abdominoplasty and found the global prevalence following the procedure approaching 11% (95% CI, 9.3-3.6%). At this time, the evidence does not support abdominoplasty when done to remove excess abdominal skin or fat, with or without tightening lax anterior abdominal wall muscles, as an effective treatment for any medical condition, though it is an effective cosmetic procedure (ASPS Practice Parameter, 2007b).

Surgical procedures to correct diastasis recti are not effective for alleviating back pain or other non-cosmetic conditions. There is insufficient evidence to support the use of surgical procedures to correct diastasis recti for purposes other than cosmetic.

Similarly, the use of liposuction has been shown to produce cosmetic benefits in terms of appearance and body contour, however, liposuction has not been shown to be an effective treatment of obesity or other medical conditions and has been associated with significant complications, including death.

## Definitions

**Abdominoplasty:** A procedure involving the removal of excess abdominal skin and fat with or without tightening lax anterior abdominal wall muscles and with or without repositioning or reconstruction of the navel.

**Bariatric surgery:** A variety of surgical procedures designed to treat obesity by either reconstructing the stomach or intestines or placing restrictive devices in or on the digestive tract.

**Cellulitis:** A diffuse, spreading inflammation of the deep tissues under the skin, and on occasion muscle, which may be associated with abscess formation.

**Diastasis recti:** A condition characterized by a separation between the left and right side of the rectus abdominis, which is the muscle covering the front surface of the chest (abdomen). A diastasis recti appears as a ridge running down the midline of the abdomen from the bottom of the breastbone to the navel.

**Hysterectomy:** Surgical removal of the uterus.

**Incisional hernia:** A condition where tissues or organs are able to push through a surgical incision or scar.

**Intertrigo:** An inflammation of the top layers of skin caused by moisture, bacteria, or fungi in the folds of the skin.

**Liposuction:** A surgical procedure designed to remove fat from under the skin via a suction device.

**Panniculectomy:** A procedure designed to remove fatty tissue and excess skin (panniculus) from the lower to middle portions of the abdomen.

**Pubis:** A part of the pelvic bone that is located in the groin, also called the pubic bone.

## References

### Peer Reviewed Publications:

1. Acarturk TO, Wachtman G, Heil B, et al. Panniculectomy as an adjuvant to bariatric surgery. *Ann Plast Surg.* 2004; 53(4):360-366.
2. Ali B, Petersen TR, McKee RG. Perioperative risk stratification model for readmission after panniculectomy. *Plast Reconstr Surg.* 2022; 150(1):181-188.
3. Arthurs ZM, Cuadrado D, Sohn V, et al. Post-bariatric panniculectomy: pre-panniculectomy body mass index impacts the complication profile. *Am J Surg.* 2007; 193(5):567-570.
4. Blomfield PI, Le T, Allen DG, Planner RS. Panniculectomy: a useful technique for the obese patient undergoing gynecological surgery. *Gynecol Oncol.* 1998; 70(1):80-86.
5. Carloni R, De Runz, Chaput B et al. Circumferential contouring of the lower trunk: indications, operative techniques, and outcomes-a systematic review. *Aesthetic Plast Surg.* 2016; 40(5):652-668.
6. Cassar K, Munro A. Surgical treatment of incisional hernia. *Br J Surg.* 2002; 89(5):534-545.
7. Coriddi MR, Koltz PF, Chen R, Gusenoff JA. Changes in quality of life and functional status following abdominal contouring in the massive weight loss population. *Plast Reconstr Surg.* 2011; 128(2):520-526.
8. Derickson M, Phillips C, Barron M, et al. Panniculectomy after bariatric surgical weight loss: analysis of complications and modifiable risk factors. *Am J Surg.* 2018; 215(5):887-890.
9. Fischer JP, Tuggle CT, Wes AM, Lovach SJ. Concurrent panniculectomy with open ventral hernia repair has added risk versus ventral hernia repair: an analysis of the ACS-NSQIP database. *J Plast Reconstr Aesthet Surg.* 2014; 67(5):693-701.
10. Giordano S, Garvey PB, Baumann DP, et al. Concomitant panniculectomy affects wound morbidity but not hernia recurrence rates in abdominal wall reconstruction: A propensity score analysis. *Plast Reconstr Surg.* 2017; 140(6):1263-1273.
11. Harth KC, Blatnik JA, Rosen MJ. Optimum repair for massive ventral hernias in the morbidly obese patient--is panniculectomy helpful? *Am J Surg.* 2011; 201(3):396-400.
12. Hopkins MP, Shriner AM, Parker MG, Scott L. Panniculectomy at the time of gynecologic surgery in morbidly obese patients. *Am J Obstet Gynecol.* 2000; 182(6):1502-1505.
13. Hughes KC. Ventral hernia repair with simultaneous panniculectomy. *Ann Surg.* 1996; 62(8):678-681.
14. Iavazzo C, Psomiadou V, Fotiou A, et al. Concurrent gynecologic surgery and panniculectomy in morbidly obese women with gynecologic cancer, a single-center experience. *Arch Gynecol Obstet.* 2021; 304(5):1271-1278.
15. Kantar RS, Rifkin WJ, Wilson SC, et al. Abdominal panniculectomy: determining the impact of diabetes on complications and risk factors for adverse events. *Plast Reconstr Surg.* 2018; 142(4):462e-471e.
16. Lesko RP, Cheah MA, Sarmiento S, et al. Postoperative complications of panniculectomy and abdominoplasty: A retrospective review. *Ann Plast Surg.* 2020; 85(3):285-289.
17. Massenburg BB, Sanati-Mehrziy P, Jablonka EM, Taub PJ. Risk factors for readmission and adverse outcomes in abdominoplasty. *Plast Reconstr Surg.* 2015; 136(5):968-977.
18. Matarasso A, Wallach SG, Rankin M, Galiano RD. Secondary abdominal contour surgery: a review of early and late reoperative surgery. *Plast Reconstr Surg.* 2005; 115(2):627-632.
19. Matory WE, O'Sullivan J, Fudem G, Dunn R. Abdominal surgery in patients with severe morbid obesity. *Plast Reconstr Surg.* 1994; 94(7):976-987.
20. Nag S, Patel T, Gaughan JP, Bonawitz SC. Panniculectomy performed in conjunction with gynecologic surgery in obese and morbidly obese patients: A National Surgical Quality Improvement Program Analysis and systematic review of the literature. *Ann Plast Surg.* 2021; 87(5):600-605. Nahas FX, Augusto SM, Ghelfond C. Should diastasis recti be corrected? *Aesth Plas Surg.* 1997; 21(4):285-289. Pearl ML, Valea FA, Disilvestro PA, Chalas E. Panniculectomy in morbidly obese gynecologic oncology patients. *Int J Surg Investig.* 2000; 2(1):59-64. Powell JL. Panniculectomy to facilitate gynecologic surgery in morbidly obese women. *Obstet Gynecol.* 1999 94(4):528-531.
21. Prodromidou A, Iavazzo C, Psomiadou V, et al. Safety and efficacy of synchronous panniculectomy and endometrial cancer surgery in obese patients: a systematic review of the literature and meta-analysis of postoperative complications. *J Turk Ger Gynecol Assoc.* 2020; 21(4):279-286.
22. Rasmussen RW, Patibandla JR, Hopkins MP. Evaluation of indicated non-cosmetic panniculectomy at time of gynecologic surgery. *Int J Gynaecol Obstet.* 2017; 138(2):207-211.
23. Rubin JP, Nguyen V, Schwentker A. Perioperative management of the post-gastric-bypass patient presenting for body contour surgery. *Clin Plast Surg.* 2004; 31(4):601-610.
24. Salari N, Fatahi B, Bartina Y, et al. The global prevalence of seroma after abdominoplasty: a systematic review and meta-analysis. *Aesthetic Plast Surg.* 2021; 45(6):2821-2836.
25. Sosin M, Termanini KM, Black CK, et al. Simultaneous ventral hernia repair and panniculectomy: A systematic review and meta-analysis of outcomes. *Plast Reconstr Surg.* 2020; 145(4):1059-1067.
26. Staalesen T, Olsén MF, Elander A. The effect of abdominoplasty and outcome of rectus fascia plication on health-related quality of life in post-bariatric surgery patients. *Plast Reconstr Surg.* 2015; 136(6):750e-761e. Tillmanns TD, Kamelle SA,

- Abudayyeh I, et al. Panniculectomy with simultaneous gynecologic oncology surgery. *Gynecol Oncol*. 2001; 83(3):518-522.
27. Vastine VL, Morgan RF, Williams GS, et al. Wound complications of abdominoplasty in obese patients. *Plast Surg*. 1999; 42(1):34-39.
  28. Warren JA, Epps M, Debrux C, et al. Surgical site occurrences of simultaneous panniculectomy and incisional hernia repair. *Am Surg*. 2015; 81(8):764-769.
  29. Winocour J, Gupta V, Ramirez JR, et al. Abdominoplasty: risk factors, complication rates, and safety of combined procedures. *Plast Reconstr Surg*. 2015; 136(5):597e-606e.
  30. Zannis J, Wood BC, Griffin LP, et al. Outcome study of the surgical management of panniculitis. *Ann Plast Surg*. 2012; 68(2):194-197.
  31. Zemlyak AY, Colavita PD, El Djouzi S, et al. Comparative study of wound complications: isolated panniculectomy versus panniculectomy combined with ventral hernia repair. *J Surg Res*. 2012; 177(2):387-391.

#### Government Agency, Medical Society, and Other Authoritative Publications:

1. American Society of Plastic and Reconstructive Surgeons (ASPS). ASPS recommended insurance coverage criteria for third-party payers: Surgical treatment of skin redundancy for obese and massive weight loss patients. 2007a. Available at: <https://www.plasticsurgery.org/Documents/medical-professionals/health-policy/insurance/Surgical-Treatment-of-Skin-Redundancy-Following.pdf>. Accessed on January 03, 2023.
2. American Society of Plastic and Reconstructive Surgeons (ASPS). Practice parameter for surgical treatment of skin redundancy for obese and massive weight loss patients. 2007b. Available at: <http://www.plasticsurgery.org/Documents/medical-professionals/health-policy/evidence-practice/Surgical-Treatment-of-Skin-Redundancy-Following-Massive-Weight-Loss.pdf>. Accessed on January 03, 2023.
3. Buchwald H; Consensus Conference Panel. Consensus conference statement bariatric surgery for morbid obesity: health implications for patients, health professionals, and third-party payers. *Surg Obes Relat Dis*. 2005; 1(3):371-381.
4. Coleman WP, Glogau RG, Klein JA, et al. American Academy of Dermatology Guidelines/Outcomes Committee. Guidelines of care for liposuction. *J Am Acad Dermatol*. 2001; 45(3):438-447.
5. Mechanick JL, Apovian C, Brethauer S, et al. Clinical Practice Guidelines for the Perioperative Nutrition, Metabolic, and Nonsurgical Support of Patients Undergoing Bariatric Procedures -2019 Update: Cosponsored by American Association of Clinical Endocrinologists/American College Of Endocrinology, The Obesity Society, American Society for Metabolic and Bariatric Surgery, Obesity Medicine Association, and American Society of Anesthesiologists – Executive Summary. 2019; 12: 1346-1359.

#### Websites for Additional Information

1. National Institutes of Health. National Heart, Lung, and Blood Institute. BMI calculator. Available at: [https://www.nhlbi.nih.gov/health/educational/lose\\_wt/BMI/bmicalc.htm](https://www.nhlbi.nih.gov/health/educational/lose_wt/BMI/bmicalc.htm). Accessed on January 03, 2023.
2. National Library of Medicine. Medical Encyclopedia: Diastasis recti. Available at: <http://www.nlm.nih.gov/medlineplus/ency/article/001602.htm>. Accessed on January 03, 2023.

#### History

Status	Date	Action
Reviewed	02/15/2024	Medical Policy & Technology Assessment Committee (MPTAC) review. Updated Discussion/General Information, References and Website sections.
Reviewed	02/16/2023	MPTAC review. Updated Discussion/General Information, References and Website sections.
Reviewed	02/17/2022	MPTAC review. Updated Discussion/General Information, References and Website sections. Updated Coding section; removed CPT anesthesia code 00802.
Reviewed	02/11/2021	MPTAC review. Revised MN definition text in the Description section. Updated Discussion/General Information, References and Website sections. Reformatted Coding section.
Reviewed	02/20/2020	MPTAC review. Updated References and Website sections.
New	03/21/2019	MPTAC review. Initial document development. Moved content of SURG.00048 Panniculectomy and Abdominoplasty to a new clinical utilization management guideline document with the same title. In the Cosmetic and Not Medically Necessary position statement section: (1) revised bullet "A" to indicate that liposuction is considered cosmetic and not medically necessary when used for the removal of excess abdominal fat; (2) revised bullet "C" by removing the words "for all indications".

Federal and State law, as well as contract language, and Medical Policy take precedence over Clinical UM Guidelines. We reserve the right to review and update Clinical UM Guidelines periodically. Clinical guidelines approved by the Medical Policy & Technology Assessment Committee are available for general adoption by plans or lines of business for consistent review of the medical necessity of services related to the clinical guideline when the plan performs utilization review for the subject. Due to variances in utilization patterns, each plan may choose whether to adopt a particular Clinical UM Guideline. To determine if review is required for this Clinical UM Guideline, please contact the customer service number on the member's card.

Alternatively, commercial or FEP plans or lines of business which determine there is not a need to adopt the guideline to review services generally across all providers delivering services to Plan's or line of business's members may instead use the clinical guideline for provider education and/or to review the medical necessity of services for any provider who has been notified that his/her/its claims will be reviewed for medical necessity due to billing practices or claims that are not consistent with other providers, in terms of frequency or in some other manner.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.