

PNAS PNAS PNAS PNAS PNAS

<sup>1</sup>Innate Immunity Unit, Immunology Department, Institut Pasteur, 75724 Paris, France; <sup>2</sup>Institut National de la Santé et de la Recherche Médicale (INSERM) U668, 75724 Paris, France; <sup>3</sup>Part of the Human Vaccine Consortium, "Grand Challenges in Global Health #4: devise reliable testing systems for new vaccines" (<http://www.hv-consortium.org/>); <sup>4</sup>Department of Cell Biology and Histology, Academic Medical Center, University of Amsterdam, 1105, Amsterdam, The Netherlands; <sup>5</sup>Lymphocyte Development Unit, Institut Pasteur, 75724 Paris, France; <sup>6</sup>INSERM U892, Groupe de Recherche Cytokines et Récepteurs, Institut de Biologie, University of Nantes, F-44093 Nantes, France; <sup>7</sup>Department of Vaccinology and Applied Microbiology, Helmholtz Centre for Infection Research, D-38124 Braunschweig, Germany; <sup>8</sup>INSERM U845, Centre de Recherche Croissance et Signalisation, Université Paris Descartes, Hôpital Necker Enfants Malades, 75015 Paris, France; <sup>9</sup>INSERM Laboratoire de Virologie, Hôpital Tenon, Assistance Publique-Hopitaux de Paris, 75020 Paris, France

Cytokine immunotherapies targeting T lymphocytes are attractive clinical interventions against viruses and tumors. In the mouse, the homeostasis of memory  $\alpha/\beta$  CD8<sup>+</sup> T cells and natural killer (NK) cells is significantly improved with increased IL-15 bioavailability. In contrast, the role of “transpresented” IL-15 on human T-cell development and homeostasis *in vivo* is unknown. We found that both CD8 and CD4 T cells in human immune system (HIS) mice are highly sensitive to transpresented IL-15 *in vivo*, with both naïve (CD62L<sup>+</sup>CD45RA<sup>+</sup>) and memory phenotype (CD62L<sup>-</sup>CD45RO<sup>+</sup>) subsets being significantly increased following IL-15 “boosting.” The unexpected global improvement in human T-cell homeostasis involved enhanced proliferation and survival of both naïve and memory phenotype peripheral T cells, which potentiated B-cell responses by increasing the frequency of antigen-specific responses following immunization. Transpresented IL-15 did not modify T-cell activation patterns or alter the global T-cell receptor (TCR) repertoire diversity. Our results indicate an unexpected effect of IL-15 on human T cells *in vivo*, in particular on CD4<sup>+</sup> T cells. As IL-15 promotes human peripheral T-cell homeostasis and increases the frequency of neutralizing antibody responses in HIS mice, IL-15 immunotherapy could be envisaged as a unique approach to improve vaccine responses in the clinical setting.

T-cell pool, and regulation of activated effector and memory T-cell compartments (6). Several signals have been implicated in controlling T-cell homeostasis, including those emanating from the T-cell receptor (TCR) following interactions with self-peptide + major histocompatibility complex (pMHC) and those induced by growth factors, including cytokines (6). The common cytokine receptor gamma chain ( $\gamma_c$ ) family of cytokines (which comprises IL-2, IL-4, IL-7, IL-9, IL-15, and IL-21) in particular have been demonstrated to play a role in T-cell homeostasis in mammals (7). Humans and mice possessing mutations in genes encoding the  $\gamma_c$ , Jak3 (both critical for signal transduction following binding  $\gamma_c$  cytokines) or the alpha chain of the IL-7 receptor (IL-7R $\alpha$ ), display a severe block in T-cell development and resulting severe combined immunodeficiency (8, 9). The  $\gamma_c$ -dependent cytokine IL-15 is unusual because its bioactive form is a functional complex associated with the IL-15R $\alpha$  chain. Thus, cells expressing IL-15 such as monocytes, dendritic cells, and stromal cells must also coexpress the IL-15R $\alpha$  to “transpresent” IL-15 to IL-15-responsive cells (that express the IL-2R $\beta$ / $\gamma_c$  complex). Accordingly, both IL-15 and IL-15R $\alpha$  are up-regulated on myeloid cells following inflammation, thereby increasing IL-15 bioavailability (10–12).

We demonstrated that transpresented murine IL-15 inefficiently triggered human natural killer (NK) cells *in vitro* and *in vivo* providing an explanation for the poor human NK cell reconstitution in BALB/c Rag2<sup>-/-</sup>  $\gamma_c$ <sup>-/-</sup> HIS mice (3). Exogenous administration of a potent human IL-15R agonist (referred to as RLI, consisting of human IL-15 covalently linked to an extended human IL-15R $\alpha$  "sushi" domain thus mimicking IL-15 transpresentation) (13–15) was sufficient to restore human NK cell development in HIS mice (3). Whereas memory CD8<sup>+</sup> T cells in mice are highly responsive to exogenous IL-15 (6, 11–14), naïve CD4<sup>+</sup> and CD8<sup>+</sup> T cells are not thought to require IL-15 for normal homeostasis (6, 11–14). However, *in vivo* effects of human

This article contains supporting information online at [www.pnas.org/lookup/suppl/doi:10.1073/pnas.1019167108/-/DCSupplemental](http://www.pnas.org/lookup/suppl/doi:10.1073/pnas.1019167108/-/DCSupplemental).


IL-15 on human T cells have not been studied, and it remained possible that the poor reactivity of human T cells to mouse IL-15 might also contribute to the low human T-cell reconstitution in the HIS mouse model. Here we show that human IL-15 transpresentation increases human T-cell reconstitution and the frequency of T-cell-dependent antibody responses in HIS mice. These studies provide a first preclinical trial of transpresented human IL-15 on human T cells in vivo and indicate that increased IL-15 bioavailability globally boosts human naïve and memory T-cell homeostasis in this humanized mouse model. Our findings offer a unique approach to study human T-cell immune responses in vivo and suggest that IL-15 immunotherapy may be useful to promote global T-cell reconstitution in humans.

## Results

**Improved Development of Human CD4<sup>+</sup> and CD8<sup>+</sup> T Cells in HIS Mice Receiving RLI.** We and others have recently reported that human fetal liver HSCs (CD34<sup>+</sup>CD38<sup>-</sup>) engrafted into newborn BALB/c Rag2<sup>-/-</sup>γC<sup>-/-</sup> mice develop into mature myeloid and lymphoid cells (1–5). We used this approach to investigate the effect of

transpresented human IL-15 on human T-cell development in vivo. Eight weeks after HSC engraftment, HIS mice were injected with the potent human IL-15/IL-15Rα agonist, RLI (13–15) (Fig. 1A). Administration of RLI resulted in a significant increase of human hematopoietic cells in the thymus and spleen and a specific increase in CD3<sup>+</sup> cells in the bone marrow (Fig. 1B). CD8<sup>+</sup> T cells were significantly augmented in the spleen ( $P = 0.025$ ), blood ( $P = 0.027$ ), and thymus ( $P = 0.033$ ) following RLI treatment in HIS mice, whereas CD4<sup>+</sup> T cells were also augmented in spleen ( $P = 0.017$ ) and lymph node ( $P = 0.025$ ) (Fig. 1B and C). In addition, NK cells were also augmented as previously reported (3), whereas B-cell numbers were not significantly altered (Fig. S1A). Thus, the increase in total human hematopoietic cells in the thymus ( $P = 0.0005$ ) and spleen ( $P = 0.002$ ) following RLI treatment is attributed to significant increases in total T-cell numbers (Fig. 1B).

**RLI Promotes Proliferation of Naïve and Memory Phenotype Peripheral CD4<sup>+</sup> and CD8<sup>+</sup> T Cells in HIS Mice.** Because RLI was effective in increasing both CD4<sup>+</sup> and CD8<sup>+</sup> T cells, we next



**Fig. 1.** RLI enhances human CD4<sup>+</sup> and CD8<sup>+</sup> T-cell development and reconstitution of lymphoid organs in HIS mice. (A) Experimental scheme. Newborn Rag2<sup>-/-</sup>γC<sup>-/-</sup> mice were irradiated with 3.3 Gy injected intrahepatically (i.h.) with  $5 \times 10^4$  CD34<sup>+</sup>CD38<sup>-</sup> human fetal liver cells. At 8, 9, 10, and 11 wk of age, HIS mice were injected intraperitoneally (i.p.) with 2.5 μg IL-15-IL-15Rα fusion protein (RLI) or PBS. Mice were killed and analyzed at 12 wk. (B) Lymphoid organs and (C) peripheral blood from HIS mice were analyzed for either total human hematopoietic cell (human CD45; hCD45<sup>+</sup>) and human T-cell (hCD45<sup>+</sup>CD3<sup>+</sup> and CD4<sup>+</sup> or CD8<sup>+</sup>) reconstitution by flow cytometry, and cellularity was enumerated. hCD45<sup>+</sup> (%) is the percentage of total cells, whereas CD3<sup>+</sup>, CD3<sup>+</sup>CD4<sup>+</sup>, and CD3<sup>+</sup>CD8<sup>+</sup> (%) are the percentages within the hCD45<sup>+</sup> population. Values represent mean  $\pm$  SEM of nine mice per group.



HIS mice, the TCR repertoire of the peripheral T-cell pool appears diverse, with T cells bearing TCRs from all V $\beta$  family members and demonstrating varying CDR3 lengths (Fig. S4A and B). Importantly, RLI administration did not skew the TCR repertoire compared with control HIS mice, suggesting that IL-15 globally boosts the available broad TCR pool. This is a valuable observation considering our model of humanized mice is becoming more popular for the study of T-cell immune responses and suggests that an extensive TCR repertoire is generated providing a cellular substrate that could react against a large number of peptide antigens.

Using CDR3 immunoscope analysis of  $\gamma$  and  $\delta$  variable chains we found that most human  $\gamma/\delta$  T cells found in the periphery of HIS mice use V $\delta$ 2 (similar to human peripheral blood mononuclear cells, PBMCs); this preferential use is unaffected by RLI treatment (Fig. S4C and D). Interestingly, we observed two large populations of V $\gamma$ 8<sup>+</sup> and V $\gamma$ 9<sup>+</sup> T cells that each represent between 40% and 60% of total  $\gamma/\delta$  T cells and who pair almost exclusively with V $\delta$ 2, although we observed occasional use of V $\delta$ 3, -5, and -8 (Fig. S4C). In vivo administration of RLI did not influence the relative  $\gamma$  and  $\delta$  chain use nor the variability in CDR3 length for any given V $\gamma$  or V $\delta$  chains (Fig. S4D).

**Improved Development of Human CD4<sup>+</sup> and CD8<sup>+</sup> T Cells in HIS Mice Receiving RLI Results in Improved Humoral Responses Following Immunization.** We next investigated whether improved human T-cell reconstitution impacted on immune functions in HIS mice. Because antigen-specific T-cell responses, particularly cytotoxic T-cell responses (cytotoxic T lymphocytes, CTLs) remain poorly elicited in HIS mice, we characterized T-dependent antigen-specific B responses that can be evoked following vaccination of HIS mice. To this end, HIS mice cohorts were immunized with commercial hepatitis B virus (HBV) and tetanus toxoid (TT) vaccines. Although serum IgM levels remained unchanged following RLI treatment, significant increases in total IgG levels were observed in RLI-treated immunized HIS mice (Fig. 3A). Using a diagnostic ELISA, we were able to detect anti-TT-specific IgG and more frequently anti-HBV surface antigen (HbsAg)-specific Ig in the sera of immunized HIS mice (Fig. 3B and C). Interestingly, although RLI treatment did not modify the concentration of antigen-specific antibody in mice that mounted an antigen-specific Ig response (Fig. 3B and C), the frequency of “responders” (HIS mice with either >0.5 IU/mL anti-TT-specific IgG or >7 IU/L anti-HbsAg-specific Ig) was significantly enhanced following RLI treatment ( $P < 0.05$ ;  $\chi^2 = 3.85$ ) (Fig. 3D). Thus, the improvement in human T-cell homeostasis in RLI-treated HIS


mice increased the likelihood of inducing antigen-specific humoral responses following immunization.

## Discussion

HIS mouse models have been continually improved over the past 30 y and are now at a stage where both antibody and cellular human immune responses can be elicited (16–19). Given that the immune system in this HIS mouse model is skewed toward B-cell development, there is a clear need to improve T lymphopoiesis and homeostasis (1, 2). We recently reported a beneficial effect on human thymopoiesis following human IL-7 treatment in HIS mice; however, peripheral T-cell numbers remained unchanged despite transient increases in thymocyte numbers, suggesting that other factors are involved (5). Given that human lymphocytes are poorly triggered by murine IL-15 (3), we hypothesized that a similar mechanism may explain the poor T-cell homeostasis that is observed in this HIS mouse model.

Previous studies in mice demonstrated increased CD8<sup>+</sup> T-cell numbers following treatment with transpresented mouse IL-15 (14, 21). Mouse CD4<sup>+</sup> T cells do not respond to IL-15 in vivo, but human CD4<sup>+</sup> T cells can express IL-2R $\beta$  and  $\gamma_c$ , raising the possibility that these cells could respond to IL-15 in vivo. In this manuscript, we validate this hypothesis by showing that transpresented human IL-15/IL-15R $\alpha$  complexes (RLI) can induce both naïve and memory human CD4<sup>+</sup> and CD8<sup>+</sup> T cells to proliferate in vivo, thereby identifying a fundamental difference between mouse and human T cells with respect to cytokine-regulated T-cell homeostasis. The RLI treatment in HIS mice resulted in a clear increase in global T-cell numbers by promoting both the survival and proliferation of peripheral human T cells.

The observation that IL-15 receptor ligation promotes naïve CD4<sup>+</sup> T cell proliferation in vivo differs from in vitro studies where IL-15 only promoted the generation of effector memory phenotype CD4<sup>+</sup> T cells from central memory phenotype CD4<sup>+</sup> T cells (20). Whereas previous studies of IL-15 actions on human T cells were performed in vitro, our results represent a first pre-clinical study of transpresented human IL-15 on human T cells in vivo. Transpresented human IL-15 may directly stimulate human naïve and memory T cells in vivo in HIS mice or, alternatively, may activate myeloid cells that then indirectly influence human T-cell homeostasis. Some evidence for the latter was provided by the observation of increased human MHC class I and II expression in RLI-treated HIS mice. Thus, our study highlights the value of the HIS model to uncover differences between mouse and human T-cell biology and to study human immunology in vivo.



**Fig. 3.** Increased frequency of T-dependent B-cell responses following RLI treatment in vivo. (A) Total serum IgM and IgG concentrations were determined by ELISA in nonimmunized HIS mice and HIS mice immunized with vaccines against hepatitis B virus (HBV) and tetanus toxoid (TT). Values represent mean  $\pm$  SEM of 16 mice per group. (B) TT-specific serum IgG or (C) HBV surface antigen (HbsAg)-specific serum Ig from immunized HIS mice were determined by ELISA. Values represent mean  $\pm$  SEM of 29 mice per treatment group. (D) Frequency of “vaccine responders” (mice with serum values >0.5 IU/mL anti-TT IgG or >7 IU/L anti-HbsAg-Ig) as determined in B and C.  $\chi^2$  value  $\chi^2 = 3.85$ ,  $P < 0.05$ .





1. Gimeno R, et al. (2004) Monitoring the effect of gene silencing by RNA interference in human CD34+ cells injected into newborn RAG2-/- gammac-/- mice: Functional inactivation of p53 in developing T cells. *Blood* 104:3886–3893.
2. Traggiai E, et al. (2004) Development of a human adaptive immune system in cord blood cell-transplanted mice. *Science* 304:104–107.
3. Huntington ND, et al. (2009) IL-15 trans-presentation promotes human NK cell development and differentiation in vivo. *J Exp Med* 206:25–34.
4. Legrand N, et al. (2006) Transient accumulation of human mature thymocytes and regulatory T cells with CD28 superagonist in “human immune system” Rag2(-/-) gammac(-/-) mice. *Blood* 108:238–245.
5. van Lent AU, et al. (2009) IL-7 enhances thymic human T cell development in “human immune system” Rag2-/-IL-2Rgammac-/- mice without affecting peripheral T cell homeostasis. *J Immunol* 183:7645–7655.
6. Surh CD, Sprent J (2008) Homeostasis of naive and memory T cells. *Immunity* 29: 848–862.
7. Di Santo JP, Kühn R, Müller W (1995) Common cytokine receptor gamma chain (gamma c)-dependent cytokines: Understanding in vivo functions by gene targeting. *Immunol Rev* 148:19–34.
8. Buckley RH (2004) Molecular defects in human severe combined immunodeficiency and approaches to immune reconstitution. *Annu Rev Immunol* 22:625–655.
9. Gilmour KC, et al. (2001) Defective expression of the interleukin-2/interleukin-15 receptor beta subunit leads to a natural killer cell-deficient form of severe combined immunodeficiency. *Blood* 98:877–879.
10. Dubois S, Mariner J, Waldmann TA, Tagaya Y (2002) IL-15Ralpha recycles and presents IL-15 in trans to neighboring cells. *Immunity* 17:537–547.
11. Ma A, Koka R, Burkett P (2006) Diverse functions of IL-2, IL-15, and IL-7 in lymphoid homeostasis. *Annu Rev Immunol* 24:657–679.
12. Waldmann TA, Dubois S, Tagaya Y (2001) Contrasting roles of IL-2 and IL-15 in the life and death of lymphocytes: Implications for immunotherapy. *Immunity* 14:105–110.
13. Stoklasek TA, Schluns KS, Lefrançois L (2006) Combined IL-15/IL-15Ralpha immunotherapy maximizes IL-15 activity in vivo. *J Immunol* 177:6072–6080.
14. Rubinstein MP, et al. (2006) Converting IL-15 to a superagonist by binding to soluble IL-15Ralpha. *Proc Natl Acad Sci USA* 103:9166–9171.
15. Mortier E, et al. (2006) Soluble interleukin-15 receptor alpha (IL-15R alpha)-sushi as a selective and potent agonist of IL-15 action through IL-15R beta/gamma. Hyperagonist IL-15 x IL-15R alpha fusion proteins. *J Biol Chem* 281:1612–1619.
16. Macchiarelli F, Manz MG, Palucka AK, Shultz LD (2005) Humanized mice: Are we there yet? *J Exp Med* 202:1307–1311.
17. Legrand N, et al. (2009) Humanized mice for modeling human infectious disease: Challenges, progress, and outlook. *Cell Host Microbe* 6:5–9.
18. Legrand N, Weijer K, Spits H (2006) Experimental models to study development and function of the human immune system in vivo. *J Immunol* 176:2053–2058.
19. Huntington ND, Di Santo JP (2008) Humanized immune system (HIS) mice as a tool to study human NK cell development. *Curr Top Microbiol Immunol* 324:109–124.
20. Geginat J, Sallusto F, Lanzavecchia A (2001) Cytokine-driven proliferation and differentiation of human naive, central memory, and effector memory CD4(+) T cells. *J Exp Med* 194:1711–1719.
21. Dubois S, Patel HJ, Zhang M, Waldmann TA, Müller JR (2008) Preassociation of IL-15 with IL-15R alpha-IgG1-Fc enhances its activity on proliferation of NK and CD8+/CD44high T cells and its antitumor action. *J Immunol* 180:2099–2106.
22. Berger C, et al. (2009) Safety and immunologic effects of IL-15 administration in nonhuman primates. *Blood* 114:2417–2426.
23. Bessard A, Solé V, Bouchaud G, Quémener A, Jacques Y (2009) High antitumor activity of RLI, an interleukin-15 (IL-15)-IL-15 receptor alpha fusion protein, in metastatic melanoma and colorectal cancer. *Mol Cancer Ther* 8:2736–2745.
24. Chapoval AI, Fuller JA, Kremlev SG, Kamdar SJ, Evans R (1998) Combination chemotherapy and IL-15 administration induce permanent tumor regression in a mouse lung tumor model: NK and T cell-mediated effects antagonized by B cells. *J Immunol* 161:6977–6984.
25. Epardaud YM, et al. (2008) Interleukin-15/interleukin-15R alpha complexes promote destruction of established tumors by reviving tumor-resident CD8+ T cells. *Cancer Res* 68:2972–2983.
26. Lu J, et al. (2002) Interleukin 15 promotes antigen-independent in vitro expansion and long-term survival of antitumor cytotoxic T lymphocytes. *Clin Cancer Res* 8: 3877–3884.
27. Alves NL, Hooibrink B, Arosa FA, van Lier RA (2003) IL-15 induces antigen-independent expansion and differentiation of human naive CD8+ T cells in vitro. *Blood* 102:2541–2546.
28. Oh S, Berzofsky JA, Burke DS, Waldmann TA, Perera LP (2003) Coadministration of HIV vaccine vectors with vaccinia viruses expressing IL-15 but not IL-2 induces long-lasting cellular immunity. *Proc Natl Acad Sci USA* 100:3392–3397.
29. Becker TC, et al. (2002) Interleukin 15 is required for proliferative renewal of virus-specific memory CD8 T cells. *J Exp Med* 195:1541–1548.
30. Teague RM, et al. (2006) Interleukin-15 rescues tolerant CD8+ T cells for use in adoptive immunotherapy of established tumors. *Nat Med* 12:335–341.
31. Mueller YM, et al. (2003) IL-15 enhances survival and function of HIV-specific CD8+ T cells. *Blood* 101:1024–1029.
32. Mueller YM, et al. (2005) Interleukin-15 increases effector memory CD8+ t cells and NK Cells in simian immunodeficiency virus-infected macaques. *J Virol* 79:4877–4885.
33. Shinkai Y, et al. (1992) RAG-2-deficient mice lack mature lymphocytes owing to inability to initiate V(D)J rearrangement. *Cell* 68:855–867.
34. DiSanto JP, Müller W, Guy-Grand D, Fischer A, Rajewsky K (1995) Lymphoid development in mice with a targeted deletion of the interleukin 2 receptor gamma chain. *Proc Natl Acad Sci USA* 92:377–381.
35. Huntington ND, Xu Y, Nutt SL, Tarlinton DM (2005) A requirement for CD45 distinguishes Ly49D-mediated cytokine and chemokine production from killing in primary natural killer cells. *J Exp Med* 201:1421–1433.
36. Lim A, et al. (2008) Many human peripheral VH5-expressing IgM+ B cells display a unique heavy-chain rearrangement. *Int Immunol* 20:105–116.