



Community Fact Sheet

Lowell, Massachusetts

Data Updated: 10/27/11

Cigarette Smoking

An estimated 18,592 smokers live in Lowell (24.2% of adults, age 18+).

The adult smoking rate is 50 percent higher in Lowell than statewide (24.2% in Lowell compared to 16.1% statewide).

Cigarette smoking among middle age adults (age 45 to 64) is 29 percent higher in Lowell (21.5%) than statewide (16.7%).

Cigarette smoking among seniors (age 65 and older) is 6 percent higher in Lowell (8.7%) than statewide (8.2%).

The rate of smoking during pregnancy in Lowell is 54 percent higher than for the overall state of Massachusetts (11.4% in Lowell compared to 7.4% statewide).

Quitting

60 percent of current smokers in Lowell made a quit attempt within the past year.

MassHealth Tobacco Cessation Benefit

Since coverage of the tobacco cessation began in July 2006, 1,763 MassHealth smokers from Lowell have used the benefit - an estimated 27.3% of MassHealth smokers living in Lowell. Statewide, more than 75,000 MassHealth smokers (41%) have used the tobacco cessation benefit since July 2006.

QuitWorks

Health care providers referred 384 smokers living in Lowell to the QuitWorks program to help them quit smoking. In addition, 1,387 people from Lowell called the Massachusetts Smokers' Helpline to quit smoking (fiscal years 2004 to 2009).

Illegal Tobacco Sales to Minors

The rate of illegal sales to minors (those under age 18) is 25 percent higher in Lowell (10.1%) compared to the state of Massachusetts (8.1%) based on data from FY 2010.



Massachusetts Department of Public Health
Tobacco Cessation and Prevention Program
(617) 624-5900 www.mass.gov/dph/mtcp



Community Fact Sheet

Lowell, Massachusetts

Data Updated: 10/27/11

Health Effects of Smoking

Mortality from lung cancer is 36 percent higher among males in Lowell compared to the state of Massachusetts.

The rate of hospitalizations for lung cancer is not significantly different among females in Lowell compared to the state of Massachusetts.

The rate of hospitalizations for lung cancer is 28 percent higher among males in Lowell compared to the state of Massachusetts.

Data Sources

Smoking figures are based on data from the 2008 Behavioral Risk Factor Surveillance System (BRFSS), Massachusetts Department of Public Health.

Figures on smoking during pregnancy are based on 2003 to 2007 Births (Vital Records), Massachusetts Department of Public Health.

Figures on quitting are based on 2008 Behavioral Risk Factor Surveillance System (BRFSS), Massachusetts Department of Public Health.

MassHealth utilization of the tobacco cessation benefit was obtained from the Executive Office of Health and Human Services, MassHealth agency covering fiscal years 2007 to 2009 (through April 30, 2009).

The number of referrals to the QuitWorks program and calls to the Massachusetts Smokers Helpline is based on data collected by the Massachusetts Tobacco Cessation and Prevention Program from fiscal years 2004 to 2009.

The rate of illegal sales to minors is based on compliance checks performed in Lowell during FY 2010 (n=368).

Figures on lung cancer mortality are based on 2003 to 2007 Deaths (Vital Records), Massachusetts Department of Public Health.

Rates of lung cancer hospitalizations are age-adjusted and based on data from the 2002 to 2006 Uniform Hospital Discharge Data System (UHDDS) maintained by the Massachusetts Division of Health Care Finance and Policy.



Massachusetts Department of Public Health
Tobacco Cessation and Prevention Program
(617) 624-5900 www.mass.gov/dph/mtcp