

ORIGINAL ARTICLE

Tobacco Smoking Trajectory and Associated Ethnic Differences Among Adolescent Smokers Seeking Cessation Treatment

MIQUN L. ROBINSON, M.D., Ph.D., IVAN BERLIN, M.D., Ph.D., AND
ERIC T. MOOLCHAN, M.D.

Purpose: To examine smoking trajectories in a clinical sample of adolescent smokers seeking cessation treatment, including: (a) smoking onset (initial, daily) and time intervals from initial to daily smoking and from daily smoking to treatment request, (b) associations between current level of tobacco dependence and smoking history, and (c) differences in smoking trajectory between African-American and non-African-American youth.

Methods: Four hundred and thirty-two adolescent smokers (aged 13–17 years, 61.8% female, 32% African-American) responding to various media advertisement completed a telephone interview as part of pre-eligibility screening for a smoking cessation trial. Smoking trajectory data included age at onset of initial and daily smoking, intervals between those time points, and cigarettes smoked per day (CPD). Tobacco dependence was assessed using the Fagerström Test for Nicotine Dependence (FTND). Data were analyzed using regression models and multiple analyses of covariance.

Results: Initial smoking occurred at a mean age of less than 12 years and daily smoking at age 13 years. Earlier onset of daily smoking was associated with higher FTND scores and longer duration from daily smoking to treatment request. For the entire sample, the time interval from initial to daily smoking was 1.14 years. When the

sample was divided into early (before age 14 years) and later (at or after age 14 years) initiators, early initiators showed a slower progression from initial to daily smoking compared with late initiators (16 months vs. 6 months). Compared with non-African-American teen smokers, African-American youth reported a 1-year delay in onset of both initial and daily smoking.

Conclusions: Early age of daily smoking and short time interval from initial to daily smoking highlight a brief window of opportunity to prevent the development of tobacco addiction and its consequences. Ethnic differences in smoking trajectory uncovered in this report call for ethnically tailored interventions to reduce youth smoking. © Society for Adolescent Medicine, 2004

KEY WORDS:

Adolescents
Cessation treatment
Ethnic differences
Smoking trajectory
Tobacco dependence

Many adolescents start smoking cigarettes despite public health efforts to dissuade youth from experimenting with tobacco [1], the leading preventable cause of morbidity and mortality in the United States. Although one-third to one-half of adolescents who try cigarettes later became regular or dependent smokers, close to two-thirds of them report having tried to quit [1–3]. Ample evidence supports the effectiveness of adult tobacco dependence treatment programs [4], but few cessation programs are available to accommodate the needs of adolescent smokers.

From the Teen Tobacco Addiction Treatment Research Clinic, National Institute on Drug Abuse Intramural Research Program, National Institutes of Health, Baltimore, Maryland (M.L.R., E.T.M.); and Department of Pharmacology, Pitié-Salpêtrière Medical School, Paris, France (I.B.).

Address correspondence to: Eric T. Moolchan, M.D., Intramural Research Program, Clinical Pharmacology and Therapeutics Research Branch, National Institute on Drug Abuse, 5500 Nathan Shock Drive, Baltimore, MD 21224. E-mail: emoolcha@intra.nida.nih.gov

Manuscript accepted October 29, 2003.

ers before smoking behavior becomes a refractory disorder [5–7]. This deficit is compounded by limited knowledge of the temporal duration of various stages of smoking behavior (smoking trajectory). Because more than one-half of the adolescent smokers who quit remain abstinent as young adults [8], early cessation treatment has significant importance. Data on smoking trajectory would facilitate the optimal timing for implementing developmentally appropriate prevention and early cessation interventions. This report examines the various time points and intervals along the smoking trajectory in a sample of adolescent smokers who request assistance for cessation, a group whose higher health risks need to be recognized because of their dependence-related consumption. Although various definitions of smoking “trajectory” may exist [9], our focus is based on a linear progression along the smoking trajectory, which refers to three time points and two intervals. Time points include the age at onset of first cigarette, age at onset of daily smoking, and age at onset of treatment request (professional assistance to quit smoking). Time intervals include initial to daily smoking and daily smoking to treatment request.

Several lines of evidence indicate that early age of smoking onset (first cigarette) predicts long-term smoking. For example, Chassin et al reported that the “early-stable” group of smokers (those with a younger age of smoking onset) attained a high rate of smoking 11–20 cigarettes a day by age 15 years and remained stable smokers through the 10-year course of the longitudinal study [10]. Smokers who started before age 14 years [11] or 16 years [12] were likely to smoke more cigarettes in adulthood. Additionally, age of smoking onset was associated with a higher level of tobacco dependence and a lower number of quit attempts [12–14]. Therefore, it is likely that there is a relationship between early smoking onset and substantial smoking history in cessation-seeking adolescent smokers.

Whereas the initiation of smoking has been studied for its significance, less is known about the temporal progression from smoking onset to daily smoking. Cross-sectional data suggest that about 30% of experimenters progress to “established” (i.e., daily) smoking within 5 years, and an identifiable subgroup show relatively higher risk of such progression [15]. Breslau et al reported a 3-year interval from first cigarette to daily smoking in a young adult cohort and this duration was independent of the number of cigarettes consumed [16]. Data characterizing the duration from daily smoking to cessation treatment are also scant [12,17]. Determining the

time frame from initial to daily smoking would seem crucial for intervening to prevent the development of dependent smoking [18,19].

In addition to studies in the general smoking population, age of smoking onset seems to vary among ethnic groups. Among non-treatment-seeking populations (both adult and adolescent smokers), African-American smokers were found to initiate smoking at a later age and to smoke fewer cigarettes per day compared with non-African-American smokers [20–23]. Later onset of smoking among African-American youth was also reported from a cessation treatment-seeking population [24]. However, once African-American teens began smoking, almost half of them developed addictive smoking, particularly in those who began smoking before age 17 years [25]. Thus, elucidating differences in the time course of smoking trajectory in a cessation-seeking sample may advance the knowledge leading to ethnically tailored treatment interventions to address tobacco-attributable health disparities [26].

Within a sample of adolescent smokers seeking assistance for cessation, this study had three objectives: (a) to examine reported age of onset of initial and daily smoking and intervals between key time points in the smoking trajectory, (b) to explore the association between current level of tobacco dependence and time course of smoking, and (c) to examine ethnic differences (African-American versus non-African-American) in smoking trajectory in this cessation-seeking sample.

Methods

This report was based on data obtained from telephone interviews that screened participants for eligibility for a combined nicotine replacement and counseling cessation study for adolescent smokers who were motivated to quit smoking. The screening protocol was approved by the National Institute on Drug Abuse (NIDA) Institutional Review Board (IRB) with a waiver of informed consent.

Subjects

From September 1, 1999 to September 30, 2001, a convenience sample of 432 adolescent participants between the ages of 13 and 17 years were recruited from the Baltimore area via several means of advertisement, including newspaper, radio, television, and word of mouth. One hundred thirty-eight participants were African-American, 282 were Caucasian, 4

Native American, 1 Asian, and 7 Hispanic. The latter four groups were collapsed into a non-African-American group. There were 88 (63.8%) females in the African-American group and 179 (60.9%) in the non-African-American group.

Procedures

Participants telephoned to inquire about the eligibility to participate in the treatment study and completed an internally developed structured telephone interview, lasting approximately 15 to 20 minutes. At the beginning of each interview, interviewers obtained permission from participants to record their information. Data collected included demographic information (age, gender, ethnicity), medical, psychiatric, and medication histories, motivation to quit (integer scale of 10, 10 being the highest), and smoking-related histories.

Measures

Smoking-related histories included age of onset of first cigarette, age of daily smoking, number of cigarettes consumed (weekdays and weekend), previous cessation attempts (number of times and duration), and current level of nicotine dependence. For data on the age of onset of smoking initiation, callers were asked, "How old were you when you smoked your first whole cigarette?" For data on age of onset of daily smoking, callers were asked, "At what age did you begin smoking daily?" The number of cigarettes smoked per day (CPD) was calculated by summing of weekly consumption and dividing by number of days of a week. Intervals of time points (e.g., from initial to daily smoking, from daily smoking to treatment request) were calculated by subtracting length of time, rounded off to the nearest year. Because of use of age in years rather than dates, for those who reported time points during the same year of age, intervals were calculated as zero. Participants who began to smoke before age 14 years were defined as "early initiators," and those who started at or later than age 14 years were categorized as "later initiators."

Tobacco dependence was assessed using the six-item Fagerström Test for Nicotine Dependence (FTND) [27]. FTND scores range from 0 to 10; suggested interpretations of FTND scores is as follows: 1–2 = very low dependence; 3–4 = low dependence; 5 = medium dependence; 6–7 = high dependence; 8–10 = very high dependence [28]. The FTND

Table 1. Smoking Trajectory of Adolescents Seeking Cessation Treatment

Variable	Mean \pm SD
Age at first cigarette (years)	11.9 \pm 2.0
Age at daily smoking (years)	13.0 \pm 1.7
Age at treatment request (years)	15.6 \pm 1.5
First cigarette to daily smoking (years)	1.1 \pm 1.2
Daily smoking to treatment request (years)	2.7 \pm 1.6
Fagerström Test for Nicotine Dependence (FTND)	5.9 \pm 2.2
Cigarette smoked per day (CPD)	14.5 \pm 7.9

have been found to have good test-retest reliability (Cronbach alpha 0.64) and higher internal consistency than FTQ in several studies [29–31].

Data Analysis

Before tabulation, we removed the personal identifiers (unlinked) of participants found ineligible for the treatment protocol. Because subsequent analyses were performed on such unlinked data, the current screening study was separately approved with an exemption by Institutional Review Board, the National Institutes of Health, Office of Human Subject Research (OHSR).

Frequencies were compared using Chi-square tests. The age at enrollment was significantly different in African-American and non-African-American adolescents, thus this variable was included in all analyses as a covariate. Standard multiple regression analysis was used to test the association of FTND with trajectory variables. Continuous variables were compared by multivariate analysis of variance (MANCOVA) followed by univariate analysis of variance to test specific variables for significant differences. Backward conditional logistic regression was used to determine which trajectory variables were independently associated with ethnicity. Relationships among variables were estimated by stepwise multiple linear regression. Data are presented as means [\pm Standard Deviation (SD) or frequencies (%)] unless otherwise indicated. Differences were considered significant at the $p < .05$ levels. SPSS version 10.0 (Chicago, IL) was used for all data analyses.

Results

Sample Smoking Characteristics

The time course of progression through various phases of the smoking cycle in our sample of cessation seekers is illustrated in Table 1. No gender

Table 2. Differences in Time Intervals of Smoking Stages in Early and Late Initiators

Variable	Early Initiator (<14 years) (Mean \pm SD)	Late Initiator (\geq 14 years) (Mean \pm SD)	F Value	p Value
First cigarette to daily smoking (years)	1.3 \pm 1.3	0.45 \pm 0.7	35.4	<.0001
Daily smoking to treatment request (years)	2.9 \pm 1.7	1.6 \pm 0.8	46.8	<.0001
First cigarette to treatment request (years)	4.2 \pm 2.0	2.0 \pm 0.9	90.0	<.0001

differences were found among any of the smoking trajectory variables.

Nicotine Dependence and Smoking Trajectory

Pearson correlation analyses showed that higher CPD was associated with a longer interval from first cigarette to seeking treatment ($r = 0.322$, $p < .0001$) and time from daily use to treatment request ($r = 0.322$, $p < .0001$). CPD did not, however, correlate with interval from first cigarette to daily smoking ($r = 0.05$, $p = .32$).

In the overall sample, the age of first cigarette use bore no relationship to current FTND scores. However, age of onset of *daily* smoking was associated with FTND score such that subjects who had an earlier onset of daily smoking had higher FTND scores (beta = 0.494, SE = 0.06, $t = -7.88$, $p < .0001$). FTND score was also positively associated with age at seeking cessation treatment request such that older participants at enrollment had higher FTND scores (beta = 0.467, SE = 0.07, $t = 6.3$, $p < .0001$). FTND score was not associated with interval from age of first cigarette to daily smoking, but was associated with interval from daily smoking to age at treatment request (beta = 0.48, SE = 0.06, $t = 8.35$, $p < .0001$) such that participants who had higher FTND scores had a longer duration from daily smoking to treatment request.

After controlling for age at treatment request, early initiators (before age 14 years) currently consumed significantly more cigarettes (Mean \pm SE 15.4

± 0.5 vs. 10.0 ± 0.9 , $F_{1,424} = 23.7$, $p = .005$) and had higher FTND scores (Mean \pm SE 6.2 ± 0.1 vs. 4.7 ± 0.2 , $F_{1,424} = 30.4$, $p < .0001$) compared with late initiators. However, differences between these groups became nonsignificant for both CPD and FTND after adjusting for the years of smoking, suggesting that dependence level and CPD are related to the duration of smoking.

As shown in Table 2, early initiators had a significantly longer interval from first cigarette smoked to daily smoking and from daily smoking to treatment request.

Ethnic Differences in Smoking Trajectory

African-American adolescents ($n = 126$, 30%) (12 African-American participants were excluded from the original sample owing to lack of age of onset of smoking) exhibited a later onset of first cigarette smoked and daily smoking than non-African-Americans (Table 3). When data were adjusted for age and gender, African-American teens had shorter intervals from any one stage to another. However, backward logistic regression analysis showed that the only trajectory variable associated with the ethnic difference was the age of the first cigarette (odds ratio: 0.736, 95% CI: 0.65 to 0.83, African-American group being the reference group, $p < .0001$, i.e., African-American teens had a later onset of smoking initiation compared with non-African-American participants).

Table 3. Ethnic Differences in Smoking Trajectory and Tobacco Dependence^a

Variable	African-American (n = 126)	Non-African-American (n = 279)	p Value
Age at first cigarette (years)	12.6 \pm 0.17	11.5 \pm 0.12	.0001
Age at daily smoking onset (years)	13.6 \pm 0.15	12.8 \pm 0.10	.0001
First cigarette to daily smoking (years)	0.95 \pm 0.11	1.2 \pm 0.7	.031
Daily smoking to treatment request (years)	2.4 \pm 1.6	2.8 \pm 1.6	.018
First cigarette to treatment (years)	3.3 \pm 0.18	4.0 \pm 0.12	.001
Cigarettes smoked per day	10.7 \pm 0.7	15.9 \pm 0.5	.0001
Fagerström Test for Nicotine Dependence (FTND)	5.2 \pm 0.2	6.2 \pm 0.1	.0001

^a Data are age-adjusted Means \pm SE.

African-American adolescents smoked significantly fewer cigarettes than non-African-American teens ($F_{1,424} = 31.9, p < .0001$) and had lower FTND scores ($F_{1,424} = 20.4, p < .0001$) than their counterparts. When the comparison of FTND was adjusted for CPD, the difference in FTND was no longer significant.

Although African-American adolescents reported lower CPD than non-African-American youth, African-Americans who began to smoke before age 14 years had greater mean daily cigarette consumption than those who initiated smoking after age 14 (Mean \pm SD 12.4 ± 7.7 vs. 9.6 ± 9.0 , $F_{1,136} = 3.70, p = 0.57$).

Discussion

In our sample of adolescent smokers seeking assistance for cessation, smoking onset occurred on average just before age 12 years and daily smoking around age 13 years. Earlier onset of daily smoking was associated with a higher level of tobacco dependence and longer duration from daily smoking to age of treatment request. Early initiators had a longer interval from first cigarette to daily smoking than later initiators. Early initiators endorsed a similar level of tobacco dependence as later initiators when adjusted for years of smoking meaning that years of smoking is an important factor in predicting dependence. Compared with non-African-American youth, African-American smokers reported a later onset of first cigarette and daily smoking.

Age of Initial and Daily Smoking

Our clinical population reported an earlier age of smoking initiation than other previous reports [13,16,32–34]. Although a caution for the generalizability of our data to other young smokers is warranted, the young age of smoking initiation and heavy use of cigarettes support the notion that adolescent smokers in our sample represent a “self-selected” subgroup who start smoking early in life, are highly dependent, and are unlikely to quit [10,14,35].

In this report we used daily smoking as a proxy measure for dependent smoking instead of using definitive values from various scales [36,37] because (a) the simplicity and familiarity of participants’ recall of daily smoking enhance accuracy; (b) there is no universally approved scale of dependence for adolescent smokers; (c) daily smoking is representative of compulsive use; and (d) daily smoking signals

a lifestyle or identity and a probable social affiliation to a smoking group [38]. The age of onset of daily smoking reported here (13 years old) is similar to one report but substantially younger than findings of other studies of non-treatment-seeking populations [39–41]. Our finding that early daily smoking predicts higher level of nicotine dependence supports the association between daily smoking and tobacco dependence [42]. Whereas some reports suggest that tobacco dependence can develop before daily smoking after a relatively short history of tobacco use, others suggest a 1-year delay from the onset of daily smoking to the development of tobacco dependence [43,44].

Duration/Progression of Smoking

The mean time interval from initial to daily smoking in our sample (1.14 years) is comparable to findings from one epidemiological study [33] but is substantially shorter than other previous findings of 2- [34] and 3-year intervals reported by Breslau [41]. This seems to point to a relatively short opportunity for offsetting the development of compulsive tobacco use. When our sample was divided into early and later initiators, early initiators progressed to daily smoking in 16 months, whereas the later initiators did so in less than 6 months. One possible explanation for a longer progression time to daily smoking in early initiators are the limited opportunities to smoke owing to parental and social disapproval of smoking at this younger age [16,45]. Additionally, reduced access to cigarettes for younger smokers may further deter the progression to daily smoking [46]. Although in some individuals, less time is needed to escalate from experimental to established smoking [43,47,48], the data showing that late initiators progress to daily smoking quicker than early initiators may be explained by experiencing contextual social influences of teenage lifestyle [38], including diminished parental supervision as a function of age, and increased identity as a social smoker. Recent data suggest that having more close friends who smoke may accelerate the progression from experimental to regular smoking [49]. Moreover, older teens may have easier access to cigarettes through smoking parents or friends who work in stores [50]. We suggest that early interventions occur before attitudinal changes that promote the onset of smoking and other risk behaviors; such as before age 10 years [51,52].

Correlates of Smoking Trajectory and Tobacco Dependence

Adjusted for age at study enrollment, early initiators appear to have a higher level of tobacco dependence measured by FTND scores and a higher rate of daily cigarette consumption. Such difference diminishes after the years of smoking were taken into account, indicating that considerations of multiple factors are needed when assessing tobacco dependence, probably length of exposition being a main factor to become dependent. As an overall sample, the longer time interval from daily smoking to time of seeking cessation treatment is associated with higher CPD and level of tobacco dependence. This finding echoes previous analyses that in both adolescent and adult smokers, heavy smokers tend to make less effort to quit smoking [17,53,54]. Fewer treatment trials among young smokers have resulted in limited information for cessation assistance to adolescent smokers [55]. This situation might have contributed to a delay in seeking cessation treatment after the onset of daily smoking.

Ethnic Differences

Consistent with several previous reports, African-American youth in our sample reported a later onset of first cigarette smoked compared with non-African-American teens (12.64 years vs. 11.55 years, respectively) [1,20,24,32,56]. In addition, African-American teens reported later onset of daily smoking (13.59 years) compared with non-African-American youth (12.78 years). Parental restrictions and less societal acceptance for cigarette smoking in the African-American population might partially explain these findings [22,57,58]. Detailed analyses showed no difference in trajectory intervals between African-American and non-African-American participants, indicating similar progression from initial to daily smoking and from daily smoking to time of treatment request in both groups. Moreover, African-American youth showed similar tobacco dependence as non-African-American participants, as reported elsewhere [26].

Limitations

Our findings should not be generalized to all adolescent smokers we used a self-selected sample of motivated cessation seekers. Because interview data were based upon recollection, our data are subject to recall bias. However, available evidence has shown

that self-reported information on smoking initiation and substance use by both adults and children are sufficiently reliable for reporting retrospective data [59,60]. Nonetheless, prospective studies of adolescent smokers would offer stronger evidence of the time course of smoking trajectories. Socioeconomic status and educational level, which are shown to be associated with smoking initiation [32,61], as well as influence of peer smoking on progression [49] were not obtained during the brief screening interview.

Conclusions

The current study identified the characteristics of smoking trajectory in a clinical sample of tobacco cessation-seeking adolescents. Our sample reported an early age of smoking onset and rapid progression from initial to daily smoking. These findings suggest that there is a relatively brief window of opportunity for intervention, thus underscoring the need for providing appropriate and prompt interventions to prevent the development of tobacco addiction. Moreover, ethnic differences in smoking trajectory uncovered here, namely later onset of initial and daily smoking, may help to design ethnically tailored smoking cessation interventions.

References

1. Trends in cigarette smoking among high school students—United States, 1991–2001. *MMWR* 2002;51:409–12.
2. Pallonen UE, Prochaska JO, Velicer WF, et al. Stages of acquisition and cessation for adolescent smoking: An empirical integration. *Addict Behav* 1998;23:303–24.
3. Prokhorov AV, Hudmon KS, de Moor CA, et al. Nicotine dependence, withdrawal symptoms, and adolescents' readiness to quit smoking. *Nicotine Tob Res* 2001;3:151–5.
4. Fiore MC, Hatsukami DK, Baker TB. Effective tobacco dependence treatment. *JAMA* 2002;288:1768–71.
5. Smith TA, House RF, Jr, Croghan IT, et al. Nicotine patch therapy in adolescent smokers. *Pediatrics* 1996;98:659–67.
6. Hurt RD, Croghan GA, Beede SD, et al. Nicotine patch therapy in 101 adolescent smokers: Efficacy, withdrawal symptom relief, and carbon monoxide and plasma cotinine levels. *Arch Pediatr Adolesc Med* 2000;154:31–7.
7. Maharaj K, Ternullo S. Using nicotine replacement therapy in treating nicotine addiction in adolescents. *J Sch Nurs* 2001;17:278–82.
8. Chassin L, Presson CC, Sherman SJ, et al. The natural history of cigarette smoking: Predicting young-adult smoking outcomes from adolescent smoking patterns. *Health Psychol* 1990;9:701–16.
9. White HR, Pandina RJ, Chen PH. Developmental trajectories of cigarette use from early adolescence into young adulthood. *Drug Alcohol Depend* 2002;65:167–78.
10. Chassin L, Presson CC, Pitts SC, et al. The natural history of cigarette smoking from adolescence to adulthood in a mid-

- western community sample: Multiple trajectories and their psychosocial correlates. *Health Psychol* 2000;19:223-31.
11. Taioli E, Wynder EL. Effect of the age at which smoking begins on frequency of smoking in adulthood. *N Engl J Med* 1991;325:968-9.
 12. Khuder SA, Dayal HH, Mutgi AB. Age at smoking onset and its effect on smoking cessation. *Addict Behav* 1999;24:673-7.
 13. Lando HA, Thai DT, Murray DM, et al. Age of initiation, smoking patterns, and risk in a population of working adults. *Prev Med* 1999;29:590-8.
 14. Breslau N, Peterson EL. Smoking cessation in young adults: Age at initiation of cigarette smoking and other suspected influences. *Am J Public Health* 1996;86:214-20.
 15. Choi WS, Patten CA, Gillin JC, et al. Cigarette smoking predicts development of depressive symptoms among U.S. adolescents. *Ann Behav Med* 1997;19:42-50.
 16. Breslau N, Fenn N, Peterson EL. Early smoking initiation and nicotine dependence in a cohort of young adults. *Drug Alcohol Depend* 1993;33:129-37.
 17. Breslau N, Johnson EO. Predicting smoking cessation and major depression in nicotine-dependent smokers. *Am J Public Health* 2000;90:1122-7.
 18. Henningfield JE, Moolchan ET, Zeller M. Regulatory strategies to reduce tobacco addiction in youth. *Tob Control* 2003;12(Suppl 1):114-24.
 19. Moolchan ET, Aung AT, Henningfield JE. Treatment of adolescent tobacco smokers: Issues and opportunities for exposure reduction approaches. *Drug Alcohol Depend* 2003;70:223-32.
 20. Faulkner DL, Escobedo LG, Zhu BP, et al. Race and the incidence of cigarette smoking among adolescents in the United States. *J Natl Cancer Inst* 1996;88:1158-60.
 21. Griesler PC, Kandel DB, Davies M. Ethnic differences in predictors of initiation and persistence of adolescent cigarette smoking in the National Longitudinal Survey of Youth. *Nicotine Tob Res* 2002;4:79-93.
 22. Robinson LA, Klesges RC. Ethnic and gender differences in risk factors for smoking onset. *Health Psychol* 1997;16:499-505.
 23. Headen SW, Bauman KE, Deane GD, et al. Are the correlates of cigarette smoking initiation different for black and white adolescents? *Am J Public Health* 1991;81:854-8.
 24. Moolchan ET, Berlin I, Robinson ML, et al. African-American teen smokers: Issues to consider for cessation treatment. *J Natl Med Assoc* 2000;92:558-62.
 25. Juon HS, Ensminger ME, Sydnor KD. A longitudinal study of developmental trajectories to young adult cigarette smoking. *Drug Alcohol Depend* 2002;66:303-14.
 26. Moolchan ET, Berlin I, Robinson ML, et al. Characteristics of African American teenage smokers who request cessation treatment: Implications for addressing health disparities. *Arch Pediatr Adolesc Med* 2003;157:533-8.
 27. Heatherton TF, Kozlowski LT, Frecker RC, et al. The Fagerstrom Test for Nicotine Dependence: A revision of the Fagerstrom Tolerance Questionnaire. *Br J Addict* 1991;86:1119-27.
 28. Fagerstrom KO, Heatherton TF, Kozlowski LT. Nicotine addiction and its assessment. *Ear Nose Throat J* 1990;69:763-5.
 29. Pomerleau CS, Carton SM, Lutzke ML, et al. Reliability of the Fagerstrom Tolerance Questionnaire and the Fagerstrom Test for Nicotine Dependence. *Addict Behav* 1994;19:33-9.
 30. Payne TJ, Smith PO, McCracken LM, et al. Assessing nicotine dependence: A comparison of the Fagerstrom Tolerance Questionnaire (FTQ) with the Fagerstrom Test for Nicotine Dependence (FTND) in a clinical sample. *Addict Behav* 1994;19:307-17.
 31. Kozlowski LT, Porter CQ, Orleans CT, et al. Predicting smoking cessation with self-reported measures of nicotine dependence: FTQ, FTND, and HSI. *Drug Alcohol Depend* 1994;34:211-6.
 32. Harrell JS, Bangdiwala SI, Deng S, et al. Smoking initiation in youth: The roles of gender, race, socioeconomic, and developmental status. *J Adolesc Health* 1998;23:271-9.
 33. Siqueira LM, Rolnitzky LM, Rickert VI. Smoking cessation in adolescents: The role of nicotine dependence, stress, and coping methods. *Arch Pediatr Adolesc Med* 2001;155:489-95.
 34. Baugh JG, Hunter SM, Webber LS, et al. Development trends of first cigarette smoking experience of children: The Bogalusa heart study. *Am J Public Health* 1982;72:1161-4.
 35. Everett SA, Warren CW, Sharp D, et al. Initiation of cigarette smoking and subsequent smoking behavior among U.S. high school students. *Prev Med* 1999;29:327-33.
 36. Breslau N, Kilbey MM, Andreski P. DSM-III-R nicotine dependence in young adults: Prevalence, correlates and associated psychiatric disorders. *Addiction* 1994;89:743-54.
 37. Moolchan ET, Radzius A, Epstein DH, et al. The Fagerstrom Test for Nicotine Dependence and the Diagnostic Interview Schedule: Do they diagnose the same smokers? *Addict Behav* 2002;27:101-13.
 38. Jones D, Schroeder JR, Moolchan ET. Time spent with friends who smoke and number of quit attempts among teen smokers. *Addict Behav* In press.
 39. DeWit DJ, Offord DR, Wong M. Patterns of onset and cessation of drug use over the early part of the life course. *Health Educ Behav* 1997;24:746-58.
 40. Giovino GA. Epidemiology of tobacco use among US adolescents. *Nicotine Tob Res* 1999;S31-40.
 41. Breslau N. Daily cigarette consumption in early adulthood: Age of smoking initiation and duration of smoking. *Drug Alcohol Depend* 1993;33:287-91.
 42. Kandel DB, Chen K. Extent of smoking and nicotine dependence in the United States: 1991-1993. *Nicotine Tob Res* 2000;2:263-74.
 43. DiFranza JR, Savageau JA, Rigotti NA, et al. Development of symptoms of tobacco dependence in youths: 30 month follow up data from the DANDY study. *Tob Control* 2002;11:228-35.
 44. Breslau N, Johnson EO, Hiripi E, et al. Nicotine dependence in the United States: Prevalence, trends, and smoking persistence. *Arch Gen Psychiatry* 2001;58:810-6.
 45. Sargent JD, Dalton M. Does parental disapproval of smoking prevent adolescents from becoming established smokers? *Pediatrics* 2001;108:1256-62.
 46. Emery S, Gilpin EA, White MM, et al. How adolescents get their cigarettes: Implications for policies on access and price. *J Natl Cancer Inst* 1999;91:184-6.
 47. McNeill AD. The development of dependence on smoking in children. *Br J Addict* 1991;86:589-92.
 48. Colby SM, Tiffany ST, Shiffman S, et al. Are adolescent smokers dependent on nicotine? A review of the evidence. *Drug Alcohol Depend* 2000;59(Suppl 1):S83-95.
 49. Blitstein JL, Robinson LA, Murray DM, et al. Rapid progression to regular cigarette smoking among nonsmoking adolescents: Interactions with gender and ethnicity. *Prev Med* 2003;36:455-63.
 50. DiFranza JR, Coleman M. Sources of tobacco for youths in communities with strong enforcement of youth access laws. *Tob Control* 2001;10:323-8.

51. Sargent JD, DiFranza JR. Tobacco control for clinicians who treat adolescents. *CA Cancer J Clin* 2003;53:102–23.
52. Reynolds AJ, Temple JA, Robertson DL, et al. Long-term effects of an early childhood intervention on educational achievement and juvenile arrest: A 15-year follow-up of low-income children in public schools. *JAMA* 2001;285:2339–46.
53. Sargent JD, Mott LA, Stevens M. Predictors of smoking cessation in adolescents. *Arch Pediatr Adolesc Med* 1998;152:388–93.
54. Garvey AJ, Bosse R, Glynn RJ, et al. Smoking cessation in a prospective study of healthy adult males: Effects of age, time period, and amount smoked. *Am J Public Health* 1983;73:446–50.
55. Fiore MC. Treating tobacco use and dependence: an introduction to the US Public Health Service Clinical Practice Guideline. *Respir Care* 2000;45:1196–9.
56. McGrady GA, Ahluwalia JS, Pederson LL. Smoking initiation and cessation in African Americans attending an inner-city walk-in clinic. *Am J Prev Med* 1998;14:130–7.
57. Koepke D FB, Johnson CA. Health behaviors in minority families: The case of cigarette smoking. *Fam Community Health* 1990;13:35–43.
58. Beech BM, Scarinci IC. Smoking attitudes and practices among low-income African-Americans: Qualitative assessment of contributing factors. *Am J Health Promot* 2003;17:240–8.
59. Patrick DL, Cheadle A, Thompson DC, et al. The validity of self-reported smoking: A review and meta-analysis. *Am J Public Health* 1994;84:1086–93.
60. Johnson TP, Mott JA. The reliability of self-reported age of onset of tobacco, alcohol and illicit drug use. *Addiction* 2001;96:1187–98.
61. Conrad KM, Flay BR, Hill D. Why children start smoking cigarettes: Predictors of onset. *Br J Addict* 1992;87:1711–24.