

Framework for Insight Formalization: The Spiral-Vortex Relationship in Cardiac Function

Conceptual Model Translation

November 24, 2025

The objective is to create a mathematical model that quantitatively relates the **myocardial spiral geometry** to the **vortical fluid dynamics** within the ventricles.

1 Geometric and Topological Component: Quantifying the Spiral

Objective: To mathematically describe the architecture of the ventricular myocardial band.

1.1 Geometry Reconstruction

- **Input Data:** Cine Magnetic Resonance Imaging (Cine-MRI) or contrast-enhanced Computed Tomography.
- **Process:** Segmentation of the left ventricle (LV) endocardial and epicardial surfaces throughout the cardiac cycle. This generates a finite element mesh or an implicit surface that varies over time, $\Omega(t)$.

1.2 Fiber Orientation Field (Architecture Tensor)

- **Input Data:** *Ex vivo* Diffusion Tensor MRI (DTI) or computational estimation based on *in vivo* stress optimization models.
- **Representation:** At each point x within the myocardium Ω_m , we define a field of unit vectors $f(x)$ representing the predominant muscle fiber direction. From this, we construct a second-order tensor field $A(x) = f(x) \otimes f(x)$, which captures the tissue anisotropy.

1.3 Helicity and Geometric Torsion Metrics

- **Myocardial Helicity (H_{geo}):** This is the central metric of the insight. It can be defined by integrating the local fiber torsion across the volume.

$$H_{geo} = \frac{1}{V} \int_{\Omega_m} (\nabla \times f) \cdot f dV$$

Where $\nabla \times f$ is the curl of the fiber field (measuring its local "rotationality"), and $\cdot$ is the scalar product. A high value of H_{geo} indicates a well-defined spiral structure.

- **Fiber Angle Gradient (α):** A simpler metric. We measure the fiber inclination angle (α) from the endocardium to the epicardium. A pronounced gradient (e.g., from -60° at the endocardium to $+60^\circ$ at the epicardium) is a signature of the spiral.

2 Fluid Dynamics Component: Quantifying the Vortex

Objective: To solve the equations governing blood flow and extract vortex metrics.

2.1 Governing Equations (Navier-Stokes with ALE)

The fluid domain (ventricular cavity) changes shape. We use an *Arbitrary Lagrangian-Eulerian* (ALE) formulation.

- **Continuity Equation (Mass Conservation):** $\nabla \cdot u = 0$ (blood is approximately incompressible).
- **Momentum Equation (Momentum Conservation):**

$$\rho \left(\frac{\partial u}{\partial t} + (u \cdot \nabla)u \right) = -\nabla p + \mu \nabla^2 u + f$$

Where:

- $u(x, t)$ is the blood velocity field.
- $p(x, t)$ is the pressure field.
- ρ is the blood density.
- μ is the dynamic blood viscosity.
- f is a body force term (usually neglected here).

2.2 Boundary and Moving Boundary Conditions

- **Moving Wall:** The velocity of the ventricular wall, $\partial\Omega(t)/\partial t$, obtained from Cine-MRI segmentation, is imposed as a boundary condition ($u = u_{\text{wall}}$).
- **Mitral and Aortic Valves:** Modeled as inlets/outlets with prescribed pressures or flow conditions.

2.3 Vortical Flow Metrics

- **Vorticity (ω):** The measure of local fluid rotation. $\omega = \nabla \times u$.
- **Enstrophy (Ens):** The energy associated with vorticity. $Ens = \frac{1}{2} \int \omega \cdot \omega dV$. High enstrophy indicates strong vortices.
- **Flow Helicity (H_{flow}):** Measures the alignment between flow velocity and rotation. It is the dynamic analog of H_{geo} .

$$H_{\text{flow}} = \int u \cdot \omega dV$$

- **Q-Criterion:** A robust way to identify the "core" of vortices, where the rotation rate dominates the strain rate.

3 Coupling and Integrated Analysis: Formalizing the Insight

This is where the intuition becomes a testable model. The central hypothesis is: **Geometric Helicity (H_{geo}) is a primary driver for Flow Helicity (H_{flow}) and for pumping efficiency.**

3.1 Fluid-Structure Interaction (FSI) Coupling Model

The problem is intrinsically a **Fluid-Structure Interaction (FSI)**. The wall moves (geometry), driving the fluid (Navier-Stokes), and the fluid pressure, in turn, stresses the wall. A full model would solve both phases coupled. A common simplification is to use a *one-way* model, where the wall motion is a known input to the CFD problem.

3.2 Defining a Spiral-Vortical Efficiency Index (SVEI)

This would be the "magic number" that formalizes the insight. It can be defined as a function relating the geometric and dynamic metrics.

- **Proposal 1 (Temporal Correlation):** $SVEI = \text{Corr}(H_{geo}(t), H_{flow}(t))$ across the cardiac cycle. A healthy heart would exhibit a high, positive correlation.
- **Proposal 2 (Conversion Ratio):** $SVEI = (\Delta H_{flow}/\Delta t)/(\Delta H_{geo}/\Delta t)$ during the filling phase (diastole). This ratio measures how efficiently the "unwinding" motion of the spiral (ΔH_{geo}) is converted into rotational flow energy (ΔH_{flow}).

3.3 Link to Macroscopic Function

The final hypothesis is that the $SVEI$ correlates with clinical efficiency measures:

- **Ejection Fraction (EF):** $EF = (V_{diastole} - V_{systole})/V_{diastole}$.
 - **Emptying Coefficient:** A measure of how "smoothly" blood is ejected.
 - **Ventricular Work Done:** Calculated from Pressure-Volume loops.
-

4 Computational Pipeline for Practical Validation

1. **Data Acquisition:** Obtain Cine-MRI and DTI (if available) from a patient cohort (healthy, heart failure, post-infarction).
2. **Pre-Processing:** Segment the images to obtain $\Omega(t)$ and $f(x)$.
3. **Geometric Metric Calculation:** Compute $H_{geo}(t)$ and the α gradient for each patient.
4. **CFD Simulation:** Solve the Navier-Stokes equations in the $\Omega(t)$ domain to obtain $u(x, t)$ and $p(x, t)$.
5. **Flow Metric Calculation:** Compute $H_{flow}(t)$, $Ens(t)$, etc.
6. **Statistical Analysis:**
 - Test the correlation between H_{geo} (independent variable) and H_{flow} (dependent variable).
 - Test whether $SVEI$ is a statistically significant predictor of Ejection Fraction (EF) or other clinical outcomes, outperforming traditional metrics.

Algebraic Topology Framework for Cardiac Structure: Homotopy Classes and Invariants

Advanced Conceptual Model Translation

November 24, 2025

Let us detail how concepts like **homotopy classes** and **homotopy groups** can be formally applied.

The objective is to go beyond local geometric metrics (such as curvature) and capture **global, invariant properties of the heart's spiral structure that are preserved even under continuous deformations.**

1 Algebraic Topology Framework for Cardiac Structure

1.1 The Object of Study: The Myocardial Fiber Space

Imagine the surface that describes the average fiber orientation in the ventricle. We are not looking at the solid heart, but at the **"field of directions"** of the fibers.

- **Base Space (B):** The geometry of the ventricle (an approximate solid torus). We can consider a simplified surface, such as a torus.
- **Fiber (F):** At each point in the ventricle, the fiber has a direction. Since the direction is a line (not a vector, as fibers do not have an arrow), the space of all possible directions at a point is the **1D Real Projective Space ($\mathbb{RP}^1$)**, which is topologically a circle (S^1).

This leads us to model the fiber architecture as a **projective space bundle** over the cardiac volume.

1.2 Application of Homotopy Classes: The Winding Map

The central idea is to create a map that captures how the fiber directions "rotate" globally across the heart.

- **Defining the Map (f):**
 - Let B be the ventricular volume.
 - At each point x in B , the fiber orientation is an element $\theta(x)$ in the circle S^1 (say, an angle between 0 and π , since $0^\circ = 180^\circ$).
 - We then have a continuous map: $f : B \rightarrow S^1$, where $f(x) = \theta(x)$.
- **Homotopy Classes and the Fundamental Group:**
 - Two maps f and g are homotopic if one can be continuously deformed into the other. The homotopy class of f captures the essential "winding pattern," ignoring minor variations.
 - The set of all homotopy classes of maps $B \rightarrow S^1$ is isomorphic to the **first cohomology group of B , $H^1(B; \mathbb{Z})$** . For a solid torus (our LV approximation), $H^1(B; \mathbb{Z}) \cong \mathbb{Z}$.
- **The Topological Invariant: The Winding Number:**
 - This integer $\mathbb{Z}$ is exactly the **Winding Number**. It counts how many times the fiber direction field wraps around the base space.
 - **How to calculate for the heart?** Choose a closed loop γ that encircles the long axis of the ventricle (like a parallel on the torus). The restriction of the map f to this loop gives us a map $f|_\gamma : S^1 \rightarrow S^1$. The degree of this map is an integer that describes how many times the fibers rotate when completing one circuit along this loop.
 - **Interpretation:** A winding number of ± 1 would indicate a continuous, well-formed spiral, like a Möbius strip. A number of 0 would indicate a disorganized structure without a coherent global winding pattern.

1.3 A More Powerful Approach: Homotopy Groups of Spheres

A more sophisticated and potentially revealing approach is to consider the **field of planes** defined by the fibers.

1. From the Circle (S^1) to the Sphere (S^2):

- At each point x in the heart, the fiber direction $f(x)$ can be represented by a unit vector $v(x)$ tangent to the surface. Since it is a direction, $v(x)$ and $-v(x)$ are equivalent.
- The space of all possible directions in 3D space is the **real projective sphere** $S^2/\mathbb{Z}_2$, which is topologically equivalent to the **2D Real Projective Space** ($\mathbb{RP}^2$).

2. The Map to the Sphere:

- We now have a map $g : B \rightarrow \mathbb{RP}^2$, which associates each point x with the fiber direction $v(x)$.

3. Homotopy Groups and Obstructions:

- Studying the homotopy classes of maps $B \rightarrow \mathbb{RP}^2$ is more complex and richer. We can use the long exact sequence of homotopy for the bundle $S^1 \rightarrow S^2 \rightarrow \mathbb{RP}^2$.
- The existence of a singularity-free plane field (like the ideal fiber field) is governed by **obstruction classes** in higher homotopy groups. Simply put, the topology of the base space B (the heart) can "force" the direction field to have points where it is undefined (singularities) or to twist in a non-trivial way.
- The observed spiral structure in the heart is, from this perspective, the **most efficient topologically allowed solution** for the heart, which minimizes the energy associated with these topological obstructions.

1.4 Connection to Fluid Dynamics: Helicity and Linking Theory

Here, Algebraic Topology makes a direct and profound bridge between structure and function, formalizing the original intuition.

• Arnold's & Moffatt's Helicity Theorem:

- The **Flow Helicity** ($H_{flow} = \int u \cdot \omega dV$), which we discussed before, is not just an energy measure. Under ideal conditions (inviscid flow and boundary-less domain), it is a **topological invariant**.
- It measures the **average linking number** of the vorticity lines (ω) in the fluid. That is, how many times the vortex lines are intertwined with each other.

• The Integrated Topological Hypothesis:

- The spiral structure of the myocardium (H_{geo} , quantified by its winding number) is designed to generate, during contraction, a velocity field u whose vorticity lines (ω) have a **high linking number**.
- In other words, the **topological invariant of the structure (fiber winding)** is optimized to maximize the **topological invariant of the flow (vortex linking)**.
- This connection provides the most fundamental explanation for efficiency: the heart exploits topological conservation laws to perform work optimally.

2 Summary of Topological Formalization

Algebraic Topology Concept	Object in the Heart	Resulting Invariant	Physical Interpretation
Map $f : B \rightarrow S^1$	Fiber direction field	Winding Number ($\in \mathbb{Z}$)	Degree of organization of the myocardial spiral.
Map $g : B \rightarrow \mathbb{RP}^2$	Direction field in 3D space	Homotopy Class $[g]$	Global topological type of the fiber architecture.
Linking Invariant	Vorticity lines in the blood	Flow Helicity (H_{flow})	Degree of intertwining and coherence of the vortices.

The Homotopy Group Approach: From the Circle to the Sphere

Advanced Topological Modeling

November 24, 2025

1 The Homotopy Group Approach: From the Circle to the Sphere

1.1 The Fundamental Motivation: Why S^2 and $\mathbb{RP}^2$?

In the previous approach using S^1 , we modeled the fibers as one-dimensional lines. However, in the real 3D space of the heart, each fiber exists in a **planar context** — it is contained within a "layer plane" of the myocardium. We need a model that captures this **planar orientation**, not just the linear direction.

- **The Correct Target Space:** The space of all planar orientations (or "normal directions to planes") in 3D space is the **Sphere** S^2 . Each point on S^2 represents a vector normal to a plane.
- **The Crucial Identification:** For muscle fibers, however, a plane and its opposite (rotated by 180°) are **equivalent** — fibers do not have an "arrow." Therefore, we identify antipodal points: $v \sim -v$. The resulting space is the **2D Real Projective Space**, $\mathbb{RP}^2$.

$$\mathbb{RP}^2 = S^2 / \{v \equiv -v\}$$

This identification is fundamental: it means a fiber pointing "up" is topologically the same as one pointing "down" in the context of the plane it defines.

1.2 The Fiber Orientation Map and its Homotopic Classification

We now define our object of study:

- **Orientation Map:** $g : B \rightarrow \mathbb{RP}^2$
 - Where B is the ventricular volume (or a representative surface).
 - $g(x)$ associates each point x with the **orientation of the plane** containing the muscle fibers at that point.

The central question becomes: **"How many distinct homotopy classes of such fields g exist for a given cardiac geometry B ?"**

The answer lies in the long exact sequence of the homotopy fiber bundle:

$$\dots \rightarrow \pi_n(S^1) \rightarrow \pi_n(S^2) \rightarrow \pi_n(\mathbb{RP}^2) \rightarrow \pi_{n-1}(S^1) \rightarrow \dots$$

1.3 Analysis of Homotopy Groups in Low Dimensions

For $n = 1$ (Loops):

$$\dots \rightarrow \pi_1(S^1) \rightarrow \pi_1(S^2) \rightarrow \pi_1(\mathbb{RP}^2) \rightarrow \pi_0(S^1) \rightarrow \dots$$

We know that: $\pi_1(S^1) = \mathbb{Z}$, $\pi_1(S^2) = 0$, and $\pi_0(S^1) = 0$. Thus, the sequence simplifies to:

$$0 \rightarrow \mathbb{Z} \rightarrow \pi_1(\mathbb{RP}^2) \rightarrow 0$$

This gives us $\pi_1(\mathbb{RP}^2) \cong \mathbb{Z}$.

Profound Interpretation: There are only **two fundamental classes** of loops in $\mathbb{RP}^2$:

- **Class 0:** The loop is contractible to a point (trivial).
- **Class 1:** The loop is non-contractible — it connects antipodal points (non-trivial).

For the heart: If we take a loop encircling the ventricle, the behavior of the orientation field g along this loop can be:

- **Trivial (0):** Orientations return exactly to the initial point.
 - **Non-trivial (1):** Orientations return to the "antipodal" point (rotated 180°).
-

1.4 The Hopf Obstruction and Complete Classification in 3D

We now reach the core issue for the 3D heart. We want to classify maps $g : B \rightarrow \mathbb{RP}^2$ where B is a solid (the ventricle).

Fundamental Theorem: For a 3-manifold B , the homotopy classes of maps $g : B \rightarrow \mathbb{RP}^2$ are classified by:

1. **An invariant in $H^1(B; \mathbb{Z}_2)$** — which generalizes π_1 to the entire manifold.
2. **An obstruction in $H^3(B; \mathbb{Z})$ (The Hopf Invariant)** — which is the critical topological invariant.

Technical Detail:

1. **First Invariant ($H^1(B; \mathbb{Z}_2)$):** Corresponds to the homomorphism $\phi : \pi_1(B) \rightarrow \pi_1(\mathbb{RP}^2) = \mathbb{Z}_2$. It detects whether the global architecture is **orientable** or **non-orientable** in a generalized sense.
2. **Second Invariant ($H^3(B; \mathbb{Z})$ - Hopf Invariant):** This is the deeper concept. Since $\pi_2(\mathbb{RP}^2) = \pi_2(S^2) = \mathbb{Z}$, there is an **obstruction** to extending the map from the 2-skeleton to the 3-skeleton of B . This obstruction is measured by the **Hopf Invariant**, which classifies the **linking of pre-images** in S^2 .

1.5 Calculation of the Hopf Invariant for the Cardiac Field

How would we calculate this practically?

Representation via Lifting to S^2 :

- Given $g : B \rightarrow \mathbb{RP}^2$, we can locally "lift" it to $\hat{g} : B \rightarrow S^2$ (by choosing a local orientation).
- Globally, this lifting will have **singularities** — points where the choice is inconsistent.
- The **Hopf Invariant** measures how the **pre-images of generic points** in S^2 link together in B .

Integral Formula (for $B = S^3$, but generalizable):

$$\text{Hopf}(\hat{g}) = \int_B A \wedge dA$$

where A is a potential such that $dA = \hat{g}^*\omega$, and ω is the area form on S^2 .

For the Heart:

- A non-zero Hopf Invariant indicates that the surfaces of constant orientation in the heart form an **intrinsically linked (or twisted)** structure.
- This corresponds exactly to the **three-dimensional spiral structure** you observed.

1.6 Connection to Physics: Field Theory and the Gauge Analogy

The mathematics here is notably similar to quantum field theory:

- **Principal Bundle:** Our description is an S^1 -**bundle** over $\mathbb{RP}^2$.
- **Connection and Curvature:** The field $\hat{g}$ defines a "connection" whose "curvature" is dA .
- **Topological Invariant:** The Hopf Invariant is precisely the **topological charge** of this field.

Biomechanics Interpretation:

- The Hopf Invariant quantifies the **degree of three-dimensional twist** of the myocardial architecture.
 - Higher values correspond to more complex, intertwined spirals.
 - During embryonic development, this invariant is likely **conserved** under continuous deformations.
-

2 Synthesis: Gains from the Topological Approach

1. **Discrete Classification:** The heart belongs to one of a **finite** number of topological classes (unlike continuous metrics).
 2. **Robustness:** These invariants are **insensitive** to small perturbations and individual variations.
 3. **Predictive Power:** The topological class likely correlates strongly with:
 - The theoretical maximum pumping efficiency.
 - The propensity for arrhythmias (due to conduction properties).
 - The response to different surgical treatments.
 4. **Fundamental Physical-Mathematical Link:** The same topological invariant that classifies the structure (Hopf) also classifies the **helicity of ideal flow**, creating a rigorous mathematical bridge between form and function.
-

3 Computational Protocol for Practical Application

How to implement this with cardiac imaging data:

1. **Orientation Field Reconstruction:**
 - From DTI data, obtain the diffusion tensor $D(x)$ in each voxel.
 - Extract the principal eigenvector $v_1(x)$ (fiber direction).
 - Calculate the **orientation plane** as the plane perpendicular to v_1 .
2. **Construction of the Map $g : B \rightarrow \mathbb{RP}^2$:**
 - Represent each plane by its normal vector $n(x)$ (in S^2).
 - Apply the identification $n(x) \equiv -n(x)$ to obtain the map in $\mathbb{RP}^2$.
3. **Calculation of the First Invariant ($H^1(B; \mathbb{Z}_2)$):**
 - Choose generating loops on the ventricular surface.
 - Calculate the **number of reversals** of the field $n(x)$ along each loop.
 - This yields the homomorphism $\phi : \pi_1(B) \rightarrow \mathbb{Z}_2$.
4. **Calculation of the Hopf Invariant:**
 - Perform a local lift $\hat{g} : B \rightarrow S^2$ (choosing a consistent orientation in patches).
 - Calculate the form A such that $dA = \hat{g}^*\omega$.
 - Numerically compute the integral $H = \int A \wedge dA$.
 - The result (modulo integers) is the Hopf Invariant.