

FOR THE TREATING VETERINARIAN · EQUICENTA® CTM

Sell the outcome. Your first 60 days.

A clinician's playbook for evaluating eCTM on a first case — what to capture at baseline, what the trial found at each follow-up window, and what to tell the owner.

01

DAYS 1–15

Case selection and baseline.

1. Confirm the case fits the trial population: intra-articular disease, performance workload.
2. Baseline exam: lameness grade, flexion, imaging, workload, owner goal.
3. Pick the endpoint you will report: return-to-work, lameness reduction, or both.
4. Set owner expectations using the language on the right; document informed consent.

02

DAYS 16–30

Injection and early follow-up.

1. Inject per label; record lot #, date, joint, and clinician.
2. 2-week check: lameness re-grade, adverse-reaction screen, owner-reported comfort.
3. 4-week check: return-to-work milestone versus pre-treatment workload.
4. Report any adverse reaction promptly; the trial showed significantly fewer than standard of care.

03

DAYS 31–60

Durability check and next case.

1. 8-week check: performance or kinematic milestone versus baseline.
2. 26-week check: is return-to-work sustained? The trial showed superiority to 26 wk.
3. Compare your outcome to the trial reference: 79% RTW at 26 wk, single injection.
4. Document the case; the next candidate is one that fits the same clinical profile.

What to tell the owner

"eCTM is a cryopreserved umbilical allograft under investigation. In a controlled trial, treated horses returned to work at a higher rate than clinician-selected standard care, with fewer reported adverse reactions and results held to 26 weeks. Single injection."

What to capture at each visit

Baseline and every checkpoint (2, 4, 8, 26 wk): AAEP lameness grade, flexion, workload, owner-reported comfort, adverse-reaction screen. One case sheet per horse. Comparing your outcome to the trial's checkpoints — not to a personal expectation — keeps the conversation objective.

The outcome you are selling

Not "injection given." The outcome is return-to-work sustained to 26 weeks. That is what the trial showed and what the owner will remember. Frame the follow-up cadence, workload progression, and durability check around that endpoint from day one.

Trial reference (Bertone 2026): 138 performance horses, pragmatic controlled design vs. clinician-selected standard care. RTW superiority to 26 wk; significantly fewer reported adverse reactions.

The outcome is the follow-up, not the injection.

Anchor every checkpoint to the trial's endpoints. Return-to-work at 26 weeks is what your owner — and your next case — will remember.

Equine Performance Labs