Equine Performance Labs, LLC
5540-B North Lamar Blvd, Austin, TX 78756
equineperformancelabs.com

WHITE PAPER

Equine Performance Labs
July 7, 2026

Clinical Case Safety Study of an Equine Umbilical-Derived Allograft

equicenta® CTM at ~25 mg/mL (commercially equicenta® MTN) for Joint and Ligament Administration in Horses

Summary of Findings

This case series safety study was a single-dose per site, prospective, unmasked clinical study with post-treatment clinical evaluations of several factors related to safety, including adverse events, lameness or swelling early after injection, and inflammation at the injection site. Veterinarians and Owners/PRs completed scores for lameness/swelling and soft measures of health, respectively, at Baseline (Day 0) and at veterinarian follow up at Day 14, such that each horse served as its own control. Six horses with 11 conditions (10 joints and 1 ligament) with a history of equicenta® CTM injection more than 30 days prior were enrolled within the defined period and completed the study at Day 14 (-3 to +7 days). No adverse events were reported by the veterinarians or the owners/PRs. No inflammation was reported at the injection site (score 0 — no inflammation). No increase in lameness or swelling early after injection was reported. These data, combined with the favorable adverse reaction rate in the main Clinical Study with 128 injections of eCTM at a higher dose, support a favorable adverse event profile for eMTN.

Secondary Observations

Conclusions regarding lameness were not made, given this study was designed for safety. Improvement in swelling did occur in 4 horses and worsened in no horse — a finding that aligned with significantly greater improvement in swelling by two weeks in the tightly controlled clinical study with the higher dose of eCTM. Owner/PR scores for overall health had a small, but positive, change in every category by two weeks, with 3 of the 6 horses improving in performance. The study was designed as a case series for safety to complement the parent Clinical Study on eCTM. Further conclusions are limited by numbers of horses, duration, and lack of contemporaneous controls.

Conclusion

In summary, equicenta® Maintain (eMTN) showed a favorable safety profile both in general and when used as a second follow-up injection to the potent eCTM.