

Notes of **Advanced Physical Chemistry II**

hebrewsnabla

January 9, 2020

Contents

12	Group Theory: the Exploitation of Symmetry	4
12.1	The Exploitation of the Symm of a Mol Can Be Used to Significantly Simplify Numerical Calculations	4
12.2	The Symm of Mols Can Be Described by a Set of Symm Elements	4
	Identity	4
	Rotation	4
	Reflection	4
	Inversion	4
	Rotation Reflection	4
12.2.1	Point Groups of Interest to Chemists	5
12.3	The Symm Operators of a Mol Form a Group	5
12.3.1	Point Group for Some Mols	5
	No Symm Axis	5
	C_n	5
	S_n	5
	C_{nv}	5
	C_{nh}	5
	D_n	5
	D_{nd}	5
	D_{nh}	5
	T_d	5
	O_h	5
	I_h	6
12.4	Symm Operators Can Be Represented by Matrices	6
12.5	The C_{3v} Point Group Has a 2-D Irreducible Representation	6
12.6	The Most Important Summary of the Properties of a Point Group Is Its Character Table	6
	basis	6
	class	6
	notations	6
12.7	Several Mathematical Relations Involve the Characters of Irreducible Representation	6
	notations	6
	order	6
	character	6
	reduce a given reducible repr Γ	7
12.8	Use Symm Arguments to Predict Which Elements in a Secular Det Equals 0 . .	7

12.9	Generating Operators Are Used to Find LCAOs That Are Bases for IrRepr . . .	7
13	Molecular Spectroscopy	7
13.1	7
13.2	Rotational Transitions Accompany Vibrational Transitions	7
13.3	7
13.4	8
13.5	Overtone Are Observed in Vibrational Spectra	8
13.6	Electronic Spectra Contains Electronic, Vibrational and Rotational Info	8
13.7	Franck-Condon Principle Predicts the Relative Intensities of Vibronic Transitions	8
13.8	The Rotational Spectrum of a Polyatomic Mols Depends Upon the Principal Mo- ments of Inertia of the Mol	8
13.9	The Vibrations of Polyatomic Mols Are Represented by Normal Coordinates . .	8
13.10	Normal Coordinates Belong to Irreducible Representations of Mol Point Groups .	8
13.11	Selection Rules Are Derived from TD Perturbation Theory	9
13.12	The Selection Rule in the Rigid-Rotator Approx Is $\Delta J = \pm 1$	9
13.13	The Harmonic-Oscillator Selection Rule Is $\Delta \nu = \pm 1$	9
14	Nuclear Magnetic Resonance Spectroscopy	9
14.1	Nuclei Have Intrinsic Spin Angular Momenta	9
14.2	Magnetic Moments Interact with Magnetic Fields	9
14.3	9
14.4	The Magnetic Field Acting upon Nuclei in Mols Is Shielded	9
14.5	Chemical Shifts Depend upon the Chemical Environment of the Nucleus	9
14.6	Spin-Spin Coupling	9
14.7	Spin-Spin Coupling Between Chemically Equivalent Protons	10
14.8	The $n + 1$ Rule	10
14.9	2nd-Order Spectra	10
15	Lasers, Laser Spectroscopy and Photochemistry	11
15.1	11
15.2	The Dynamics of Spectroscopic Transitions between the Electronic States	11
	absorption	11
	emission	11
15.3	Population Inversion, 2-Level System	11
15.4	Population Inversion, 3-Level System	11
15.5	What is Inside a Laser?	11
15.6	He-Ne Laser	11
15.7	High-Resolution Laser Spectroscopy	11
15.8	The Dynamics of Photochemistry Process	11
16	The Properties of Gases	11
17	The Boltzmann Factor and Partition Functions	11
17.1	11
17.2	Partition Function	11
17.3	Thermodynamic Quantities	12
	17.3.1 Energy	12
	17.3.2 Work and Heat	12
17.4	Pressure	12

17.4.1	Heat Capacity	12
17.4.2	Entropy	12
17.5	12
17.6	The Partition Function of a System of Independent, Distinguishable Mols	12
17.7	The Partition Function of a System of Independent, Indistinguishable Mols . . .	12
17.8	12
18	Partition Functions and Ideal Gases	12
18.1	The Translational PF of Monatomic Ideal Gas	12
18.2	The Electronic PF	12
18.3	12
18.4	The Rotational PF	12
	Symmetry Number	13
25	The Kinetic Theory of Gases	13
25.1	13
25.2	Speed Distribution	13
25.3	Maxwell Distribution	13
25.4	The Frequency of Collisions with a Wall	13
25.5	13
25.6	Inter-collision and MFP	13
26	Chemical Kinetics I: Rate Laws	14
26.8	Reaction Rate Constants	14
27	14
28	Gas Phase Reaction Dynamics	14
28.1	Hard-sphere Collision Theory	14
28.2	Reaction Cross Section Depending on the Impact Parameter	14
	line-of-centers model	14
	Example 28-3	14

Introduction

TA: 刘琼 G403

12 Group Theory: the Exploitation of Symmetry

Matrices

$\det(\mathbf{A}) = 0 \Rightarrow \mathbf{A}$ is a singular matrix.

12.1 The Exploitation of the Symm of a Mol Can Be Used to Significantly Simplify Numerical Calculations

12.2 The Symm of Mols Can Be Described by a Set of Symm Elements

E	
C_n	Rotation by $360^\circ/n$
σ	
i	
S_n	

Table 1: Symmetry elements and operators

Identity

Rotation

σ_h	horizontal
σ_v	vertical
σ_d	diagonal (vertical and bisects the angle between C_2 axis)

Table 2

Reflection

Inversion

Rotation Reflection

$$\hat{S}_n = \hat{\sigma}_h \times \hat{C}_n \quad (12.1)$$

12.2.1 Point Groups of Interest to Chemists

C_{nv}	Rotation by $360^\circ/n$
C_{nh}	
D_{nh}	
D_{nv}	
D_{nd}	
T_d	

Table 3: Symmetry elements and operators

12.3 The Symm Operators of a Mol Form a Group

A set of operators form a group if they satisfy:

1. closed under multiplication 乘法封闭
2. associative multiplication 乘法结合律
3. only one identity operator 单位元
4. everyone has only one inverse 逆元

12.3.1 Point Group for Some Mols

No Symm Axis

C_1 – nothing

C_s – σ

C_i – i

C_n

S_n

C_{nv} – C_n and $n\sigma_v$

C_{nh} – C_n and σ_h

D_n – C_n and $nC_2 \perp C_n$

e.g. 一点点交错的 C_3H_6 , C_2 在 3 个角平分线处

D_{nd} – C_n (also S_{2n}) and $nC_2 \perp C_n$ and $n\sigma_d$

D_{nh} – C_n and $nC_2 \perp C_n$ and σ_h

T_d 主轴是 S_4

O_h

I_h

12.4 Symm Operators Can Be Represented by Matrices

12.5 The C_{3v} Point Group Has a 2-D Irreducible Representation

12.6 The Most Important Summary of the Properties of a Point Group Is Its Character Table

basis

class same characters – in a class.

of class = # of irred representn.

notations

1. $A_1, B_1, E:2D, T:3D$
2. A_1 : symm wrt C_2/σ_v , A_2 : antisymm wrt that.
3. A' : symm wrt σ_h , A'' : antisymm wrt that.
4. A_g, A_u :

12.7 Several Mathematical Relations Involve the Characters of Irreducible Representation

notations

XU G.X.	McQuarrie	
$D^{(\nu)}(R)$		
$\chi^{(\nu)}(R)$	$\chi_j(R)$	
n_ν	d_j	dimension of repr matrix
a_ν	a_j	
g	h	

Table 4

order

$$\sum_{\nu} n_{\nu}^2 = g \quad (12.2)$$

character

$$\sum_R D_{il}^{(\nu)} D_{jm}^{*(\mu)} = \frac{g}{n_{\nu}} \delta_{\mu\nu} \delta_{ij} \delta_{lm} \quad (12.3)$$

$$\sum_R \chi^{(\nu)}(R) \chi^{*(\mu)}(R) = g \delta_{\mu\nu} \quad (12.4)$$

$$\sum_R \chi^{(\nu)}(R) = 0 \quad (\nu \neq A_1) \quad (12.5)$$

reduce a given reducible repr Γ

Suppose

$$\chi(R) = \sum_{\nu} a_{\nu} \chi^{(\nu)}(R) \quad (12.6)$$

thus

$$a_{\nu} = \frac{1}{g} \sum_R \chi(R) \chi^{(\nu)}(R) \quad (12.7)$$

12.8 Use Symm Arguments to Predict Which Elements in a Secular Det Equals 0

12.9 Generating Operators Are Used to Find LCAOs That Are Bases for IrRepr

$$\hat{\mathbf{P}}_j = \frac{d_j}{h} \sum_{\hat{\mathbf{R}}} \chi_j(\hat{\mathbf{R}}) \hat{\mathbf{R}} \quad (12.8)$$

13 Molecular Spectroscopy

13.1

	micro	far IR	IR	visible & UV
f/Hz				
λ/m				
$\tilde{\nu}/\text{cm}^{-1}$				
$E/\text{J mol}^{-1}$				
process				

Table 5

13.2 Rotational Transitions Accompany Vibrational Transitions

Vib & rot energy

$$\tilde{E} = G(\nu) + F(J) \quad (13.1)$$

$$= (v + 1/2)\tilde{\nu} + \tilde{B}J(J + 1) \quad (13.2)$$

selex rule:

$$\Delta v = \pm 1 \quad \Delta J = \pm 1 \quad (13.3)$$

P – left – $\Delta J = -1$ – wide

R – right – $\Delta J = +1$ – narrow

13.3

$$\tilde{B}_v = \tilde{B}_e - \tilde{\alpha}_e(v + 1/2) \quad (13.4)$$

$$\tilde{B}_0 > \tilde{B}_1 > \dots \quad (13.5)$$

which makes P-branches wider.

13.4

$$F(J) = \tilde{B}J(J+1) - \tilde{D}J^2(J+1)^2 \quad (13.6)$$

\tilde{D} :

13.5 Overtones Are Observed in Vibrational Spectra

$$G(v) = \tilde{\nu}_e \left(v + \frac{1}{2} \right) - \tilde{x}_e \tilde{\nu}_e \left(v + \frac{1}{2} \right)^2 \quad (13.7)$$

\tilde{x}_e : anharmonicity cons.

$$\tilde{\nu}_{obs} = \tilde{\nu}_e v - \tilde{x}_e \tilde{\nu}_e v(v+1) \quad (13.8)$$

$v = 1, 2, \dots$

13.6 Electronic Spectra Contains Electronic, Vibrational and Rotational Info

$$\tilde{E} = n\tilde{u}_{el} + \tilde{\nu}_e(v+1/2) - \tilde{x}_e \tilde{\nu}_e(v+1/2)^2 + \dots (rot) \quad (13.9)$$

vibronic transitions:

$0 \rightarrow 0$

$$\tilde{\nu}_{0,0} = \tilde{T}_e + \frac{1}{2}(\tilde{\nu}'_e - \tilde{\nu}''_e) - \frac{1}{4}(\tilde{x}'_e \tilde{\nu}'_e - \tilde{x}''_e \tilde{\nu}''_e) \quad (13.10)$$

13.7 Franck-Condon Principle Predicts the Relative Intensities of Vibronic Transitions

13.8 The Rotational Spectrum of a Polyatomic Mols Depends Upon the Principal Moments of Inertia of the Mol

$$\begin{pmatrix} I_{xx} & I_{xy} & I_{xz} \\ I_{xy} & I_{yy} & I_{yz} \\ I_{xz} & I_{yz} & I_{zz} \end{pmatrix} \xrightarrow{\text{diagonalization}} \begin{pmatrix} I_A & & \\ & I_B & \\ & & I_C \end{pmatrix} \quad (13.11)$$

	top	requisition
$I_C = I_B > I_A = 0$		
$I_C = I_B = I_A$	sph top	$2C_n, n \geq 3$
$I_C = I_B > I_A$	prolate symm top	
$I_C > I_B = I_A$	oblate symm top	
$I_C \neq I_B \neq I_A$	asymm	

Table 6

13.9 The Vibrations of Polyatomic Mols Are Represented by Normal Coordinates

13.10 Normal Coordinates Belong to Irreducible Representations of Mol Point Groups

Contribution to $\chi(R)$ per unmoved atom

$\widehat{\mathbf{R}}$ contribution per unmoved atom
--

Table 7

Now we get Γ_{3N} .

Subtract the irreducible representations corresponding to translational (x, y, z) and rotational (R_x, R_y, R_z) degrees of freedom, we get Γ_{vib} .

13.11 Selection Rules Are Derived from TD Perturbation Theory

Consider a mol interacting w/ EM radiation. The EM field

$$\mathbf{E} = \mathbf{E}_0 \cos 2\pi\nu t \quad (13.12)$$

$$\widehat{H}^{(1)} = -\boldsymbol{\mu} \cdot \mathbf{E} = -\boldsymbol{\mu} \mathbf{E}_0 (\mathrm{e}^{\mathrm{i} 2\pi\nu t} + \mathrm{e}^{-\mathrm{i} 2\pi\nu t})/2 \quad (13.13)$$

$$\Psi(t) = a_1(t)\Psi_1(t) + a_2(t)\Psi_2(t) \quad (13.14)$$

$$a_1(t)\widehat{H}^{(1)}\Psi_1 + a_2(t)\widehat{H}^{(1)}\Psi_2 = \mathrm{i}\hbar \left(\Psi_1 \frac{\mathrm{d}a_1}{\mathrm{d}t} + \Psi_2 \frac{\mathrm{d}a_2}{\mathrm{d}t} \right) \quad (13.15)$$

$$a_1(t) \left\langle \psi_2 \left| \widehat{H}^{(1)} \right| \Psi_1 \right\rangle + a_2(t) \left\langle \psi_2 \left| \widehat{H}^{(1)} \right| \Psi_2 \right\rangle = \mathrm{i}\hbar \left(0 + \frac{\mathrm{d}a_2}{\mathrm{d}t} \mathrm{e}^{-\mathrm{i} E t/\hbar} \right) \quad (13.16)$$

...

$$\mathrm{i}\hbar \frac{\mathrm{d}a_2}{\mathrm{d}t} = \mathrm{e}^{-\mathrm{i}(E_1-E_2)t/\hbar} \left\langle \psi_2 \left| \widehat{H}^{(1)} \right| \psi_1 \right\rangle \quad (13.17)$$

$$\frac{\mathrm{d}a_2}{\mathrm{d}t} \approx \dots \quad (13.18)$$

13.12 The Selection Rule in the Rigid-Rotator Approx Is $\Delta J = \pm 1$

$$\begin{aligned} \langle J', M' | \mu_z | J, M \rangle &= \int_0^{2\pi} \mathrm{d}\phi \int_0^\pi Y_{J'}^{M'*} \mu_z Y_J^M \sin\theta \mathrm{d}\theta \\ &= \dots \end{aligned} \quad (13.19)$$

13.13 The Harmonic-Oscillator Selection Rule Is $\Delta\nu = \pm 1$

14 Nuclear Magnetic Resonance Spectroscopy

14.1 Nuclei Have Intrinsic Spin Angular Momenta

14.2 Magnetic Moments Interact with Magnetic Fields

14.3

14.4 The Magnetic Field Acting upon Nuclei in Mols Is Shielded

14.5 Chemical Shifts Depend upon the Chemical Environment of the Nucleus

14.6 Spin-Spin Coupling

$$\widehat{H} = -\gamma B_0(1 - \sigma_1)\widehat{\mathbf{I}}_{z1} - \gamma B_0(1 - \sigma_2)\widehat{\mathbf{I}}_{z2} + 2\pi J_{12}\widehat{\mathbf{I}}_1 \cdot \widehat{\mathbf{I}}_2 \quad (14.1)$$

$$\hat{H}^{(0)} = -\gamma B_0(1 - \sigma_1)\hat{I}_{z1} - \gamma B_0(1 - \sigma_2)\hat{I}_{z2} \quad \hat{H}^{(1)} = 2\pi J_{12}\hat{I}_1 \cdot \hat{I}_2 \quad (14.2)$$

$$\psi_1^{(0)} = \alpha(1)\alpha(2) \quad \psi_2^{(0)} = \beta(1)\alpha(2) \quad (14.3)$$

$$\psi_3^{(0)} = \alpha(1)\beta(2) \quad \psi_4^{(0)} = \beta(1)\beta(2) \quad (14.4)$$

$$E_1^{(0)} = -\gamma B_0 \left(1 - \frac{\sigma_1 + \sigma_2}{2}\right) \quad E_2^{(0)} = -\gamma B_0(\sigma_1 - \sigma_2) \quad (14.5)$$

$$E_3^{(0)} = \gamma B_0(\sigma_1 - \sigma_2) \quad E_4^{(0)} = \gamma B_0 \left(1 - \frac{\sigma_1 + \sigma_2}{2}\right) \quad (14.6)$$

perturbed to 1st order

$$E_1 = -\gamma B_0 \left(1 - \frac{\sigma_1 + \sigma_2}{2}\right) + \frac{2\pi J_{12}}{4} \quad (14.7)$$

$$E_2 = -\gamma B_0(\sigma_1 - \sigma_2) - \frac{2\pi J_{12}}{4} \quad (14.8)$$

$$E_3 = \gamma B_0(\sigma_1 - \sigma_2) - \frac{2\pi J_{12}}{4} \quad (14.9)$$

$$E_4 = \gamma B_0 \left(1 - \frac{\sigma_1 + \sigma_2}{2}\right) + \frac{2\pi J_{12}}{4} \quad (14.10)$$

Since

$$\nu_0 = \frac{\gamma B_0}{2\pi} \quad (14.11)$$

$$\nu_{1 \rightarrow 2} = \nu_0(1 - \sigma_1) - \frac{J_{12}}{2} \quad (14.12)$$

$$\nu_{1 \rightarrow 3} = \nu_0(1 - \sigma_2) - \frac{J_{12}}{2} \quad (14.13)$$

$$\nu_{2 \rightarrow 4} = \nu_0(1 - \sigma_2) + \frac{J_{12}}{2} \quad (14.14)$$

$$\nu_{3 \rightarrow 4} = \nu_0(1 - \sigma_1) + \frac{J_{12}}{2} \quad (14.15)$$

14.7 Spin-Spin Coupling Between Chemically Equivalent Protons

14.8 The $n + 1$ Rule

14.9 2nd-Order Spectra

Only for the case in which

$$J \ll \nu_0 |\sigma_1 - \sigma_2| \quad (14.16)$$

the $n + 1$ spectra.

15 Lasers, Laser Spectroscopy and Photochemistry

15.1

15.2 The Dynamics of Spectroscopic Transitions between the Electronic States

absorption

$$-\frac{dN_1(t)}{dt} = B_{12}\rho_\nu(\nu_{12})N_1(t) \quad (15.1)$$

emission

15.3 Population Inversion, 2-Level System

15.4 Population Inversion, 3-Level System

15.5 What is Inside a Laser?

15.6 He-Ne Laser

15.7 High-Resolution Laser Spectroscopy

Hyperfine structure

15.8 The Dynamics of Photochemistry Process

def: quantum yield

$$\Phi = \frac{\# \text{ mols undergoing reaction}}{\# \text{ photons absorbed}} \quad (15.2)$$

16 The Properties of Gases

17 The Boltzmann Factor and Partition Functions

17.1

17.2 Partition Function

$$Q = \quad (17.1)$$

$$p_j = \frac{e^{-E_j\beta}}{Q} \quad (17.2)$$

17.3 Thermodynamic Quantities

17.3.1 Energy

17.3.2 Work and Heat

17.4 Pressure

17.4.1 Heat Capacity

17.4.2 Entropy

17.5

17.6 The Partition Function of a System of Independent, Distinguishable Mols

$$Q(N, V, T) = \sum_{i,j,k,\dots} e^{-\beta(\varepsilon_i + \varepsilon_j + \varepsilon_k)} = \sum_i e^{-\beta\varepsilon_i} \sum_j e^{-\beta\varepsilon_j} \dots \quad (17.3)$$

17.7 The Partition Function of a System of Independent, Indistinguishable Mols

$$Q(N, V, T) = \frac{q(V, T)^N}{N!} \quad (17.4)$$

17.8

$$\langle \varepsilon \rangle = \sum_j \frac{\varepsilon_j e^{-\beta\varepsilon_j}}{q} \quad (17.5)$$

$$\varepsilon = \varepsilon_i^{trans} + \varepsilon_j^{rot} + \varepsilon_k^{vib} + \varepsilon_l^{elec} \quad (17.6)$$

$$\pi_{ijkl} = \frac{e^{-\beta\varepsilon_i^{trans}} e^{-\beta\varepsilon_j^{rot}} e^{-\beta\varepsilon_k^{vib}} e^{-\beta\varepsilon_l^{elec}}}{q_{trans} q_{rot} q_{vib} q_{elec}} \quad (17.7)$$

18 Partition Functions and Ideal Gases

18.1 The Translational PF of Monatomic Ideal Gas

$$q_{trans} = \left(\frac{2\pi m k_B T}{h^2} \right)^{3/2} T \quad (18.1)$$

18.2 The Electronic PF

18.3

18.4 The Rotational PF

$$q_{rot} = \sum_{J=0} (2J+1) e^{-J(J+1)\Theta_{rot}/T} \quad (18.2)$$

where

$$\Theta_{rot} = \frac{\hbar^2}{2Ik_B} \quad (18.3)$$

At low Θ_{rot}

$$q_{rot} = \dots = \frac{T}{\Theta_{rot}} \quad (18.4)$$

Symmetry Number

$$q_{\text{rot}} = \frac{T}{\sigma \Theta_{\text{rot}}} \quad (18.5)$$

Spherical top

$$q_{\text{rot}} = \frac{\sqrt{\pi}}{\sigma} \left(\frac{T}{\Theta_{\text{rot}}} \right)^{3/2} \quad (18.6)$$

25 The Kinetic Theory of Gases

25.1

25.2 Speed Distribution

$$f(u_x) = \sqrt{\frac{m}{2\pi k_B T}} e^{-mu_x^2/2k_B T} \quad (25.1)$$

$$\langle u_x^2 \rangle = \frac{k_B T}{m} = \frac{RT}{M} \quad (25.2)$$

25.3 Maxwell Distribution

$$F(u) = 4\pi u^2 \left(\frac{m}{2\pi k_B T} \right)^{3/2} e^{-mu^2/2k_B T} \quad (25.3)$$

$$\langle u \rangle = \sqrt{\frac{8k_B T}{\pi m}} \quad (25.4)$$

$$\langle u^2 \rangle = \frac{3k_B T}{m} \quad (25.5)$$

$$u_{mp} = \sqrt{\frac{2k_B T}{m}} \quad (25.6)$$

$$F(\varepsilon) = \frac{2\pi}{(\pi k_B T)^{3/2}} \varepsilon^{1/2} e^{-\varepsilon/k_B T} \quad (25.7)$$

$$\langle \varepsilon \rangle = \frac{3}{2} k_B T \quad (25.8)$$

25.4 The Frequency of Collisions with a Wall

$$dz = \frac{1}{A} \frac{dN}{dt} \quad (25.9)$$

freq per area

$$z = \frac{\rho}{4} \langle u \rangle \quad (25.10)$$

25.5

25.6 Inter-collision and MFP

$$z_A = \rho \sigma \langle u_r \rangle = \rho \sigma \sqrt{2} \langle u \rangle \quad (25.11)$$

$$l = \frac{u}{z_A} = \frac{1}{\sqrt{2} \rho \sigma} \quad (25.12)$$

$$p(x) dx = \frac{1}{l} e^{-x/l} dx \quad (25.13)$$

26 Chemical Kinetics I: Rate Laws

26.8 Reaction Rate Constants

$$k(T) = \frac{k_B T}{h c^\circ} e^{-\Delta^\ddagger G^\circ / RT} \quad (26.1)$$

27

28 Gas Phase Reaction Dynamics

28.1 Hard-sphere Collision Theory

Naïve hard-sphere rate cons.

$$k = \sigma_{AB} N_A \langle u_r \rangle \quad (\text{dm}^3 \text{mol}^{-1} \text{s}^{-1}) \quad (28.1)$$

experimental

$$k = A e^{-E_a / RT} \quad (28.2)$$

Taking into account the dependence of the rate on $\langle u_r \rangle$

$$\sigma_r(E_r) = \begin{cases} 0 & E_r < E_0 \\ \pi d_{AB}^2 & E_r \geq E_0 \end{cases} \quad (28.3)$$

...

$$k = \sigma_{AB} N_A \langle u_r \rangle e^{-E_0 / k_B T} \left(1 + \frac{E_0}{k_B T} \right) \quad (28.4)$$

28.2 Reaction Cross Section Depending on the Impact Parameter

line-of-centers model

energy dependence of the reaction CS

$$\sigma_r(E_r) = \begin{cases} 0 & E_r < E_0 \\ \pi d_{AB}^2 \left(1 - \frac{E_0}{E_r} \right) & E_r \geq E_0 \end{cases} \quad (28.5)$$

$$k = \sigma_{AB} N_A \langle u_r \rangle e^{-E_0 / k_B T} \quad (28.6)$$

Example 28-3

$$k = A e^{-E_a / k_B T} \Rightarrow E_a = k_B T^2 \frac{d \ln k}{dT} \quad (28.7)$$

...

$$E_a = E_0 + \frac{1}{2} k_B T \quad (28.8)$$

with (28.6),

$$\sigma_{AB} N_A \langle u_r \rangle = A e^{-1/2} \Rightarrow A = \sigma_{AB} N_A \langle u_r \rangle e^{1/2} \quad (28.9)$$