



A Brief Population Analysis on Polarized Water

Shirong Wang

Kuang Yaming Honors School

October 14, 2019





Contents

Non-polarized Water

Water Polarized by Another 2 Water Molecule

Water Polarized by Background Charges

Solvated Water

Summary





Non-polarized Water



Atom charges	Mulliken	NPA	ESP(MK)
O	-0.603	-0.936	-0.724
H	0.302	0.468	0.362
H	0.302	0.468	0.362


Table: Atomic charges (a.u.) on each atom of non-polarized water

All results are calculated under M06-2X/def2-QZVP with Gaussian16.





Case I – Polarized by Another 2 Water Molecule




Atom charges	Mulliken	NPA	ESP(MK)
O	-0.642	-1.011	-0.877
H	0.295	0.486	0.449
H	0.274	0.483	0.379

Table: Atomic charges (a.u.) on each atom of polarized water, Case I





Case II – Polarized by Another 2 Water Molecule




Atom charges	Mulliken	NPA	ESP(MK)
O	-0.655	-0.950	-0.550
H	0.353	0.496	0.358
H	0.352	0.495	0.341

Table: Atomic charges (a.u.) on each atom of polarized water, Case II





Case III – Polarized by Background Charges



Atom charges	Mulliken	NPA	ESP(MK)
O	-0.365	-0.787	-0.523
H	0.720	0.647	0.725
H	-0.354	0.139	-0.202

Table: Atomic charges (a.u.) on each atom of polarized water, Case III



Case IV – Solvated Water

Atom charges	Mulliken	NPA	ESP(MK)
O	-0.640	-0.950	-0.763
H	0.320	0.475	0.381
H	0.320	0.475	0.381


Table: Atomic charges (a.u.) on each atom of water in cyclohexane solvent (PCM model)

The results show that water molecule is a little bit more polarized in cyclohexane.





Summary



Atom	non-polarized	I	II	III	IV
O	0	-0.153	0.174	0.201	-0.039
H	0	0.087	-0.004	0.363	0.019
H	0	0.017	-0.021	-0.564	0.019

Table: changes of ESP charges on each atom (set non-polarized results as zero)



Thank You

