

Catalyzing the Hydrodefluorination of CF₃-Substituted Alkenes by PhSiH₃. H• Transfer from a Nickel Hydride

Chengbo Yao, Shuai Wang, Jack Norton,* and Matthew Hammond

Cite This: <https://dx.doi.org/10.1021/jacs.9b13757>

Read Online

ACCESS |



Metrics & More




Article Recommendations



Supporting Information

ABSTRACT: The hydrodefluorination of CF₃-substituted alkenes can be catalyzed by a nickel(II) hydride bearing a pincer ligand. The catalyst loading can be as low as 1 mol%. *gem*-Difluoroalkenes containing a number of functional groups can be formed in good to excellent yields by a radical mechanism initiated by H• transfer from the nickel hydride. The relative reactivity of various substrates supports the proposed mechanism, as does a TEMPO trapping experiment.



INTRODUCTION

Fluorine chemistry is gaining increasing attention because of the importance of fluorine-containing compounds in medicinal chemistry and agrochemistry.^{1–6} Among fluorine-containing functional groups, *gem*-difluoroalkenes are intriguing, contained in a series of biologically active compounds,^{7–10} and well established as a bioisostere of carbonyl compounds with increased metabolic stability and thus improved pharmaceutical performance.^{11–14} In the case of artemisinin, the replacement of a carbonyl with a *gem*-difluoroalkene gives enhanced antimalarial activity. In some cases, the *gem*-difluoroalkene moiety reverses the regioselectivity of enzyme-catalyzed hydride reduction (Figure 1). *gem*-Difluoroalkenes can also serve as versatile building blocks for the synthesis of other fluorine-containing molecules.^{15–20}

The growing interest in the *gem*-difluoroalkene moiety has led to a number of strategies for its preparation (Scheme 1). The conventional approach relies on functional group interconversion, i.e., the difluoromethylation of carbonyl or diazo compounds (Scheme 1a).^{16,17} However, these functional group interconversion strategies typically involve highly reactive intermediates or harsh reaction conditions, limiting their substrate scope.

There are several ways in which *gem*-difluoroalkenes can be prepared from the readily available^{21–25} trifluoromethyl-substituted alkenes. In one convergent approach, nucleophilic attack on a CF₃ can lead to fluoride loss, but an S_N2' reaction with strong nucleophiles, such as Grignard reagents or organolithium reagents, will suffer from poor functional group tolerance (Scheme 1b). Recently, radical chemistry has been used for the synthesis of *gem*-difluoroalkenes, with defluorina-




Figure 1. Representative applications of *gem*-difluoroalkenes.

tion of CF₃ by either photocatalysis or Ni catalysis (Scheme 1c,d).^{26–36}

Typical Ni-catalyzed defluorinations of trifluoromethyl alkenes for the synthesis of *gem*-difluoroalkenes begin with single electron transfer from the nickel to an alkyl radical precursor. The resulting alkyl radical adds to another CF₃ alkene, producing a new radical which is then quenched by the formation of a Ni–C bond; β -F elimination gives the final

Received: December 20, 2019

Published: January 14, 2020



ACS Publications



© XXXX American Chemical Society

A


<https://dx.doi.org/10.1021/jacs.9b13757>
J. Am. Chem. Soc. XXXX, XXX, XXX–XXX

Scheme 1. Typical Synthetic Routes to *gem*-Difluoroalkenes


a. function group interconversion

b. S_N2' type

c. photocatalysis



d. Ni catalysis




product. Other routes to functionalized *gem*-difluoroalkenes, such as alkenylation,³⁷ arylation,³⁸ and borylation,³⁹ have also been reported.

In general, C–F bond activation provides an easy approach to the synthesis of partially fluorinated compounds from readily available polyfluorinated species.^{15,40,41} The simplest transformation of this sort, hydrodefluorination, has attracted much attention and features a unique mechanistic diversity.^{42–45} However, most hydrodefluorination reactions promoted by transition metals are limited to aromatic or olefinic C–F bonds and show little selectivity among such bonds. The Hiseida group has reported a (Co)₁₂–TiO₂ hybrid catalyst for the photochemical hydrodefluorination of substituted α -CF₃ styrenes,⁴⁶ although a hydrogenation byproduct is always generated along with the *gem*-difluoroalkene. Zhang and co-workers reported a copper-catalyzed reductive defluorination of β -trifluoromethylated enones.⁴⁷ However, the use of Grignard reagents limited its functional group tolerance. Herein, we report that the iso-PmBox Ni(II) hydride **1a** can catalyze the synthesis of *gem*-difluoroalkenes by the hydrodefluorination of trifluoromethyl-substituted alkenes with silanes.

RESULTS AND DISCUSSION

The iso-PmBox nickel hydride system **1a** was developed by, and has been studied by, the Gade group.⁴⁸ It is well established that the Ni(II) hydride is in dynamic equilibrium with Ni(I) metalloradical. The Ni(I) can abstract halides from organic compounds and make Ni(II) halides, from which Ni(II)-H can be regenerated with silanes and boron hydrides.^{49–52}

While investigating hydrogen atom transfer (HAT) from **1a**, we found that it carried out the hydrodefluorination of α -CF₃ styrene **2a** (Scheme 2). During that reaction, the characteristic

Scheme 2. Hydrodefluorination by PhSiH₃ of α -CF₃ Styrene **2a** by isoPmBox Ni(II)-H **1a** in a Stoichiometric and a Catalytic Manner

¹⁹F NMR resonance of the Ni(II)-F complex **1b** was observed at δ –444.3.⁴⁹ Moreover, the disappearance of **2a** (a ¹⁹F singlet at δ –64.50) was accompanied by the appearance of an ABX₃ pattern centered at δ –90.69 (²J_{F,F} = 44.1 Hz, ⁴J_{H,F} = 3.3 Hz), belonging to the *gem*-difluoroalkene **3a**. After the addition of PhSiH₃, the ¹⁹F peak of **1b** disappeared and the ¹H NMR peak of **1a** reappeared. (Et₃SiH did not regenerate **1a**.) Indeed, **1a** was able to catalyze, in quantitative yield (as determined by ¹⁹F NMR) at room temperature, the dehydrofluorination of **2a** with a stoichiometric amount of PhSiH₃.

Table 1 displays the scope of our reaction. Various substituents, either electron-donating or electron-withdrawing, and different substitution patterns on the aromatic ring are well tolerated. All the substrates give yields ranging from good to near quantitative. No substantial amount of hydrogenation products was observed for any of the substrates, demonstrating a satisfying chemoselectivity. A thioether **3c**, an ether **3d**, a tertiary amine **3m**, and the heteroaromatic rings in **3g** and **3p** remain intact. Even the acidic protons of an amide **3e** or the carboxylic acid **3f** do not interfere with the reaction. An exocyclic *gem*-difluoroalkene **3h**, and the 2,2-difluorostyrene **3r**, can be obtained from trisubstituted alkenes bearing a CF₃ substituent, although an elevated temperature is required. Interestingly, only the *E* isomer of the starting material **2r** gives product, with elevated temperature and extended reaction time, while the *Z* isomer remains unreacted.⁵³ A monofluoroalkene **3i** can be obtained from an alkene bearing a difluoromethyl substituent. Nitrile **3j**, ester **3k**, ketone **3n**, and aldehyde **3o**, which are not compatible with Wittig or Julia-type olefinations or with strong nucleophiles in S_N2' -type reactions, are all well tolerated by our method. Product **3l** shows that our reaction can achieve chemoselective activation of the C–F bonds in trifluoromethyl alkenes without attacking an aryl fluoride C–F bond. Other radical stabilizing groups, like a carboalkoxy substituent, can also facilitate the reaction, as shown by the formation of product **3q**. Unfortunately, the reaction does not work on CF₃ alkenes with aliphatic substituents, even at elevated temperatures—a result that is to be expected from the mechanism we propose below.

The control experiment in Table 2 (entry 2) shows that the nickel hydride **1a** is required for the reaction. Attempts at replacing **1a** with metal hydrides previously used in our lab (entry 3), such as HCpCr(CO)₃ and HV(CO)₄(dppe) (dppe = 1,2-bis(diphenylphosphino)ethane), have been unsuccessful,⁵⁴ so the reactivity of **1a** is unique. The catalyst loading can be reduced (entry 4) to 1 mol% without diminishing the yield, although a longer reaction time is necessary. The number of equivalents of PhSiH₃ can be reduced without affecting the yield


Table 1. Substrate Scope of the Nickel-Hydride-Catalyzed Hydrodefluorination of Trifluoromethyl-Substituted Alkenes^a

$\begin{array}{c} \text{R} \\ \\ \text{Ar}-\text{C}=\text{CF}_3 \\ \text{2a-q} \end{array} \xrightarrow[\text{C}_6\text{D}_6, \text{rt, 24h}]{\begin{array}{c} 5 \text{ mol\% } \mathbf{1a} \\ 1 \text{ equiv PhSiH}_3 \end{array}} \begin{array}{c} \text{R} \\ \\ \text{Ar}-\text{C}=\text{CF}_2 \\ \text{3a-q} \end{array}$		

^aIsolated yields, unless otherwise noted. ^b70 °C. ^c50 °C. ^d95 °C, 10 days, only from the *E* isomer of starting material. The yield is determined by ¹⁹F NMR.

(entry 5), which suggests that all three silane hydrides can be used.

Two mechanisms for this reaction seem worth considering. One (shown in the top of Scheme 3) is similar to Gade's proposal for the hydrodefluorination (eq 3) of geminal difluorocyclopropanes.⁴⁹ The Ni(I) (complex 1c) may abstract



an F atom from the substrate **2a** to form the Ni(II) fluoride **1b** and the organic radical **4**; H• transfer from the Ni(II) hydride **1a**


Table 2. Control Experiments^a

$\begin{array}{c} \text{Ph}-\text{C}=\text{CF}_3 \\ \text{2a} \end{array} \xrightarrow[\text{C}_6\text{D}_6, \text{rt, 24h}]{\begin{array}{c} 5 \text{ mol\% } \mathbf{1a} \\ 1 \text{ equiv PhSiH}_3 \end{array}} \begin{array}{c} \text{Ph}-\text{C}=\text{CF}_2 \\ \text{3a} \end{array}$		
entry	deviation from "standard conditions"	yield ^b (%)
1	none	>95
2	no 1a	<5
3	20 mol% HCpCr(CO) ₃ or 20 mol% HV(CO) ₄ (dppe) instead of 1a	<5
4	1 mol% 1a , 72 h	>95
5	0.4 equiv PhSiH ₃	>95


^aAll reactions are performed on 0.5 mmol scale. ^bDetermined by ¹⁹F NMR.

Scheme 3. Two Possible Mechanisms Initiated by Fluorine Atom Abstraction and Hydrogen Atom Transfer, Respectively

Fluorine Atom Abstraction Initiation




Hydrogen Atom Transfer Initiation



will then give the product **3a** and regenerate **1c**, while the silane will reduce the fluoride **1b** back to the hydride **1a**. The other possible mechanism (shown at the bottom of Scheme 3) involves the sort of H• transfer to olefins that we have used to generate radicals for cyclization and isomerization.^{55–58} Transfer to the methylene of **2a** from the hydride **1a** is expected,^{59,60} generating the organic radical **5** while leaving the Ni(I) complex **1c**. Abstraction of an F atom from **5** by **1c** gives the product **3a** and yields the Ni fluoride **1b**,⁶¹ which can be reduced by the silane back to **1a**.

The second mechanism is supported by several lines of evidence. First, it explains why aliphatic alkenes do not work (Scheme 4a), even at an elevated temperature. The aryl group is

Scheme 4. Evidence in Favor of HAT-Initiating Mechanism^a



^aThe structure of **6** has been confirmed by single-crystal X-ray diffraction (Figure 2).

essential for stabilizing the organic radical resulting from HAT, given that CF₃ is a radical destabilizing group;^{62,63} however, the fluorine atom abstraction in the first mechanism would not require an aryl substituent. Second, the slow reaction of trisubstituted alkenes (in Scheme 4b) is more easily explained by the second mechanism—using the established^{60,64} effects of olefin substitution on the rate of HAT to an olefin from a metal hydride. A methyl substituent on the carbon receiving the H• (in the second mechanism) is known to slow HAT by about 3 orders of magnitude, while the rate of fluorine atom abstraction (in the first mechanism) should not change much with the extra substituent on carbon. Third, and the most conclusive, is the successful trapping of the radical **5** by TEMPO (2,2,6,6-tetramethylpiperidin-1-yl)oxyl; Scheme 4c). The addition of 3 equiv of TEMPO to the reaction results in the formation of the TEMPO adduct **6** (Figure 2) in 73% isolated yield.

CONCLUSION

gem-Difluoroalkenes with a variety of functional groups can be generated by the nickel-hydride-catalyzed hydrodefluorination of CF₃ alkenes. The reaction is initiated by H• transfer from Ni to the substrate. Trapping of the radical **5** with TEMPO demonstrates a new mechanism for the previously reported⁴⁸ NNN-pincer nickel(I/II) system.




Figure 2. Molecular structure of TEMPO-adduct **6**. Hydrogen atoms are omitted for clarity.

EXPERIMENTAL SECTION

General Procedures. All manipulations were carried out in an inert atmosphere box (O₂ < 1 ppm) or under Ar by standard Schlenk techniques unless otherwise noted. Glassware was oven-dried or flame-dried prior to use. All commercial reagents were used as received without further purification unless specified. Deuterated benzene (C₆D₆) was distilled from molten potassium and benzophenone ketyl. Benzene (C₆H₆) and tetrahydrofuran (THF) were distilled from sodium-benzophenone ketyl. isoPmbox-Ni(II)-H **1a**,⁴⁸ CpCr(CO)₃H,⁶⁵ HV(CO)₄(dppe),⁶⁶ and Co(dmgBF₂)₂(THF)₂⁶⁷ were synthesized according to the literature procedures and stored in an argon atmosphere glovebox (O₂ < 1 ppm). ¹H NMR, ¹³C NMR and ¹⁹F NMR spectra were recorded using a Bruker 500 Ascend, DRX 500, DRX 400, or DRX 300 spectrometer. Peaks are referenced relative to solvent residual peaks in benzene-*d*₆, THF-*d*₈, CD₃CN, and CDCl₃. The data are reported as follows: chemical shift in parts per million from internal tetramethylsilane on the δ scale, integration, multiplicity (br = broad, s = singlet, d = doublet, t = triplet, q = quartet, p = pentet, m = multiplet), and coupling constants (Hz). High-resolution mass spectra were acquired on a Waters XEVO G2-XS QToF mass spectrometer equipped with a UPC2 SFC inlet and a LockSpray source with one of three probes: electrospray ionization (ESI) probe, atmospheric pressure chemical ionization (APCI) probe, or atmospheric pressure solids analysis probe (ASAP). X-ray diffraction data were collected on a Bruker Apex II diffractometer. Crystal data, data collection and refinement parameters are summarized in Table S1. The structure was solved using direct methods and standard difference map techniques, and was refined by full-matrix least-squares procedures on F² with SHELXTL (Version 2013/4).^{68–70}

General Procedure of NiH-Catalyzed Hydrodefluorination. In an inert atmosphere glovebox, CF₃ substituted alkenes (0.25 or 0.5 mmol), PhSiH₃ (1 equiv), and isoPmbox Ni(II)-H **1a** (0.05 equiv) were weighed in a glass vial and transferred to a J-Young tube using 1 mL of dry and degassed C₆D₆. The reaction was carried out at room temperature for 24 h unless otherwise noted. The crude reaction mixture was directly subjected to flash column chromatography for purification. Spectroscopic details of all the reaction products can be found in the Supporting Information.

Reaction with Other Metal Hydrides. In an inert atmosphere glovebox, (1,1-difluoroprop-1-en-2-yl)benzene **2a** (0.25 mmol), PhSiH₃ (0.25 mmol, 1 equiv), and HCpCr(CO)₃ (10 mg, 0.05 mmol, 0.2 equiv), Co(dmgBF₂)₂(THF)₂ (27 mg, 0.05 mmol, 0.2 equiv), or HV(CO)₄(dppe) (28 mg, 0.05 mmol, 0.2 equiv) were weighed in a glass vial and transferred to a J-Young tube using 1 mL of dry and degassed C₆D₆. The reaction was carried out at room temperature for 24 h. Crude ¹H NMR and ¹⁹F NMR were taken directly or after silica plug.

TEMPO Trapping Experiment. In an inert atmosphere glovebox, (1,1-difluoroprop-1-en-2-yl)benzene **2a** (0.5 mmol), PhSiH₃ (0.5 mmol, 1 equiv), TEMPO (1.5 mmol, 3 equiv), and isoPmbox Ni(II)-H **1a** (0.025 mmol, 0.05 equiv) were weighed in a glass vial and transferred to a J-Young tube using 1 mL of dry and degassed C₆D₆. The reaction was carried out at room temperature for 144 h. The reaction conversion was 56%, 77%, and 89% at 3, 17, and 144 h, respectively. The crude reaction mixture was directly subjected to flash column chromatography for purification. Flash column chromatography was done using pure hexane. Product was obtained with 73% yield.

2,2,6,6-Tetramethyl-1-((1,1,1-trifluoro-2-phenylpropan-2-yl)oxy)piperidine (6). ¹H NMR (400 MHz, chloroform-*d*): δ 7.68–7.62 (m, 2H), 7.46–7.34 (m, 3H), 1.95 (q, *J* = 1.2 Hz, 3H), 1.69–1.50 (m, 3H), 1.47–1.41 (m, 2H), 1.29–1.36 (m, 7H), 1.13 (s, 3H), 0.43 (s, 3H). ¹⁹F NMR (376 MHz, chloroform-*d*) δ –74.83. ¹³C NMR (101 MHz, chloroform-*d*): δ 140.86, 128.27, 127.76, 127.68, 126.00 (q, *J* = 287.6 Hz), 82.54 (q, *J* = 26.4 Hz), 60.98, 60.26, 41.68, 41.56, 33.13, 33.08 (q, *J* = 4.1 Hz), 20.89, 20.80, 16.92, 16.35 (q, *J* = 1.7 Hz). HRMS-ASAP+ (*m/z*): calcd for C₁₈H₂₇F₃NO [M+H]⁺: 330.2045, found: 330.2025.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/jacs.9b13757>.

Starting material preparation, NMR spectra of substrates and products, and 2D NMR characterization of compound **6** (PDF)

X-ray crystallographic data for **6** (CIF)

■ AUTHOR INFORMATION

Corresponding Author

Jack Norton – Department of Chemistry, Columbia University, New York, New York 10027, United States; orcid.org/0000-0003-1563-9555; Email: jrn11@columbia.edu

Authors

Chengbo Yao – Department of Chemistry, Columbia University, New York, New York 10027, United States; orcid.org/0000-0003-4470-7414

Shuai Wang – Department of Chemistry, Columbia University, New York, New York 10027, United States; orcid.org/0000-0001-9464-6758

Matthew Hammond – Department of Chemistry, Columbia University, New York, New York 10027, United States

Complete contact information is available at: <https://pubs.acs.org/doi/10.1021/jacs.9b13757>

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We thank Rebecca Wiles and Gary Molander for providing some of the CF₃ alkenes (**2b–2h** in Table 1). Lutz Gade and Chris Lorenc are acknowledged for helpful discussions. Research reported in this publication was supported by the National

Institute of General Medical Sciences of the National Institutes of Health under Award R01GM124295.

■ REFERENCES

- (1) Kirsch, P. *Modern Fluoroorganic Chemistry: Synthesis, Reactivity, Applications*. Wiley-VCH: Weinheim, Germany, 2004.
- (2) Müller, K.; Faeh, C.; Diederich, F. Fluorine in Pharmaceuticals: Looking Beyond Intuition. *Science* **2007**, *317* (5846), 1881.
- (3) Hagmann, W. K. The Many Roles for Fluorine in Medicinal Chemistry. *J. Med. Chem.* **2008**, *51* (15), 4359–4369.
- (4) Purser, S.; Moore, P. R.; Swallow, S.; Gouverneur, V. Fluorine in medicinal chemistry. *Chem. Soc. Rev.* **2008**, *37* (2), 320–330.
- (5) Zhou, Y.; Wang, J.; Gu, Z.; Wang, S.; Zhu, W.; Aceña, J. L.; Soloshonok, V. A.; Izawa, K.; Liu, H. Next Generation of Fluorine-Containing Pharmaceuticals, Compounds Currently in Phase II–III Clinical Trials of Major Pharmaceutical Companies: New Structural Trends and Therapeutic Areas. *Chem. Rev.* **2016**, *116* (2), 422–518.
- (6) Wang, J.; Sánchez-Roselló, M.; Aceña, J. L.; del Pozo, C.; Sorochinsky, A. E.; Fustero, S.; Soloshonok, V. A.; Liu, H. Fluorine in Pharmaceutical Industry: Fluorine-Containing Drugs Introduced to the Market in the Last Decade (2001–2011). *Chem. Rev.* **2014**, *114* (4), 2432–2506.
- (7) Pan, Y.; Qiu, J.; Silverman, R. B. Design, Synthesis, and Biological Activity of a Difluoro-Substituted, Conformationally Rigid Vigabatrin Analogue as a Potent γ-Aminobutyric Acid Aminotransferase Inhibitor. *J. Med. Chem.* **2003**, *46* (25), S292–S293.
- (8) Altenburger, J.-M.; Lassalle, G. Y.; Matrougui, M.; Galtier, D.; Jetha, J.-C.; Bocskai, Z.; Berry, C. N.; Lunven, C.; Lorrain, J.; Herault, J.-P.; Schaeffer, P.; O'Connor, S. E.; Herbert, J.-M. SSR182289A, a selective and potent orally active thrombin inhibitor. *Bioorg. Med. Chem.* **2004**, *12* (7), 1713–1730.
- (9) Messaoudi, S.; Tréguier, B.; Hamze, A.; Provot, O.; Peyrat, J.-F.; De Losada, J. R.; Liu, J.-M.; Bignon, J.; Wdziejczak-Bakala, J.; Thoret, S.; Dubois, J.; Brion, J.-D.; Alami, M. Isocombretastatins A versus Combretastatins A: The Forgotten isoCA-4 Isomer as a Highly Promising Cytotoxic and Antitubulin Agent. *J. Med. Chem.* **2009**, *52* (14), 4538–4542.
- (10) Bobek, M.; Kawai, I.; De Clercq, E. Synthesis and biological activity of 5-(2,2-difluorovinyl)-2'-deoxyuridine. *J. Med. Chem.* **1987**, *30* (8), 1494–1497.
- (11) Meanwell, N. A. Synopsis of Some Recent Tactical Application of Bioisosteres in Drug Design. *J. Med. Chem.* **2011**, *54* (8), 2529–2591.
- (12) Magueur, G.; Crousse, B.; Ourévitich, M.; Bonnet-Delpon, D.; Bégue, J.-P. Fluoro-artemisinins: When a gem-difluoroethylene replaces a carbonyl group. *J. Fluorine Chem.* **2006**, *127* (4), 637–642.
- (13) Leriche, C.; He, X.; Chang, C.-w. T.; Liu, H.-w. Reversal of the Apparent Regiospecificity of NAD(P)H-Dependent Hydride Transfer: The Properties of the Difluoromethylene Group, A Carbonyl Mimic. *J. Am. Chem. Soc.* **2003**, *125* (21), 6348–6349.
- (14) Zhao, Z.; Liu, H.-w. Synthesis of a Deoxysugar Dinucleotide Containing an exo-Difluoromethylene Moiety As a Mechanistic Probe for Studying Enzymes Involved in Unusual Sugar Biosynthesis. *J. Org. Chem.* **2001**, *66* (20), 6810–6815.
- (15) Fujita, T.; Fuchibe, K.; Ichikawa, J. Transition-Metal-Mediated and -Catalyzed C–F Bond Activation by Fluorine Elimination. *Angew. Chem., Int. Ed.* **2019**, *58* (2), 390–402.
- (16) Zhang, X.; Cao, S. Recent advances in the synthesis and CF functionalization of gem-difluoroalkenes. *Tetrahedron Lett.* **2017**, *58* (5), 375–392.
- (17) Chelucci, G. Synthesis and Metal-Catalyzed Reactions of gem-Dihalovinyl Systems. *Chem. Rev.* **2012**, *112* (3), 1344–1462.
- (18) Hu, J.; Han, X.; Yuan, Y.; Shi, Z. Stereoselective Synthesis of Z Fluoroalkenes through Copper-Catalyzed Hydrodefluorination of gem-Difluoroalkenes with Water. *Angew. Chem., Int. Ed.* **2017**, *56* (43), 13342–13346.
- (19) Hu, J.; Zhao, Y.; Shi, Z. Highly tunable multi-borylation of gem-difluoroalkenes via copper catalysis. *Nat. Catal.* **2018**, *1* (11), 860–869.
- (20) Yoo, W.-J.; Kondo, J.; Rodríguez-Santamaría, J. A.; Nguyen, T. V. Q.; Kobayashi, S. Efficient Synthesis of α-Trifluoromethyl Carboxylic

Acids and Esters through Fluorocarboxylation of gem-Difluoroalkenes. *Angew. Chem., Int. Ed.* **2019**, *58* (20), 6772–6775.

(21) Hamlin, T. A.; Kelly, C. B.; Cywar, R. M.; Leadbeater, N. E. Methylenation of Perfluoroalkyl Ketones using a Peterson Olefination Approach. *J. Org. Chem.* **2014**, *79* (3), 1145–1155.

(22) Jiang, B.; Xu, Y. Trifluoroisopropenylzinc reagent as a useful α -(trifluoromethyl)ethenyl carbanion synthetic equivalent. Preparation and palladium-catalyzed coupling with aryl halides. *J. Org. Chem.* **1991**, *56* (26), 7336–7340.

(23) Kobayashi, O.; Uruguchi, D.; Yamakawa, T. Synthesis of α -trifluoromethylstyrene derivatives via Ni-catalyzed cross-coupling of 2-bromo-3,3,3-trifluoropropene and aryl Grignard reagents. *J. Fluorine Chem.* **2009**, *130* (6), 591–594.

(24) Phelan, J. P.; Wiles, R. J.; Lang, S. B.; Kelly, C. B.; Molander, G. A. Rapid access to diverse, trifluoromethyl-substituted alkenes using complementary strategies. *Chem. Sci.* **2018**, *9* (12), 3215–3220.

(25) Trost, B. M.; Debien, L. Palladium-Catalyzed Trimethylene-methane Cycloaddition of Olefins Activated by the σ -Electron-Withdrawing Trifluoromethyl Group. *J. Am. Chem. Soc.* **2015**, *137* (36), 11606–11609.

(26) Lang, S. B.; Wiles, R. J.; Kelly, C. B.; Molander, G. A. Photoredox Generation of Carbon-Centered Radicals Enables the Construction of 1,1-Difluoroalkene Carbonyl Mimics. *Angew. Chem., Int. Ed.* **2017**, *56* (47), 15073–15077.

(27) Phelan, J. P.; Lang, S. B.; Sim, J.; Berritt, S.; Peat, A. J.; Billings, K.; Fan, L.; Molander, G. A. Open-Air Alkylation Reactions in Photoredox-Catalyzed DNA-Encoded Library Synthesis. *J. Am. Chem. Soc.* **2019**, *141* (8), 3723–3732.

(28) He, Y.; Anand, D.; Sun, Z.; Zhou, L. Visible-Light-Promoted Redox Neutral γ,γ -Difluoroallylation of Cycloketone Oxime Ethers with Trifluoromethyl Alkenes via C–C and C–F Bond Cleavage. *Org. Lett.* **2019**, *21* (10), 3769–3773.

(29) Xia, P.-J.; Ye, Z.-P.; Hu, Y.-Z.; Song, D.; Xiang, H.-Y.; Chen, X.-Q.; Yang, H. Photocatalytic, Phosphoranyl Radical-Mediated N–O Cleavage of Strained Cycloketone Oximes. *Org. Lett.* **2019**, *21* (8), 2658–2662.

(30) Ding, D.; Lan, Y.; Lin, Z.; Wang, C. Synthesis of gem-Difluoroalkenes by Merging Ni-Catalyzed C–F and C–C Bond Activation in Cross-Electrophile Coupling. *Org. Lett.* **2019**, *21* (8), 2723–2730.

(31) Lu, X.; Wang, X.-X.; Gong, T.-J.; Pi, J.-J.; He, S.-J.; Fu, Y. Nickel-catalyzed allylic defluorinative alkylation of trifluoromethyl alkenes with reductive decarboxylation of redox-active esters. *Chem. Sci.* **2019**, *10* (3), 809–814.

(32) Lin, Z.; Lan, Y.; Wang, C. Synthesis of gem-Difluoroalkenes via Nickel-Catalyzed Reductive C–F and C–O Bond Cleavage. *ACS Catal.* **2019**, *9* (1), 775–780.

(33) Lan, Y.; Yang, F.; Wang, C. Synthesis of gem-Difluoroalkenes via Nickel-Catalyzed Allylic Defluorinative Reductive Cross-Coupling. *ACS Catal.* **2018**, *8* (10), 9245–9251.

(34) Xiao, T.; Li, L.; Zhou, L. Synthesis of Functionalized gem-Difluoroalkenes via a Photocatalytic Decarboxylative/Defluorinative Reaction. *J. Org. Chem.* **2016**, *81* (17), 7908–7916.

(35) Chen, H.; Anand, D.; Zhou, L. Photoredox Defluorinative Alkylation of 1-Trifluoromethyl Alkenes and 1,3-Butadienes with 1,4-Dihydropyridines as Alkylation Reagents. *Asian J. Org. Chem.* **2019**, *8* (5), 661–664.

(36) Wiles, R. J.; Phelan, J. P.; Molander, G. A. Metal-free defluorinative arylation of trifluoromethyl alkenes via photoredox catalysis. *Chem. Commun.* **2019**, *55* (53), 7599–7602.

(37) Ichitsuka, T.; Fujita, T.; Ichikawa, J. Nickel-Catalyzed Allylic C(sp³)–F Bond Activation of Trifluoromethyl Groups via β -Fluorine Elimination: Synthesis of Difluoro-1,4-dienes. *ACS Catal.* **2015**, *5* (10), 5947–5950.

(38) Huang, Y.; Hayashi, T. Rhodium-Catalyzed Asymmetric Arylation/Defluorination of 1-(Trifluoromethyl)alkenes Forming Enantioenriched 1,1-Difluoroalkenes. *J. Am. Chem. Soc.* **2016**, *138* (38), 12340–12343.

(39) Liu, Y.; Zhou, Y.; Zhao, Y.; Qu, J. Synthesis of gem-Difluoroallylboronates via FeCl₂-Catalyzed Boration/ β -Fluorine Elimination of Trifluoromethyl Alkenes. *Org. Lett.* **2017**, *19* (4), 946–949.

(40) Ahrens, T.; Kohlmann, J.; Ahrens, M.; Braun, T. Functionalization of Fluorinated Molecules by Transition-Metal-Mediated C–F Bond Activation To Access Fluorinated Building Blocks. *Chem. Rev.* **2015**, *115* (2), 931–972.

(41) Amii, H.; Uneyama, K. C–F Bond Activation in Organic Synthesis. *Chem. Rev.* **2009**, *109* (5), 2119–2183.

(42) Whittlesey, M. K.; Peris, E. Catalytic Hydrodefluorination with Late Transition Metal Complexes. *ACS Catal.* **2014**, *4* (9), 3152–3159.

(43) Kuehnle, M. F.; Lentz, D.; Braun, T. Synthesis of Fluorinated Building Blocks by Transition-Metal-Mediated Hydrodefluorination Reactions. *Angew. Chem., Int. Ed.* **2013**, *52* (12), 3328–3348.

(44) Hu, J.-Y.; Zhang, J.-L., Hydrodefluorination Reactions Catalyzed by Transition-Metal Complexes. In *Organometallic Fluorine Chemistry*, Braun, T., Hughes, R. P., Eds.; Springer International Publishing: Cham, 2015; pp 143–196.

(45) Andrella, N. O.; Xu, N.; Gabidullin, B. M.; Ehm, C.; Baker, R. T. Selective Copper Complex-Catalyzed Hydrodefluorination of Fluoroalkenes and Allyl Fluorides: A Tale of Two Mechanisms. *J. Am. Chem. Soc.* **2019**, *141* (29), 11506–11521.

(46) Tian, H.; Shimakoshi, H.; Imamura, K.; Shiota, Y.; Yoshizawa, K.; Hisaeda, Y. Photocatalytic alkene reduction by a B₁₂–TiO₂ hybrid catalyst coupled with C–F bond cleavage for gem-difluoroolefin synthesis. *Chem. Commun.* **2017**, *53* (68), 9478–9481.

(47) Wu, X.; Xie, F.; Gridnev, I. D.; Zhang, W. A Copper-Catalyzed Reductive Defluorination of β -Trifluoromethylated Enones via Oxidative Homocoupling of Grignard Reagents. *Org. Lett.* **2018**, *20* (6), 1638–1642.

(48) Rettenmeier, C.; Wadepohl, H.; Gade, L. H. Stereoselective Hydrodehalogenation via a Radical-Based Mechanism Involving T-Shaped Chiral Nickel(I) Pincer Complexes. *Chem. - Eur. J.* **2014**, *20* (31), 9657–9665.

(49) Wenz, J.; Rettenmeier, C. A.; Wadepohl, H.; Gade, L. H. Catalytic C–F bond activation of geminal difluorocyclopropanes by nickel(I) complexes via a radical mechanism. *Chem. Commun.* **2016**, *52* (1), 202–205.

(50) Rettenmeier, C. A.; Wenz, J.; Wadepohl, H.; Gade, L. H. Activation of Aryl Halides by Nickel(I) Pincer Complexes: Reaction Pathways of Stoichiometric and Catalytic Dehalogenations. *Inorg. Chem.* **2016**, *55* (16), 8214–8224.

(51) Wenz, J.; Kochan, A.; Wadepohl, H.; Gade, L. H. A Readily Accessible Chiral NNN Pincer Ligand with a Pyrrole Backbone and Its Ni(II) Chemistry: Syntheses, Structural Chemistry, and Bond Activations. *Inorg. Chem.* **2017**, *56* (6), 3631–3643.

(52) Wenz, J.; Wadepohl, H.; Gade, L. H. Regioselective hydrosilylation of epoxides catalyzed by nickel(II) hydrido complexes. *Chem. Commun.* **2017**, *53* (31), 4308–4311.

(53) This difference in reactivity is probably due to a steric effect on the initial HAT step (see the mechanism we propose in the bottom half of Scheme 3).

(54) When substrate **2a** is treated with 20 mol% CpCr(CO)₃H at 70 °C for 24 h, the hydrogenation product is obtained in 23% yield; no hydrodefluorination product is observed. The difference in results between CpCr(CO)₃H and **1a** is probably due to steric effects: the Ni complex **1a** has a bulky ligand environment and thus cannot transfer a second hydrogen atom to **5**. In contrast, CpCr(CO)₃H is able to transfer a second hydrogen atom to give the hydrogenation product. The bulkiness of the Ni hydride **1a** is also demonstrated by the HAT experiments in footnote 59.

(55) Li, G.; Kuo, J. L.; Han, A.; Abuyuan, J. M.; Young, L. C.; Norton, J. R.; Palmer, J. H. Radical Isomerization and Cycloisomerization Initiated by H• Transfer. *J. Am. Chem. Soc.* **2016**, *138* (24), 7698–7704.

(56) Kuo, J. L.; Hartung, J.; Han, A.; Norton, J. R. Direct Generation of Oxygen-Stabilized Radicals by H• Transfer from Transition Metal Hydrides. *J. Am. Chem. Soc.* **2015**, *137* (3), 1036–1039.

(57) Kuo, J. L.; Lorenc, C.; Abuyuan, J. M.; Norton, J. R. Catalysis of Radical Cyclizations from Alkyl Iodides under H₂: Evidence for

Electron Transfer from $[\text{CpV}(\text{CO})_3\text{H}]^-$. *J. Am. Chem. Soc.* **2018**, *140* (13), 4512–4516.

(58) Hartung, J.; Pulling, M. E.; Smith, D. M.; Yang, D. X.; Norton, J. R. Initiating radical cyclizations by H transfer from transition metals. *Tetrahedron* **2008**, *64* (52), 11822–11830.

(59) Compound **1a** is able to catalyze the hydrogenation of styrene or methyl methacrylate in C_6D_6 at 60 °C under 70 psig H_2 . H/D exchange is observed when α -methylstyrene is treated with **1a** under 70 psig D_2 .

(60) Choi, J.; Tang, L.; Norton, J. R. Kinetics of Hydrogen Atom Transfer from $(\eta^5\text{-C}_5\text{H}_5)\text{Cr}(\text{CO})_3\text{H}$ to Various Olefins: Influence of Olefin Structure. *J. Am. Chem. Soc.* **2007**, *129* (1), 234–240.

(61) At this stage we are not sure if the Ni(I) metalloradical **1c** recombines with the carbon-centered radical and then undergoes β -fluorine elimination.

(62) Henry, D. J.; Parkinson, C. J.; Mayer, P. M.; Radom, L. Bond Dissociation Energies and Radical Stabilization Energies Associated with Substituted Methyl Radicals. *J. Phys. Chem. A* **2001**, *105* (27), 6750–6756.

(63) Hioe, J.; Zipse, H. Radical Stability—Thermochemical Aspects. *Encyclopedia of Radicals in Chemistry, Biology and Materials*; Wiley, 2012. DOI: 10.1002/9781119953678.rad012.

(64) Estes, D. P.; Norton, J. R.; Jockusch, S.; Sattler, W. Mechanisms by which Alkynes React with $\text{CpCr}(\text{CO})_3\text{H}$. Application to Radical Cyclization. *J. Am. Chem. Soc.* **2012**, *134* (37), 15512–15518.

(65) Yao, C.; Dahmen, T.; Gansäuer, A.; Norton, J. Anti-Markovnikov alcohols via epoxide hydrogenation through cooperative catalysis. *Science* **2019**, *364* (6442), 764–767.

(66) Choi, J.; Pulling, M. E.; Smith, D. M.; Norton, J. R. Unusually Weak Metal–Hydrogen Bonds in $\text{HV}(\text{CO})_4(\text{P–P})$ and Their Effectiveness as $\text{H}\bullet$ Donors. *J. Am. Chem. Soc.* **2008**, *130* (13), 4250–4252.

(67) Estes, D. P.; Grills, D. C.; Norton, J. R. The Reaction of Cobaloximes with Hydrogen: Products and Thermodynamics. *J. Am. Chem. Soc.* **2014**, *136* (50), 17362–17365.

(68) Sheldrick, G. M. A short history of SHELX. *Acta Crystallogr., Sect. A: Found. Crystallogr.* **2008**, *64*, 112–122.

(69) Sheldrick, G. M. *SHELXTL*, An Integrated System for Solving, Refining, and Displaying Crystal Structures from Diffraction Data; University of Göttingen: Göttingen, Federal Republic of Germany, 1981.

(70) Sheldrick, G. M. Crystal structure refinement with SHELXL. *Acta Crystallogr., Sect. C: Struct. Chem.* **2015**, *71* (1), 3–8.