



NE, DA, and 5-HT neurotransmission influences cortical and subcortical function, suggesting key roles in the neurobiology of ADHD [1][2]

Physiological Function of Monoamine Neurotransmitters




The interplay between NE, DA, and 5-HT



The Optimal Range of Monoaminergic Activity: NE and DA

There is an inverted U-shaped relationship between arousal and task performance, with task-directed behavior requiring optimal regulation of arousal to achieve good performance



References:

- Faraone SV, Bellgrove MA, Brikell I, et al. Attention-deficit/hyperactivity disorder. Nat Rev Dis Primers. 2024;10(1):11. Published 2024 Feb 22. doi:10.1038/s41572-024-00495-0.
- Faraone SV, Radonjić NV. Neurobiology of attention deficit hyperactivity disorder. In: Tasman A, Riba MB, Alarcón RD, et al, eds. Tasman's Psychiatry. Springer, Cham; 2023:1-28.
- da Silva BS, Grevet EH, Silva LCF, Ramos JKN, Rovaris DL, Bau CHD. An overview on neurobiology and therapeutics of attention-deficit/hyperactivity disorder. Discov Ment Health. 2023;3(1):2. Published 2023 Jan 5. doi:10.1007/s44192-022-00030-1.
- Pourhamzeh M, Moravej FG, Arabi M, et al. The roles of serotonin in neuropsychiatric disorders. Cell Mol Neurobiol. 2022;42(6):1671-1692. doi:10.1007/s10571-021-01064-9.
- American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders. 5th ed, text revision. American Psychiatric Association; 2022.
- Koirala S, Grimsrud G, Mooney MA, et al. Neurobiology of attention-deficit hyperactivity disorder: historical challenges and emerging frontiers. Nat Rev Neurosci. 2024;25(12):759-775. doi:10.1038/s41583-024-00869-z.