TITLE:- STABILITY IN THE SHADOWS: MODELING DRIFT OF AUTONOMOUS NODES IN LAGRANGIAN ORBITS

AUTHOR:- ARJUN A.K.A HEET TRIVEDI

EMAIL: - <u>heettrivedio2@gmail.com</u>

Abstract: – This Paper Investigates the dynamics of Orbital Drift of Autonomous Nodes operating near Lagrangian Points (L1 – L5). These are Gravitational Equilibrium zones used by DysonNode Satellites for Orbital Positioning. Even with stabilization protocols, gradual displacement due to micro-gravity, solar pressure, and quantum curvature instabilities leads to drift — Threating Long-term System Autonomy.

Introduction: Lagrangian Points allow Stable orbital positioning with minimal fuel. But nodes placed there are not fully static – they undergo liberation, Chaotic drift, and Non-linear due to;

- ★ 3 Body Gravitational Instabilities
- ★ Solar Wind + Radiation pressure
- ★ Momentum Build-up via micro-thrusts
- ★ Curved Space Warping due to nearby mass activity.

What are lagrangian Points?:- They are points in a two-body gravitational System (e.g., Sun-Earth) where a Third Body can Remain in relative equilibrium.

Points	Туре	Stability
L1, L2, L3	Collinear	Unstable (Require Connections)
L4 / L5	Triangular	Conditionally Stable (But Susceptible to Perturbations)

Drift Dynamics near L1 - L5:- Lagrangian Zones are not flat- they're dynamic potential Wells.

 \bigstar L1 - L3 - > Unstable equilibrium Nodes placed here drift Rapidly without Corrections.

 \bigstar L4 / L5 L - > Stable but can drift under Small Perturbations (e.g., mass loss, radiation).

Drift Grows Exponentially if left Uncontrolled; $\delta r(t) \propto e^{\lambda t}$

Mathematical Orbit Perturbations Model: - Use Hill- Clohessy - Wiltshire equations for Small Oscillations;

$$egin{cases} \ddot{x}-2n\dot{y}-3n^2x=0\ \ddot{y}+2n\dot{x}=0\ \ddot{z}+n^2z=0 \end{cases}$$

Where., x,y,z = Position from equilibrium n = Mean Motion of Central Body

Drift from Solar Radiation Pressure: Solar Photons import tiny but Cumulative force;

$$F = Psolar *A * (1 - P) / C$$

Where., A = Area facing the Sun

P = Reflectivity Coefficient

c = Speed of Light

Drift Momentum Metric:- Define Node Drift Index (NDI);

$$NDI(t) = \sqrt{(x(t) - x0)^2 + (y(t) - y0)^2}$$

If NDI(t) > Rsafe, trigger fallback

Set., Rsafe = 10000 Km

$$\lambda = 1.4 * 10^{-5} 1/second$$

Fallback Conditions:-

Conditions	Trigger
$NDI \ge R$ safe	Auto thrust Vector Correction
Rate of drift $d(NDI) / dt > 3 \text{ km/sec}$	Shutdown Non-linear
Positional Sensor error > 2%	Switch to Redundant Inertial Navigation

Dyson Node Inertial fallback delay: - Node Control Lag» Drift not Corrected Instantly.

Even 1 - 2 second fallback delay at 1 AU - > Micro-Oscillations grow Chaotic over time.

Autonomous Correction Algorithms: - Our Model Includes;

- ★ Predictive Correction using Velocity Curve.
- ★ AI Controlled micro-thruster burst in opposite

Direction.

★ Fuel Less Corrections using Solar Sail orientation.

Curvature Feedback Model (SOLIS Original): - We Integrate Curvature – based drift feedback;

Let;

 $\mathcal{K} = Local Curvature$

Then define;

 $\delta_{drift} \propto \nabla \mathcal{K} * \hat{r}$

That is., If Spacetime Curvature Changes near the node, Drift Direction aligns with Curvature Gradient.

Simulation Parameters: -

parameter	Value
Inertial drift velocity	0.01 m/s
Time Window	6000 Orbital Cycles
Correction Latency	o.2 Second
Radiation Pressure	9 μN/m ²

Fallback Protocols: - If Drift exceeds Critical boundary;

- ★ Switch to Station keeping emergency mode
- ★ Cut off Non-Critical Sensors
- ★ Focus only on Orientation And thrust Correction

Real world parallels:- \star SOHO< \jmath WST, Gaia all perform Constant Corrections around L1/L2.

SOLIS Nodes.

★ L5 Mission (Planned) will need AI Stablizations like

★ The model could literally guide future lagrangian Stabilizers.

Theoretical Hypothesis: - "A dyson node is not in orbit – it's surfing the fabric of Curvature. It's drift is not Random, it is Spacetime's whisper of Imbalance".

— > I propose drift is practically geometric – based not only on Mass and Informational Entropy Gradients in Curved geometry.

Conclusion: - Lagrangian drift is Inevitable – but not uncontrollable. With Real time modeling Curvature based predictions, and predictive Correction, SOLIS Nodes can maintain for decades without failure.

— > This scenario Proves., Autonomy is not about staying balance while drifting inside chaos.

References:-

1. Roy, A. E.

"Orbital Motion," Institute of Physics Publishing, 2004.

2. Farquhar, R.W., Muhonen, D.P., Church, L.C.

"The Trajectory Design of the ISEE-3 Mission," AIAA/AAS Astrodynamics Conference, 1977.

3. NASA Goddard Space Flight Center,

"Lagrange Points: Gateway to Space Exploration," https://solarsystem.nasa.gov

4. Koon, W.S., Lo, M.W., Marsden, J.E., Ross, S.D.

"Dynamical Systems, the Three-Body Problem and Space Mission Design,"

California Institute of Technology, 2000.

5. Szebehely, V. and Giacaglia, G.E.O.

"Stability of L4 and L5 in the Elliptic Restricted Problem of Three Bodies,"

The Astronomical Journal, 1964.

6. ESA Gaia Mission Documentation,

"Orbit Maintenance at L2: Station-Keeping for Billion-Star Mapping,"

https://www.esa.int

7. F.W. Dyson, A.S. Eddington, and C. Davidson,

"A Determination of the Deflection of Light by the Sun's Gravitational Field,"

Philosophical Transactions of the Royal Society, 1920. (Included due to curvature linkage)

8. Szebehely, V.

"Theory of Orbits: The Restricted Problem of Three Bodies," Academic Press, 1967.

9. Project SOLIS — Heet Trivedi (2025)

"Curvature-Based Autonomous Drift Correction Algorithms in Lagrangian-Equilibrium Nodes,"

Self-Published Research Archive, 2025.