

An Exact Cooperation Formula for Introspection Dynamics in the Heterogeneous Public Goods Game

Harry Foster¹, Vince Knight^{1,*}, and Sebastian Krapohl²

¹School of Mathematics, Cardiff University

²Faculty of Social and Behavioural Sciences, University of Amsterdam

April 14, 2026

Abstract

We study introspection dynamics on the heterogeneous public goods game, where N players may differ in their contributions α_i , public goods multipliers r_i , and choice intensities β_i . We prove that the payoff difference a player evaluates when considering a strategy switch is independent of all other players' current actions, a consequence of the linear payoff structure. This state-independence implies that the introspection Markov chain decomposes as a random-scan product of N independent two-state chains, one per player. The stationary distribution is therefore a product measure, and the long-run cooperation probability admits the exact closed form

$$p_C^{\text{intro}} = \frac{1}{N} \sum_{i=1}^N \left[\frac{1 - \mu_{i0} - \mu_{i1}}{1 + e^{\beta_i \alpha_i (1 - r_i/N)}} + \mu_{i0} \right],$$

with no asymptotic approximation. Several structural consequences follow immediately: a player-specific cooperation threshold at $r_i = N$ (under symmetric mutation), payoff-neutrality under zero choice intensity, and the sign of each player's sensitivity to their own parameters.

1 Introduction

The public goods game is a canonical model of cooperation in evolutionary game theory: players decide whether to contribute to a common resource, which is multiplied and redistributed equally, creating a tension between individual incentive and collective benefit [5]. Classical analyses assume a homogeneous population, but real groups are heterogeneous: players differ in their productivity, their contributions, and how sensitively they respond to payoff differences [4].

A natural learning rule for such settings is *introspection dynamics*, introduced in [1]: at each time step a player compares their current payoff to the payoff they *would have received* had they played the alternative action (keeping all other players fixed), and switches with a probability governed by this difference. Because the comparison is self-referential rather than imitative, introspection applies to arbitrary asymmetric games where role-model imitation is unavailable or unnatural.

In [2] the authors extended introspection dynamics to multiplayer asymmetric games and derived a formula for the stationary distribution that can be evaluated numerically for any game. In their general framework the transition rate for player i switching depends on the actions of all other players through the payoff function, so the stationary distribution must be computed by solving a linear system of dimension 2^N ; no closed form is available in general.

In this paper we show that the specific payoff structure of the heterogeneous public goods game admits an exact closed-form analysis. The key observation is that the payoff difference $\Delta\pi_i$ a player evaluates depends only on their own current action and individual parameters, not on the actions of others. This is a consequence of the *linearity* of the public goods payoff in contributions. As a result, the introspection chain decomposes exactly as a product of N independent two-state chains (one per player), the stationary distribution is a product measure, and the cooperation probability p_C^{intro} admits an explicit closed form that scales as $O(N)$ rather than $O(2^N)$.

The paper is organised as follows. Section 2 introduces the model. Section 3 establishes the payoff identity. Section 4 proves the product decomposition and derives the exact cooperation formula. Section 5 records several structural corollaries. Section 6 concludes.

2 Setup and notation

Consider a well-mixed population of N ordered players with contributions $\alpha_1, \dots, \alpha_N > 0$, player-specific public goods multipliers $r_1, \dots, r_N > 1$, player-specific choice intensities $\beta_1, \dots, \beta_N > 0$, and two actions $\{C, D\}$ coded as $\{1, 0\}$. The state space is $S = \{0, 1\}^N$ with $|S| = 2^N$. The payoff to player i in state $\mathbf{a} = (a_1, \dots, a_N)$ is

$$\pi_i(\mathbf{a}) = \frac{r_i}{N} \sum_{j=1}^N \alpha_j a_j - \alpha_i a_i. \quad (1)$$

We write $C(\mathbf{a}) = \sum_j \alpha_j a_j$ for the total contribution. To give the definition of introspection dynamics we define the following two quantities:

- **Fermi function.** Each player i has a personal Fermi function $\phi_i(x) = (1 + e^{\beta_i x})^{-1}$. Since $\beta_i > 0$, the exponential $e^{\beta_i x}$ is strictly increasing in x , so ϕ_i is strictly decreasing with $\phi_i(0) = \frac{1}{2}$ and $\phi_i(x) + \phi_i(-x) = 1$.
- **Mutation.** Each player i has mutation probabilities $\mu_{i0}, \mu_{i1} > 0$ with $\mu_{i0} + \mu_{i1} < 1$, where μ_{i0} is the probability of mutating to cooperation and μ_{i1} is the probability of mutating to defection, regardless of the intended choice of the dynamic. Having $\mu_{i0}, \mu_{i1} > 0$ ensures ergodicity of the Markov chain.

Definition 1 (Cooperation probability). *Given an ergodic chain on S with stationary distribution π , the cooperation probability is $p_C = \pi \cdot s$, where $s_{\mathbf{a}} = \frac{1}{N} \sum_i a_i$ is the fraction of cooperators in state $\mathbf{a}$.*

Introspection dynamics. At each step, one player i is selected uniformly at random. Player i considers switching from action a_i to the alternative $\bar{a}_i = 1 - a_i$. Writing $\mathbf{a}_{i \rightarrow \bar{a}_i}$ for the state obtained by flipping only coordinate i , the transition probability for the switch is

$$T_{\mathbf{a}, \mathbf{a}_{i \rightarrow \bar{a}_i}}^{\text{intro}} = \frac{1}{N} [(1 - \mu_{i0} - \mu_{i1}) \phi_i(\Delta\pi_i) + \mu_{i, \bar{a}_i}], \quad (2)$$

where $\Delta\pi_i := \pi_i(\mathbf{a}) - \pi_i(\mathbf{a}_{i \rightarrow \bar{a}_i})$ is the payoff gain from keeping action a_i , and $\mu_{i, \bar{a}_i}$ denotes μ_{i0} when $\bar{a}_i = C$ and μ_{i1} when $\bar{a}_i = D$. Since ϕ_i is strictly decreasing, a larger payoff gain from the current action yields a smaller switching probability.

3 The payoff identity

Lemma 2 (State-independence of payoff differences). *For the payoff (1), the difference player i evaluates when considering a switch from a_i to $\bar{a}_i$ is*

$$\Delta\pi_i = (2a_i - 1) \alpha_i \left(\frac{r_i}{N} - 1 \right). \quad (3)$$

This depends only on a_i , α_i , r_i , and N ; in particular, it is independent of all other players' actions $\mathbf{a}_{-i}$.

Proof. When player i switches from a_i to $\bar{a}_i = 1 - a_i$, only their own contribution changes, so the total contribution becomes $C(\mathbf{a}_{i \rightarrow \bar{a}_i}) = C(\mathbf{a}) + \alpha_i(1 - 2a_i)$. Expanding both payoffs:

$$\begin{aligned} \pi_i(\mathbf{a}) &= \frac{r_i}{N} C(\mathbf{a}) - \alpha_i a_i, \\ \pi_i(\mathbf{a}_{i \rightarrow \bar{a}_i}) &= \frac{r_i}{N} [C(\mathbf{a}) + \alpha_i(1 - 2a_i)] - \alpha_i(1 - a_i). \end{aligned}$$

Subtracting:

$$\begin{aligned} \Delta\pi_i &= -\frac{r_i}{N} \alpha_i(1 - 2a_i) + \alpha_i(1 - 2a_i) \\ &= \alpha_i(1 - 2a_i) \left(1 - \frac{r_i}{N} \right) \\ &= \alpha_i(2a_i - 1) \left(\frac{r_i}{N} - 1 \right). \end{aligned}$$

The term $C(\mathbf{a})$ cancels entirely, so $\Delta\pi_i$ is independent of $\mathbf{a}_{-i}$. □

Remark 3. *The cancellation of $C(\mathbf{a})$ relies on the payoff (1) being linear in contributions. The heterogeneity of r_i does not affect this cancellation: each player's comparison depends only on their own parameters.*

4 Individual cooperation probabilities

By Lemma 2, player i 's update depends only on their own action a_i , not on $\mathbf{a}_{-i}$. We use this to compute each player's long-run cooperation probability directly from their two-state update rule.

Theorem 4 (Individual cooperation probabilities). *Under introspection dynamics with player-specific β_i , r_i , and mutation probabilities $\mu_{i0}, \mu_{i1} > 0$ with $\mu_{i0} + \mu_{i1} < 1$, and payoff (1), the long-run probability that player i cooperates is*

$$p_i = \phi_i(\alpha_i(1 - r_i/N))(1 - \mu_{i0} - \mu_{i1}) + \mu_{i0}. \quad (4)$$

Proof. By Lemma 2, $\Delta\pi_i$ depends only on a_i . When player i is selected, the probability their next action is C equals p_i regardless of their current action: if $a_i = D$, the switch probability is $\phi_i(\alpha_i(1 - r_i/N))(1 - \mu_{i0} - \mu_{i1}) + \mu_{i0} = p_i$; if $a_i = C$, the probability of remaining at C is $1 - [\phi_i(\alpha_i(r_i/N - 1))(1 - \mu_{i0} - \mu_{i1}) + \mu_{i1}] = p_i$, using $\phi_i(x) + \phi_i(-x) = 1$. Since each selection of player i places them in state C with probability p_i independently of their current state, the long-run cooperation probability of player i is p_i . □

Theorem 5 (Stationary distribution). *The stationary distribution of the introspection chain on $S = \{0, 1\}^N$ is the product measure*

$$\pi_{\mathbf{a}} = \prod_{i=1}^N p_i^{a_i} (1 - p_i)^{1-a_i}, \quad \mathbf{a} \in S, \quad (5)$$

where p_i is given by (4).

Proof. By Theorem 4, each selection of player i places them in state C with probability p_i independently of every other player's action. In stationarity the players' actions are therefore independent, with player i cooperating with probability p_i and defecting with probability $1 - p_i$. The probability of state $\mathbf{a}$ is the product of these marginals, giving (5). □

Corollary 6 (Exact cooperation probability). *The long-run cooperation probability is*

$$p_C^{\text{intro}} = \frac{1}{N} \sum_{i=1}^N \left[\frac{1 - \mu_{i0} - \mu_{i1}}{1 + e^{\beta_i \alpha_i (1 - r_i/N)}} + \mu_{i0} \right]. \quad (6)$$

Proof. By Theorem 4:

$$p_C^{\text{intro}} = \frac{1}{N} \sum_{i=1}^N p_i.$$

Substituting $\phi_i(\alpha_i(1 - r_i/N)) = (1 + e^{\beta_i \alpha_i (1 - r_i/N)})^{-1}$ into (4) gives (6). □

Figure 1 validates Corollary 6 against exact values and simulation across group sizes. Table 1 verifies Theorem 5: for $N = 3$ with per-player multipliers $r_1 = 1$, $r_2 = 3 = N$, $r_3 = 9$, uniform contributions $\alpha_i = 1$, $\beta = 2$, and $\mu_{i0} = \mu_{i1} = 0.1$, the individual cooperation probabilities are $p_1 \approx 0.267$, $p_2 = 0.500$, $p_3 \approx 0.886$, and the formula and exact values of $\pi_{\mathbf{a}}$ agree to five decimal places for all eight states.

State $\mathbf{a}$	Formula $\pi_{\mathbf{a}}$	Exact $\pi_{\mathbf{a}}$
DDD	0.04193	0.04193
DDC	0.32463	0.32462
DCD	0.04193	0.04193
DCC	0.32463	0.32462
CDD	0.01526	0.01527
CDC	0.11818	0.11818
CCD	0.01526	0.01527
CCC	0.11818	0.11818

Table 1: Stationary-distribution probabilities $\pi_{\mathbf{a}}$ for all eight states $\mathbf{a} \in \{D, C\}^3$ with $N = 3$, $\alpha_i = 1$, $\beta = 2$, $\mu_{i0} = \mu_{i1} = 0.1$, and per-player multipliers $r_1 = 1 < N = r_2 = 3 < r_3 = 9$. Formula: product measure (5) (Theorem 5); Exact: values from the 2^N linear system obtained using [3]. Both columns to five decimal places.




Figure 1: Validation of formula (6) on a heterogeneous group with $\alpha_i = i$, $\beta = 0.5$, $\mu_{i0} = \mu_{i1} = 0.1$. *Left:* $r = 2N$ (so $r > N$ for all N). *Right:* $r = N/2$ (so $r < N$ for all N). Orange dots: exact p_C from the full 2^N linear system (feasible only for small N). Blue curve: formula (6) (Corollary 6). Blue bands: p_C estimated from 10^5 simulation steps across 99 independent runs. All values obtained using [3].

5 Structural consequences

The closed form (6) allows immediate structural conclusions about how cooperation depends on game parameters.

Corollary 7 (Player-specific cooperation thresholds). *Player i cooperates with probability greater than $\frac{1}{2}$ if and only if*

$$\phi_i(\alpha_i(1 - r_i/N)) > \frac{\frac{1}{2} - \mu_{i0}}{1 - \mu_{i0} - \mu_{i1}}.$$

When $\mu_{i0} = \mu_{i1}$ this reduces to $r_i > N$. A sufficient condition for $p_C^{\text{intro}} > \frac{1}{2}$ is that the inequality holds for every i .

Proof. Since $1 - \mu_{i0} - \mu_{i1} > 0$:

$$p_i > \frac{1}{2} \iff (1 - \mu_{i0} - \mu_{i1})\phi_i(\alpha_i(1 - r_i/N)) > \frac{1}{2} - \mu_{i0} \iff \phi_i(\alpha_i(1 - r_i/N)) > \frac{\frac{1}{2} - \mu_{i0}}{1 - \mu_{i0} - \mu_{i1}}.$$

When $\mu_{i0} = \mu_{i1}$, the right-hand side equals $\frac{1}{2}$, and since ϕ_i is strictly decreasing with $\phi_i(0) = \frac{1}{2}$, this is equivalent to $\alpha_i(1 - r_i/N) < 0$, i.e. $r_i > N$. The claim about p_C^{intro} follows by averaging. $\square$

Corollary 8 (Neutral drift). *If $\beta_i = 0$ for all i , then game parameters r_i and α_i have no effect on p_C ; cooperation is determined entirely by mutation:*

$$p_i = \frac{1 + \mu_{i0} - \mu_{i1}}{2}, \quad p_C^{\text{intro}} = \frac{1}{2} + \frac{1}{2N} \sum_{i=1}^N (\mu_{i0} - \mu_{i1}).$$

In particular, $p_C^{\text{intro}} = \frac{1}{2}$ if and only if $\sum_i (\mu_{i0} - \mu_{i1}) = 0$.

Proof. When $\beta_i = 0$, the Fermi function satisfies $\phi_i(x) = \frac{1}{2}$ for all x . Substituting into (4):

$$p_i = \frac{1 - \mu_{i0} - \mu_{i1}}{2} + \mu_{i0} = \frac{1}{2} + \frac{\mu_{i0} - \mu_{i1}}{2} = \frac{1 + \mu_{i0} - \mu_{i1}}{2}. \quad \square$$

Corollary 9 (Role of heterogeneity). *For player i with $r_i \neq N$:*

- (i) $\frac{\partial p_i}{\partial \alpha_i}$ is positive when $r_i > N$ and negative when $r_i < N$.
- (ii) $\frac{\partial p_i}{\partial \beta_i}$ is positive when $r_i > N$ and negative when $r_i < N$.

Proof. Write $p_i = (1 - \mu_{i0} - \mu_{i1})\phi_i(\alpha_i(1 - r_i/N)) + \mu_{i0}$.

Part (i). Differentiating with respect to α_i :

$$\frac{\partial p_i}{\partial \alpha_i} = (1 - \mu_{i0} - \mu_{i1})(1 - r_i/N) \phi'_i(\alpha_i(1 - r_i/N)).$$

Since $\phi'_i(x) < 0$ for all x and $1 - \mu_{i0} - \mu_{i1} > 0$, the sign equals $-\text{sgn}(1 - r_i/N) = \text{sgn}(r_i/N - 1)$, which is positive if and only if $r_i > N$.

Part (ii). Write $g(t) = (1 + e^t)^{-1}$, so $\phi_i(x) = g(\beta_i x)$. Differentiating with respect to β_i :

$$\frac{\partial p_i}{\partial \beta_i} = (1 - \mu_{i0} - \mu_{i1}) \alpha_i(1 - r_i/N) g'(\beta_i \alpha_i(1 - r_i/N)).$$

Since $g' < 0$ and $\alpha_i > 0$, the sign equals $-\text{sgn}(1 - r_i/N) = \text{sgn}(r_i/N - 1)$, which is positive if and only if $r_i > N$. $\square$

Remark 10. *Corollaries 7–9 show that each player's long-run cooperation is fully determined by their individual triple (α_i, β_i, r_i) and the group size N . Players with $r_i > N$ cooperate more than half the time, and increasing α_i or β_i raises their cooperation probability further; players with $r_i < N$ show the opposite sensitivity. The parameters of other players have no effect on player i 's marginal cooperation probability.*

6 Conclusion

We have proved that introspection dynamics on the heterogeneous public goods game is exactly solvable. The linear payoff structure implies that each player's payoff comparison is independent of all other players' actions (Lemma 2), so when player i is selected their next action is C with probability p_i regardless of current state (Theorem 4). The long-run cooperation probability therefore has a closed form with N terms rather than 2^N (Corollary 6).




Figure 2: Structural properties of p_i for a single player with $N = 5$, $\mu_{i0} = \mu_{i1} = 0.1$. *Top left:* p_i as a function of β_i ($\alpha_i = 2$) for $r < N$, $r = N$, and $r > N$; all curves pass through $p_i = \frac{1}{2}$ at $\beta_i = 0$ (Corollary 8). *Top centre:* p_i as a function of α_i for two values of r and β_i ; curves above (below) $\frac{1}{2}$ correspond to $r > N$ ($r < N$). *Top right:* $p_i(\alpha_i, r_i)$ at $\beta_i = 0$, confirming $p_i = \frac{1}{2}$ everywhere (Corollary 8); the dashed line marks $r_i = N$. *Bottom row:* Heatmaps of the partial derivatives $\partial p_i / \partial \mu_i$, $\partial p_i / \partial \alpha_i$, and $\partial p_i / \partial \beta_i$ over the (r_i, α_i) or (r_i, β_i) plane ($\beta_i = 0.5$ or $\alpha_i = 2$ as fixed parameter). Red indicates a positive derivative; blue indicates negative. Each derivative changes sign exactly at $r_i = N$ (dashed line), consistent with Corollary 9.

References

- [1] Marta C. Couto, Stefano Giaimo, and Christian Hilbe. Introspection dynamics: a simple model of counterfactual learning in asymmetric games. *New Journal of Physics*, 24(6):063010, 2022.
- [2] Marta C. Couto and Saptarshi Pal. Introspection dynamics in asymmetric multiplayer games. *Dynamic Games and Applications*, 13:1215–1246, 2023.
- [3] Harry Foster and Vince Knight. hefos/ludics: v0.1.1, April 2026.
- [4] Oliver P. Hauser, Christian Hilbe, Krishnendu Chatterjee, and Martin A. Nowak. Social dilemmas among unequals. *Nature*, 572:524–527, 2019.
- [5] Martin A. Nowak. Five rules for the evolution of cooperation. *Science*, 314:1560–1563, 2006.