

# An Exact Cooperation Formula for Introspection Dynamics in the Heterogeneous Public Goods Game

Harry Foster<sup>1</sup>, Vince Knight<sup>1,\*</sup>, and Sebastian Krapohl<sup>2</sup>

<sup>1</sup>School of Mathematics, Cardiff University

<sup>2</sup>Faculty of Social and Behavioural Sciences, University of Amsterdam

May 21, 2026

## Abstract

Cooperation in heterogeneous groups, where individuals differ in resources, productivity, and behavioural responsiveness, underpins collective action across many social and biological systems. Introspection dynamics, in which each player compares their payoff to what they would have received under the alternative action, provides a natural learning rule for such asymmetric settings. We study introspection dynamics on multiplayer games in which the payoff difference  $\Delta f_i$  evaluated by a player when considering a strategy switch is independent of all other players' current actions, a property we call *state-independence*. Under this condition the introspection Markov chain decomposes as a random-scan product of  $N$  independent two-state chains, one per player, and the stationary distribution is a product measure. As our main application we consider the heterogeneous public goods game, where  $N$  players may differ in their contributions  $\alpha_i$ , public goods multipliers  $r_i$ , and choice intensities  $\beta_i$ . We prove that the linear payoff structure implies state-independence, and the long-run cooperation probability admits the exact closed form

$$p_C^{\text{intro}} = \frac{1}{N} \sum_{i=1}^N \left[ \frac{1 - \mu_{i0} - \mu_{i1}}{1 + e^{\beta_i \alpha_i (1 - r_i/N)}} + \mu_{i0} \right],$$

with no asymptotic approximation. Several structural consequences follow immediately: a player-specific cooperation threshold at  $r_i = N$  (under symmetric mutation), payoff-neutrality under zero choice intensity, and the sign of each player's sensitivity to their own parameters.

## 1 Introduction

The public goods game is a canonical model of cooperation in evolutionary game theory: players decide whether to contribute to a common resource, which is multiplied and redistributed, creating a tension between individual incentive and collective benefit [5]. Classical analyses assume a homogeneous population, but real groups are heterogeneous: players differ in their productivity, their contributions, and how sensitively they respond to payoff differences [4].

A natural learning rule for such settings is *introspection dynamics*, introduced in [1]: at each time step a player compares their current payoff to the payoff they *would have received* had they played the alternative action (keeping all other players fixed), and switches with a probability governed by this difference. Because the comparison is self-referential rather than imitative, introspection applies to arbitrary asymmetric games where role-model imitation is unavailable or unnatural.

In [2] the authors extended introspection dynamics to multiplayer asymmetric games and derived a formula for the stationary distribution that can be evaluated numerically for any game. In their general framework the transition rate for player  $i$  switching depends on the actions of all other players through the payoff function, so the stationary distribution must be computed by solving a linear system of dimension  $2^N$ ; no closed form is available in general.

In this paper we show that whenever the payoff difference  $\Delta f_i$  a player evaluates when considering a switch is independent of all other players' current actions, a property we call *state-independence*, the introspection chain decomposes exactly as a product of  $N$  independent two-state chains (one per player), the stationary distribution is a product measure, and the cooperation probability  $p_C^{\text{intro}}$  admits an explicit closed form scaling as  $O(N)$  rather than  $O(2^N)$ . We then show that the heterogeneous public goods game (PGG) satisfies state-independence as a consequence of the *linearity* of its payoff structure, yielding the exact formula (8).

The paper is organised as follows. Section 2 introduces the general framework and the state-independence condition, noting that whilst the donation-form Prisoner's Dilemma is state-independent, the Stag Hunt and the classical (RPST) Prisoner's Dilemma are not. Section 3 proves, for any state-independent game, that individual cooperation probabilities are given by (4) and the stationary distribution is a product measure. Section 4 establishes state-independence for the PGG and derives the exact cooperation formula. Section 5 records several structural corollaries. Section 6 concludes.

## 2 Setup and notation

Consider  $N$  ordered players, each choosing from two actions  $\{C, D\}$  coded as  $\{1, 0\}$ . The state space is  $S = \{0, 1\}^N$  with  $|S| = 2^N$ . Each player  $i$  has a payoff function  $f_i : S \rightarrow \mathbb{R}$  and player-specific choice intensity  $\beta_i > 0$ . Writing  $\mathbf{a}_{i \rightarrow \bar{a}_i}$  for the state with player  $i$ 's action flipped to  $\bar{a}_i = 1 - a_i$ , the *payoff difference* for player  $i$  at state  $\mathbf{a}$  is

$$\Delta f_i(\mathbf{a}) := f_i(\mathbf{a}) - f_i(\mathbf{a}_{i \rightarrow \bar{a}_i}).$$

A payoff function  $f_i$  is *state-independent* if  $\Delta f_i(\mathbf{a})$  depends only on  $a_i$ , not on  $\mathbf{a}_{-i}$ , the actions of all individuals other than  $i$ . When this holds there is a constant  $\delta_i \in \mathbb{R}$  such that

$$\Delta f_i(\mathbf{a}) = (1 - 2a_i) \delta_i, \tag{1}$$

where  $\delta_i = \Delta f_i|_{a_i=0}$  is the payoff difference when player  $i$  defects. When  $a_i = 0$ , (1) follows directly from the definition of  $\delta_i$ ; when  $a_i = 1$ ,  $\Delta f_i|_{a_i=1} = -\Delta f_i|_{a_i=0} = -\delta_i$ , giving (1).

Note that not all games lead to *state-independent* payoff functions. For example in [1] two of the 2 player, 2 strategy games considered to illustrate introspection dynamics are: the Prisoner's Dilemma (2 -  $M_1$ ) and the Stag-hunt (2 -  $M_2$ ) for  $0 < c_1, c_2 < b$ .

$$M_1 = \begin{pmatrix} b - c_1, b - c_2 & -c_1, b \\ b, -c_2 & 0, 0 \end{pmatrix} \quad M_2 = \begin{pmatrix} b - c_1, b - c_2 & -c_1, 0 \\ 0, -c_2 & 0, 0 \end{pmatrix} \tag{2}$$

For  $M_1$ :  $\delta_1 = c_1$  and  $\delta_2 = c_2$  however for  $M_2$ ,  $\Delta f_1((1, 1)) = b - c_1$  but  $\Delta f_1((1, 0)) = -c_1$ . The Prisoner's Dilemma (as defined by  $M_1$ ) is *state-independent* but the Stag-hunt game (as defined by  $M_2$ ) is not. Further games considered in [1] are the Volunteer's dilemma and the Volunteer's timing dilemma (a generalisation with more than 2 strategies) which are both not *state-independent*.

More generally, the classical Prisoner's Dilemma is parameterised by the reward  $R$ , temptation  $T$ , sucker payoff  $S$ , and punishment  $P$  with  $T > R > P > S$  and the social-efficiency condition

$2R > T + S$ . State-independence for player 1 requires the *equality*

$$T - R = P - S \quad (\text{equivalently } P + R = S + T),$$

that is, the gain from defecting against a cooperator must equal the gain from defecting against a defector. The PD axioms are strict inequalities and do not force this equality; a generic RPST Prisoner's Dilemma therefore fails it. For instance,  $T = 5$ ,  $R = 3$ ,  $P = 1$ ,  $S = 0$  satisfies  $T > R > P > S$  and  $2R = 6 > 5 = T + S$ , yet  $T - R = 2 \neq 1 = P - S$ . The donation form  $M_1$  above is the exceptional case where the equality holds by construction (both differentials equal  $c$ ).

To define introspection dynamics we introduce the following two quantities:

- **Fermi function.** Each player  $i$  has a personal Fermi function  $\phi_i(x) = (1 + e^{\beta_i x})^{-1}$ . Since  $\beta_i > 0$ , the exponential  $e^{\beta_i x}$  is strictly increasing in  $x$ , so  $\phi_i$  is strictly decreasing with  $\phi_i(0) = \frac{1}{2}$  and  $\phi_i(x) + \phi_i(-x) = 1$ .
- **Mutation.** Each player  $i$  has mutation probabilities  $\mu_{i0}, \mu_{i1} > 0$  with  $\mu_{i0} + \mu_{i1} < 1$ , where  $\mu_{i0}$  is the probability of mutating to cooperation and  $\mu_{i1}$  is the probability of mutating to defection, regardless of the intended choice of the dynamic.

**Introspection dynamics.** At each step, one player  $i$  is selected uniformly at random. Player  $i$  considers switching from action  $a_i$  to the alternative  $\bar{a}_i = 1 - a_i$ . The transition probability for the switch is

$$T_{\mathbf{a}, \mathbf{a}_i \rightarrow \bar{a}_i}^{\text{intro}} = \frac{1}{N} [(1 - \mu_{i0} - \mu_{i1})\phi_i(\Delta f_i(\mathbf{a})) + \mu_{i, \bar{a}_i}], \quad (3)$$

where  $\mu_{i, \bar{a}_i}$  denotes  $\mu_{i0}$  when  $\bar{a}_i = C$  and  $\mu_{i1}$  when  $\bar{a}_i = D$ . Since  $\phi_i$  is strictly decreasing, a larger payoff gain from keeping the current action yields a smaller switching probability.

**Definition 1** (Cooperation probability). *Given an ergodic chain on  $S$  with stationary distribution  $\pi$ , the cooperation probability is  $p_C = \pi \cdot s$ , where  $s_{\mathbf{a}} = \frac{1}{N} \sum_i a_i$  is the fraction of cooperators in state  $\mathbf{a}$ .*

### 3 Individual cooperation probabilities

For a state-independent game,  $\Delta f_i(\mathbf{a})$  depends only on  $a_i$ , not on  $\mathbf{a}_{-i}$ . We use this to compute each player's long-run cooperation probability and the stationary distribution directly from the two-state update rule.

**Theorem 1** (Individual cooperation probabilities). *For any state-independent game with constants  $\delta_i$ , under introspection dynamics with player-specific choice intensity  $\beta_i$  and mutation probabilities  $\mu_{i0}, \mu_{i1} > 0$  with  $\mu_{i0} + \mu_{i1} < 1$ , the long-run probability that player  $i$  cooperates is*

$$p_i = \phi_i(\delta_i)(1 - \mu_{i0} - \mu_{i1}) + \mu_{i0}. \quad (4)$$

*Proof.* By state-independence,  $\Delta f_i(\mathbf{a}) = (1 - 2a_i)\delta_i$  depends only on  $a_i$ . When player  $i$  is selected, the probability their next action is  $C$  equals  $p_i$  regardless of their current action: if  $a_i = D$ , the switch probability is  $p_i = \phi_i(\delta_i)(1 - \mu_{i0} - \mu_{i1}) + \mu_{i0}$ ; if  $a_i = C$ , the probability of remaining at  $C$  is  $1 - [\phi_i(-\delta_i)(1 - \mu_{i0} - \mu_{i1}) + \mu_{i1}]$ , using  $\phi_i(x) + \phi_i(-x) = 1$  this gives:  $1 - [(1 - \phi_i(\delta_i))(1 -$

$\mu_{i0} - \mu_{i1} + \mu_{i1}] = p_i$ . Since each selection of player  $i$  places them in state  $C$  with probability  $p_i$  independently of their current state, the long-run cooperation probability of player  $i$  is  $p_i$ .  $\square$

**Theorem 2** (Stationary distribution). *For any state-independent game with constants  $\delta_i$ , the stationary distribution of the introspection chain on  $S = \{0, 1\}^N$  is the product measure*

$$\pi_{\mathbf{a}} = \prod_{i=1}^N p_i^{a_i} (1 - p_i)^{1-a_i}, \quad \mathbf{a} \in S, \quad (5)$$

where  $p_i$  is given by (4).

*Proof.* By Theorem 1, each selection of player  $i$  places them in state  $C$  with probability  $p_i$  independently of every other player's action. In stationarity the players' actions are therefore independent, with player  $i$  cooperating with probability  $p_i$  and defecting with probability  $1 - p_i$ . The probability of state  $\mathbf{a}$  is the product of these marginals, giving (5).  $\square$

**Remark 3.** *For the Prisoner's Dilemma  $M_1$  (2) of [1] the payoff  $f_1(\mathbf{a}) = -c_1 a_1 + b a_2$  is separable with  $\delta_1 = c_1$ , and similarly  $\delta_2 = c_2$ . With  $b = 1$ ,  $c_1 = 0.6$ ,  $c_2 = 0.1$ ,  $\beta = 5$ , and  $\mu_{ij} = 0$ , Theorems 1 and 2 give*

$$p_1 = \frac{1}{1 + e^3} \approx 0.047, \quad p_2 = \frac{1}{1 + e^{0.5}} \approx 0.378,$$

and the stationary distribution over  $\{DD, DC, CD, CC\}$  is the product measure

$$\pi_{DD} \approx 0.593, \quad \pi_{DC} \approx 0.360, \quad \pi_{CD} \approx 0.030, \quad \pi_{CC} \approx 0.018.$$

## 4 The heterogeneous public goods game

The *heterogeneous public goods game* is played by  $N$  players with player-specific contributions  $\alpha_1, \dots, \alpha_N > 0$  and public goods multipliers  $r_1, \dots, r_N > 1$ . The payoff to player  $i$  in state  $\mathbf{a} = (a_1, \dots, a_N)$  is

$$f_i(\mathbf{a}) = \frac{r_i}{N} \sum_{j=1}^N \alpha_j a_j - \alpha_i a_i. \quad (6)$$

We write  $C(\mathbf{a}) = \sum_j \alpha_j a_j$  for the total contribution.

**Lemma 4** (State-independence of the PGG). *The payoff (6) is state-independent for every player  $i$ , with*

$$\delta_i = \alpha_i \left(1 - \frac{r_i}{N}\right), \quad (7)$$

i.e.  $\Delta f_i(\mathbf{a}) = (1 - 2a_i) \alpha_i (1 - r_i/N)$ , independent of  $\mathbf{a}_{-i}$ .

*Proof.* When player  $i$  switches from  $a_i$  to  $\bar{a}_i = 1 - a_i$ , only their own contribution changes, so the total contribution becomes  $C(\mathbf{a}_{i \rightarrow \bar{a}_i}) = C(\mathbf{a}) + \alpha_i (1 - 2a_i)$ . Expanding both payoffs:

$$\begin{aligned} f_i(\mathbf{a}) &= \frac{r_i}{N} C(\mathbf{a}) - \alpha_i a_i, \\ f_i(\mathbf{a}_{i \rightarrow \bar{a}_i}) &= \frac{r_i}{N} [C(\mathbf{a}) + \alpha_i (1 - 2a_i)] - \alpha_i (1 - a_i). \end{aligned}$$

Subtracting:

$$\begin{aligned}\Delta f_i &= -\frac{r_i}{N}\alpha_i(1-2a_i) + \alpha_i(1-2a_i) \\ &= \alpha_i(1-2a_i)\left(1 - \frac{r_i}{N}\right).\end{aligned}$$

The term  $C(\mathbf{a})$  cancels entirely, confirming state-independence with  $\delta_i = \alpha_i(1 - r_i/N)$ .  $\square$

**Corollary 5** (Exact cooperation probability). *For the heterogeneous public goods game (6), the long-run cooperation probability is*

$$p_C^{\text{intro}} = \frac{1}{N} \sum_{i=1}^N \left[ \frac{1 - \mu_{i0} - \mu_{i1}}{1 + e^{\beta_i \alpha_i (1 - r_i/N)}} + \mu_{i0} \right]. \quad (8)$$

*Proof.* By Lemma 4,  $\delta_i = \alpha_i(1 - r_i/N)$ . By Theorem 1:  $p_C^{\text{intro}} = \frac{1}{N} \sum_{i=1}^N p_i$ . Substituting  $\phi_i(\delta_i) = (1 + e^{\beta_i \alpha_i (1 - r_i/N)})^{-1}$  into (4) gives (8).  $\square$

Figure 1 validates Corollary 5 against exact values and simulation across group sizes. Table 1 verifies Theorem 2: for  $N = 3$  with per-player multipliers  $r_1 = 1$ ,  $r_2 = 3 = N$ ,  $r_3 = 9$ , uniform contributions  $\alpha_i = 1$ ,  $\beta = 2$ , and  $\mu_{i0} = \mu_{i1} = 0.1$ , the individual cooperation probabilities are  $p_1 \approx 0.267$ ,  $p_2 = 0.500$ ,  $p_3 \approx 0.886$ , and the formula and exact values of  $\pi_{\mathbf{a}}$  agree to five decimal places for all eight states.




Figure 1: Validation of formula (8) on a heterogeneous group with  $\alpha_i = i$ ,  $\beta = 0.5$ ,  $\mu_{i0} = \mu_{i1} = 0.1$ . *Left:*  $r = 2N$  (so  $r > N$  for all  $N$ ). *Right:*  $r = N/2$  (so  $r < N$  for all  $N$ ). Orange dots: exact  $p_C$  from the full  $2^N$  linear system (feasible only for small  $N$ ). Blue curve: formula (8) (Corollary 5). Blue boxes: distribution of  $p_C$  across 99 independent runs of  $10^5$  simulation steps each (outliers shown). The negligible variability across runs reflects the large number of simulation steps per run. All values obtained using [3].

| State $\mathbf{a}$ | Formula $\pi_{\mathbf{a}}$ | Exact $\pi_{\mathbf{a}}$ |
|--------------------|----------------------------|--------------------------|
| DDD                | 0.04193                    | 0.04193                  |
| DDC                | 0.32463                    | 0.32462                  |
| DCD                | 0.04193                    | 0.04193                  |
| DCC                | 0.32463                    | 0.32462                  |
| CDD                | 0.01526                    | 0.01527                  |
| CDC                | 0.11818                    | 0.11818                  |
| CCD                | 0.01526                    | 0.01527                  |
| CCC                | 0.11818                    | 0.11818                  |

Table 1: Stationary-distribution probabilities  $\pi_{\mathbf{a}}$  for all eight states  $\mathbf{a} \in \{D, C\}^3$  with  $N = 3$ ,  $\alpha_i = 1$ ,  $\beta = 2$ ,  $\mu_{i0} = \mu_{i1} = 0.1$ , and per-player multipliers  $r_1 = 1 < N = r_2 = 3 < r_3 = 9$ . Formula: product measure (5) (Theorem 2); Exact: values from the  $2^N$  linear system obtained using [3]. Both columns to five decimal places.

## 5 Structural consequences

The formulas (4) and (8) allow immediate structural conclusions about individual and aggregate cooperation.

**Corollary 6** (Player-specific cooperation thresholds). *Player  $i$  cooperates with probability greater than  $\frac{1}{2}$  if and only if*

$$\phi_i(\delta_i) > \frac{\frac{1}{2} - \mu_{i0}}{1 - \mu_{i0} - \mu_{i1}}.$$

*For the PGG,  $\delta_i = \alpha_i(1 - r_i/N)$ ; when  $\mu_{i0} = \mu_{i1}$  this reduces to  $r_i > N$ . A sufficient condition for  $p_C^{\text{intro}} > \frac{1}{2}$  is that the inequality holds for every  $i$ .*

*Proof.* Since  $1 - \mu_{i0} - \mu_{i1} > 0$ :

$$p_i > \frac{1}{2} \iff (1 - \mu_{i0} - \mu_{i1})\phi_i(\delta_i) > \frac{1}{2} - \mu_{i0} \iff \phi_i(\delta_i) > \frac{\frac{1}{2} - \mu_{i0}}{1 - \mu_{i0} - \mu_{i1}}.$$

When  $\mu_{i0} = \mu_{i1}$ , the right-hand side equals  $\frac{1}{2}$ , and since  $\phi_i$  is strictly decreasing with  $\phi_i(0) = \frac{1}{2}$ , this is equivalent to  $\delta_i < 0$ . For the PGG,  $\delta_i = \alpha_i(1 - r_i/N) < 0$  iff  $r_i > N$ . The claim about  $p_C^{\text{intro}}$  follows by averaging. □

With symmetric mutation, a player cooperates the majority of the time precisely when their personal multiplier on the public good exceeds the group size: that is, when they extract more from the pool per unit contributed than they put in. The top centre panel of Figure 2 illustrates this, with curves at  $r > N$  lying above  $p_i = \frac{1}{2}$  and curves at  $r < N$  lying below it.

**Corollary 7** (Neutral drift). *If  $\beta_i = 0$  for all  $i$ , then  $\phi_i$  is constant and the payoff parameters  $\delta_i$  have no effect on  $p_C$ ; cooperation is determined entirely by mutation:*

$$p_i = \frac{1 + \mu_{i0} - \mu_{i1}}{2}, \quad p_C^{\text{intro}} = \frac{1}{2} + \frac{1}{2N} \sum_{i=1}^N (\mu_{i0} - \mu_{i1}).$$

*In particular,  $p_C^{\text{intro}} = \frac{1}{2}$  if and only if  $\sum_i (\mu_{i0} - \mu_{i1}) = 0$ .*

*Proof.* When  $\beta_i = 0$ , the Fermi function satisfies  $\phi_i(x) = \frac{1}{2}$  for all  $x$ . Substituting into (4):

$$p_i = \frac{1 - \mu_{i0} - \mu_{i1}}{2} + \mu_{i0} = \frac{1}{2} + \frac{\mu_{i0} - \mu_{i1}}{2} = \frac{1 + \mu_{i0} - \mu_{i1}}{2}. \quad \square$$

Players who are insensitive to payoff differences ignore the underlying game; their long-run behaviour is set entirely by their mutation bias, and group-level cooperation reduces to the average asymmetry between mutation rates. The top right panel of Figure 2 illustrates this for symmetric mutation: at  $\beta_i = 0$ ,  $p_i = \frac{1}{2}$  for every choice of  $\alpha_i$  and  $r_i$ . The top left panel shows the same from a different angle: all three curves converge to  $p_i = \frac{1}{2}$  at  $\beta_i = 0$  before diverging as sensitivity increases, with the direction of divergence set by  $r_i$  relative to  $N$ .

**Corollary 8** (Role of heterogeneity). *For the PGG, for player  $i$  with  $r_i \neq N$ :*

- (i)  $\frac{\partial p_i}{\partial \alpha_i}$  is positive when  $r_i > N$  and negative when  $r_i < N$ .
- (ii)  $\frac{\partial p_i}{\partial \beta_i}$  is positive when  $r_i > N$  and negative when  $r_i < N$ .

*Proof.* Write  $p_i = (1 - \mu_{i0} - \mu_{i1})\phi_i(\alpha_i(1 - r_i/N)) + \mu_{i0}$ .

*Part (i).* Differentiating with respect to  $\alpha_i$ :

$$\frac{\partial p_i}{\partial \alpha_i} = (1 - \mu_{i0} - \mu_{i1})(1 - r_i/N) \phi'_i(\alpha_i(1 - r_i/N)).$$

Since  $\phi'_i(x) < 0$  for all  $x$  and  $1 - \mu_{i0} - \mu_{i1} > 0$ , the sign equals  $-\text{sgn}(1 - r_i/N) = \text{sgn}(r_i/N - 1)$ , which is positive if and only if  $r_i > N$ .

*Part (ii).* Write  $g(t) = (1 + e^t)^{-1}$ , so  $\phi_i(x) = g(\beta_i x)$ . Differentiating with respect to  $\beta_i$ :

$$\frac{\partial p_i}{\partial \beta_i} = (1 - \mu_{i0} - \mu_{i1}) \alpha_i(1 - r_i/N) g'(\beta_i \alpha_i(1 - r_i/N)).$$

Since  $g' < 0$  and  $\alpha_i > 0$ , the sign equals  $-\text{sgn}(1 - r_i/N) = \text{sgn}(r_i/N - 1)$ , which is positive if and only if  $r_i > N$ .  $\square$

For a player who benefits from the public good ( $r_i > N$ ), contributing more (larger  $\alpha_i$ ) or responding more strongly to payoff differences (larger  $\beta_i$ ) raises their cooperation rate; the same changes lower it for a player who does not benefit ( $r_i < N$ ). Heterogeneity in any one of these parameters therefore translates into heterogeneity in cooperation, with the sign of the effect set by  $r_i$  relative to  $N$ . The bottom centre and right panels of Figure 2 show this directly: the heatmaps of  $\partial p_i / \partial \alpha_i$  and  $\partial p_i / \partial \beta_i$  change sign exactly at  $r_i = N$ . The bottom left panel shows that  $\partial p_i / \partial \mu$  obeys the same rule: raising the mutation rate pulls  $p_i$  towards  $\frac{1}{2}$ , lowering cooperation when  $r_i > N$  and raising it when  $r_i < N$ .

**Remark 9.** *Corollaries 6–8 show that each player’s long-run cooperation is fully determined by their individual triple  $(\alpha_i, \beta_i, r_i)$  and the group size  $N$ . Players with  $r_i > N$  cooperate more than half the time, and increasing  $\alpha_i$  or  $\beta_i$  raises their cooperation probability further; players with  $r_i < N$  show the opposite sensitivity. The parameters of other players have no effect on player  $i$ ’s marginal cooperation probability.*




Figure 2: Structural properties of  $p_i$  for a single player with  $N = 5$ ,  $\mu_{i0} = \mu_{i1} = 0.1$ . *Top left:*  $p_i$  as a function of  $\beta_i$  ( $\alpha_i = 2$ ) for  $r < N$ ,  $r = N$ , and  $r > N$ ; all curves pass through  $p_i = \frac{1}{2}$  at  $\beta_i = 0$  (Corollary 7). *Top centre:*  $p_i$  as a function of  $\alpha_i$  for two values of  $r$  and  $\beta_i$ ; curves above (below)  $\frac{1}{2}$  correspond to  $r > N$  ( $r < N$ ). *Top right:*  $p_i(\alpha_i, r_i)$  at  $\beta_i = 0$ , confirming  $p_i = \frac{1}{2}$  everywhere (Corollary 7); the dashed line marks  $r_i = N$ . *Bottom row:* Heatmaps of the partial derivatives  $\partial p_i / \partial \mu_i$ ,  $\partial p_i / \partial \alpha_i$ , and  $\partial p_i / \partial \beta_i$  over the  $(r_i, \alpha_i)$  or  $(r_i, \beta_i)$  plane ( $\beta_i = 0.5$  or  $\alpha_i = 2$  as fixed parameter). Red indicates a positive derivative; blue indicates negative. Each derivative changes sign exactly at  $r_i = N$  (dashed line), consistent with Corollary 8.

## 6 Conclusion

We have proved that introspection dynamics on any state-independent game is exactly solvable. When  $\Delta f_i$  depends only on  $a_i$ , each selection of player  $i$  places them in state  $C$  with probability  $p_i$  regardless of current state (Theorem 1), and the stationary distribution is a product measure (Theorem 2). For the heterogeneous public goods game, the linear payoff structure implies state-independence (Lemma 4), and the long-run cooperation probability has a closed form with  $N$  terms rather than  $2^N$  (Corollary 5).

The approach does not extend to games that are not state-independent. When  $\Delta f_i(\mathbf{a})$  depends on  $\mathbf{a}_{-i}$ , the per-player chains are coupled and the stationary distribution is no longer a product measure; the full  $2^N$  linear system must be solved in general. Games such as the Stag Hunt ( $M_2$  of (2)) contain payoff cross terms  $a_i a_{-i}$  that cannot be eliminated by rescaling, so no reformulation brings them within the scope of the present results.



## 7 Acknowledgements

Harry Foster’s research was supported by EPSRC grant EP/Z535126/1.

## References

- [1] Marta C. Couto, Stefano Giaimo, and Christian Hilbe. Introspection dynamics: a simple model of counterfactual learning in asymmetric games. *New Journal of Physics*, 24(6):063010, 2022.
- [2] Marta C. Couto and Saptarshi Pal. Introspection dynamics in asymmetric multiplayer games. *Dynamic Games and Applications*, 13:1215–1246, 2023.
- [3] Harry Foster and Vince Knight. hefos/ludics: v0.1.1, April 2026.
- [4] Oliver P. Hauser, Christian Hilbe, Krishnendu Chatterjee, and Martin A. Nowak. Social dilemmas among unequals. *Nature*, 572:524–527, 2019.
- [5] Martin A. Nowak. Five rules for the evolution of cooperation. *Science*, 314:1560–1563, 2006.