

3 Quadratics and factoring, division

3.1 Tues, Feb. 27: General form of quadratics

Polynomials are important in mathematics and life. It's worth thinking about why polynomials appear so often... I don't expect that you fully understand, but I do want us to spend some time thinking about how they relate important ideas about space and distance. Does this connection seem familiar? It is related to the two interpretations (graphic, algebraic) of linear equations in the form $y = mx + b$.

Why are quadratic polynomials useful? There seem to be two different but interacting reasons. The first is that quadratic functions of a real variable are always either convex or concave and therefore have a unique maximum or minimum. The second is that quadratic functions are intimately related to bilinear forms and therefore can be accessed using linear algebra. —Paul Siegel, *Mathoverflow*

Polynomials have the form:

$$a_n x^n + \dots + a_3 x^3 + a_2 x^2 + a_1 x + a_0 \quad (27)$$

Where the “poly” means many and the “nomial” refers to terms... or things added to each other. A polynomial with only one term is called a monomial, one with two is called a binomial... and the highest order (or power) term designates whether it is quadratic, cubic, etc. The behavior of a polynomial is largely determined by its first term. Why is this? Think about properties of exponents. Is there a way we could say that x^5 is more “powerful” than x ? This gives us the “leading coefficient test”. And tells us how to graph polynomials. (Weird behavior often happens between the zeros, not outside of them). We begin investigating polynomials by thinking about quadratics. The standard form of a quadratic is:

$$f(x) = a(x - h)^2 + k \quad (28)$$

How could we derive this form from (12)? In this form, (h, k) will be the vertex. What does f look like if $a > 0$? If $a < 0$? Will the vertex be a min or max? For $ax^2 + bx + c = 0$, where $a, b, c \in \mathbb{R}$ and $a \neq 0$, $h = -\frac{b}{2a}$ and $k = f(h) = f(-\frac{b}{2a})$.

Exercise 8. Write $f(x) = 2x^2 + 8x + 7$ in standard form by factoring around x , then working on the $x^2 + x$ term and inventing a constant value to make the factorization work... then adding that value to k .