
Notes on Simplicial Neural Networks for Path Complexes

Heitor Baldo*

University of São Paulo, Brazil

Institute for Globally Distributed Open Research and Education (IGDORE)

Abstract

Simplicial neural networks (SNNs) generalize graph neural networks by performing message passing along simplices arranged in a simplicial complex. Here, we present a brief note on a straightforward extension of SNNs to path complexes.

1 Introduction

Topological deep learning [7] extends the graph neural networks (GNNs) paradigm to higher-order relational structures, including simplicial complexes, cellular complexes, and hypergraphs. In particular, simplicial neural networks (SNNs) [1, 2, 3] associate a feature vector with each k -simplex and perform message passing through the simplicial Hodge Laplacian, enabling information exchange across different simplex dimensions. However, simplicial complexes are still limited structures since every face of a simplex must be present in the complex. This assumption is often unsuitable for directed relational data, such as causal networks, where only selected directed paths are meaningful. To model such data, a more general higher-order structure is required. Path complexes [4, 5] provide such a framework by replacing simplices with directed paths as the fundamental building blocks of the complex.

2 Path Complexes and Path Homologies

Throughout this text, $G = (V, E)$ denotes a finite digraph with vertex set V and arc set $E \subseteq V \times V$ (no self-loops), and $\mathbb{K} = \mathbb{R}$. The primary bibliography for this part is [4, 5].

Definition 2.1. *Let V be a finite non-empty set. For any integer $p \geq 0$, an elementary p -path on V is any sequence $i_0 i_1 \cdots i_p$ of $p + 1$ elements of V (repetitions are permitted a priori). For $p = -1$ the elementary p -path is the empty path, denoted e . Its length is p and it has $p + 1$ vertices.*

Unlike a simplex, vertices in an elementary path may repeat (if cycles exist in G), and the ordering is essential.

Definition 2.2. *A p -path over $\mathbb{K}$ is any formal $\mathbb{K}$ -linear combination of elementary p -paths. We denote $\Lambda_p = \Lambda_p(V)$ the $\mathbb{K}$ -linear space of all p -paths. For $i_0 i_1 \cdots i_p$, the corresponding basis element of Λ_p is written $e_{i_0 \dots i_p}$. Thus, any p -path $u \in \Lambda_p$ has the form*

$$u = \sum u^{i_0 \dots i_p} e_{i_0 \dots i_p},$$

*Email address: heitor.baldo@igdore.org

where $u^{i_0 \dots i_p} \in \mathbb{K}$.

Example 2.3. $\Lambda_0 = \text{Span}\{e_i : i \in V\}$, $\Lambda_1 = \text{Span}\{e_{ij} : i, j \in V\}$, $\Lambda_2 = \text{Span}\{e_{ijk} : i, j, k \in V\}$.

Moreover, we say that an elementary path $e_{i_0 \dots i_p}$ is *regular* if $i_k \neq i_{k+1}$ for all k , and *irregular* otherwise (some *consecutive* repeat).

Definition 2.4. For any $p \geq 1$, we define the p -th boundary operator $\partial_p : \Lambda_p \rightarrow \Lambda_{p-1}$ by

$$\partial_p e_{i_0 \dots i_p} = \sum_{q=0}^p (-1)^q e_{i_0 \dots \hat{i}_q \dots i_p}.$$

For instance, $\partial_1 e_{01} = e_1 - e_0$ and $\partial_2 e_{012} = e_{12} - e_{02} + e_{01}$.

Definition 2.5. An elementary p -path $i_0 \dots i_p$ on V is called *allowed* if $i_k \rightarrow i_{k+1}$ for any $k = 0, \dots, p-1$ and *non-allowed* otherwise.

Definition 2.6. Let $\mathcal{A}_p$ the $\mathbb{K}$ -linear space spanned by allowed elementary p -paths:

$$\mathcal{A}_p = \text{Span}\{e_{i_0 \dots i_p} : i_0 \dots i_p \text{ is allowed}\},$$

and consider the following subspace of $\mathcal{A}_p$:

$$\Omega_p = \Omega_p(G) := \{u \in \mathcal{A}_p : \partial u \in \mathcal{A}_{p-1}\}.$$

The elements of Ω_p are called ∂ -invariant p -paths.

Hence, we obtain a chain complex of ∂ -invariant paths, $\Omega_\bullet = \Omega_\bullet(G)$:

$$0 \leftarrow \Omega_0 \xleftarrow{\partial} \Omega_1 \xleftarrow{\partial} \dots \xleftarrow{\partial} \Omega_{p-1} \xleftarrow{\partial} \Omega_p \xleftarrow{\partial} \dots$$

Example: A directed triangle is a sequence of three vertices a, b, c such that $a \rightarrow b \rightarrow c$, $a \rightarrow c$. A directed triangle determines a 2-path $e_{abc} \in \Omega_2$ because $e_{abc} \in \mathcal{A}_2$ and $\partial e_{abc} = e_{bc} - e_{ac} + e_{ab} \in \mathcal{A}_1$.

Moreover, we have $\partial^2 = 0$ and $\partial \Omega_p \subseteq \Omega_{p-1}$.

Definition 2.7. A path complex over a set V is a non-empty collection $\mathcal{P}$ of (regular) elementary paths on V with the following property:

- for any $n \geq 0$, if $i_0 \dots i_n \in \mathcal{P}$ then also the truncated paths $i_0 \dots i_{n-1}$ and $i_1 \dots i_n$ belong to $\mathcal{P}$.

Definition 2.8. Given a digraph $G = (V, E)$, $E \subseteq \{(i, j) : i \neq j\}$ (no loops), its path complex $\mathcal{P}(G)$ consists of all regular paths $i_0 \dots i_p$ that are walks: $(i_{k-1}, i_k) \in E$ for every k .

Note that deleting the first or last vertex of a walk is still a walk, so $\mathcal{P}(G)$ is a path complex. Hence every digraph has a path homology.

Definition 2.9. The path homology of $\mathcal{P}(G)$ is the homology of $(\Omega_\bullet, \partial)$:

$$H_p(G) = H_p(\mathcal{P}(G)) = \frac{\ker \partial|_{\Omega_p}}{\text{Im } \partial|_{\Omega_{p+1}}}.$$

The p -th Betti number is defined as the dimension of the p -th path homology: $\beta_p(G) = \dim H_p(G)$.

3 Combinatorial Hodge Laplacians

We can build Hodge Laplacian operators for path complexes associated with digraphs as follows [6].

Consider the standard inner product in the $\mathbb{R}$ -linear spaces Λ_k , $k \geq 0$. Since Ω_k is a subspace of Λ_k , it inherits the inner product. Let $\partial_k : \Omega_k \rightarrow \Omega_{k-1}$ be the k -th boundary operator and $\partial_k^* : \Omega_{k-1} \rightarrow \Omega_k$ be its adjoint operator. The *Hodge p -Laplacian operator*, $\Delta_k : \Omega_k \rightarrow \Omega_k$, is defined by

$$\Delta_k = \partial_{k+1} \circ \partial_{k+1}^* + \partial_k^* \circ \partial_k.$$

The matrix representation of the Hodge k -Laplacian operator is given by

$$[\Delta_k] = B_{k+1} B_{k+1}^T + B_k^T B_k,$$

where $B_k = [\partial_k]$ is the matrix representation of the operator ∂_k . Let δ be the *coboundary* operator and δ^* its adjoint operator. Since $[\partial_{k+1}]^T = [\delta_k]$ and $[\delta_k^*] = [\delta_k]^\dagger$, and if $[\delta_k] = A_k$, we can verify the following equalities: $[\delta_k^* \delta_k] = A_k^\dagger A_k$, $[\delta_{k-1} \delta_{k-1}^*] = A_{k-1} A_{k-1}^\dagger$, thus

$$[\Delta_k] = A_{k-1} A_{k-1}^\dagger + A_k^\dagger A_k. \quad (1)$$

4 Simplicial Neural Networks for Path Complexes

In this part, we propose a variant of the simplicial neural network (SNN) [3] for path complexes. This model is a higher-order extension of the graph neural networks (GNNs) paradigm.

Let's use the previously mentioned Hodge k -Laplacians (1) of a path complex $\mathcal{P}$ in terms of their matrices A_k , as they carry valuable topological information about the complex.

If $*$ denotes a convolution, a convolutional layer can be expressed by $\psi \circ (f * \phi_W)$, where ϕ_W is a function with small support parameterized by learnable weights W , and ψ is some nonlinearity and bias. Following Ebli et al. [3], we define the Fourier transform of real p -cochains on a path complex,

$$\mathcal{F}_k(c) = (\langle c, e_1 \rangle_k, \langle c, e_2 \rangle_k, \dots, \langle c, e_{\dim \Omega_k} \rangle_k), \quad (2)$$

where e_i are the eigenvectors of the k -Laplacian $[\Delta_k]$, ordered by eigenvalues $\lambda_1 \leq \dots \leq \lambda_{\dim \Omega_k}$.

The function $\mathcal{F}_k$ is invertible since $[\Delta_k]$ is diagonalizable. Thus, for cochains $c, c' \in C^k(\mathcal{P})$ we define their convolution as the cochain $c *_k c' = \mathcal{F}_k^{-1}(\mathcal{F}_k(c) \mathcal{F}_k(c'))$ (pointwise product); we define a simplicial convolutional layer with input p -cochain c and weights W as being of the form $\psi \circ (\mathcal{F}_k^{-1}(\phi_W) *_k c)$. To ensure certain properties, we put, for a small N ,

$$\phi_W = \sum_{i=0}^N W_i \Lambda^i = \sum_{i=0}^N W_i (\lambda_1^i, \dots, \lambda_{\dim \Omega_k}^i). \quad (3)$$

Then we can write

$$\mathcal{F}_k^{-1}(\phi_W) *_k c = \sum_{i=0}^N W_i U \text{diag}(\Lambda^i) U^T c = \sum_{i=0}^N W_i (U \text{diag}(\Lambda) U^T)^i c = \sum_{i=0}^N W_i [\Delta_k]^i c. \quad (4)$$

References

- [1] C. Bodnar, F. Frasca, Y. Wang, N. Otter, G. Montufar, P. Lio, and M. Bronstein, “Weisfeiler and Lehman go topological: Message passing simplicial networks,” *ICML*, 2021.
- [2] E. Bunch, Q. You, G. Fung, and V. Singh, “Simplicial 2-complex convolutional neural nets,” *NeurIPS Workshop on TDA and Beyond*, 2020.
- [3] S. Ebli, M. Defferrard, and G. Spreemann, “Simplicial neural networks,” *NeurIPS Workshop on TDA and Beyond*, 2020.
- [4] A. Grigor’yan, Y. Lin, Y. Muranov, and S.-T. Yau, “Homologies of path complexes and digraphs,” *arXiv:1207.2834*, 2012.
- [5] A. Grigor’yan, Y. Lin, Y. Muranov, and S.-T. Yau, “Path complexes and their homologies,” *Journal of Mathematical Sciences*, vol. 248, no. 5, pp. 564–599, 2020.
- [6] A. Grigor’yan. “Advances in path homology theory of digraphs.” *Notices of the International Consortium of Chinese Mathematicians*, 10(2), 61–124, 2022.
- [7] M. Hajij *et al.*, “Topological deep learning: Going beyond graph data,” *arXiv:2206.00606*, 2023.